


Lecture Notes on Motivic Cohomology

Clay Mathematics Monographs

Volume 2

Lecture Notes on Motivic Cohomology

**Carlo Mazza
Vladimir Voevodsky
Charles Weibel**



**American Mathematical Society
Clay Mathematics Institute**

2000 *Mathematics Subject Classification*. Primary 14F42;
Secondary 19E15, 14C25, 14-01, 14F20.

For additional information and updates on this book, visit
www.ams.org/bookpages/cmim-2

Library of Congress Cataloging-in-Publication Data

Mazza, Carlo, 1974-

Lecture notes on motivic cohomology / Carlo Mazza, Vladimir Voevodsky, Charles A. Weibel.
p. cm. — (Clay mathematics monographs, ISSN 1539-6061 ; v. 2)

Includes bibliographical references and index.

ISBN-10: 0-8218-3847-4 (alk. paper)

ISBN-13: 978-0-8218-3847-1 (alk. paper)

1. Homology theory. I. Voevodsky, Vladimir. II. Weibel, Charles A., 1950- III. Title.
IV. Series.

QA612.3.M39 2006
514'.23—dc22

2006045973

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the Clay Mathematics Institute. Requests for such permission should be addressed to the Clay Mathematics Institute, One Bow Street, Cambridge, MA 02138, USA. Requests can also be made by e-mail to permissions@claymath.org.

© 2006 by the authors. All rights reserved.

Published by the American Mathematical Society, Providence, RI,
for the Clay Mathematics Institute, Cambridge, MA.

Printed in the United States of America.

⊗ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

Visit the Clay Mathematics Institute home page at <http://www.claymath.org/>

10 9 8 7 6 5 4 3 2 1 11 10 09 08 07 06

Contents

Preface	vii
Introduction	xi
Part 1. Presheaves with Transfers	1
Lecture 1. The category of finite correspondences	3
Appendix 1A. The category $Cors$	7
Lecture 2. Presheaves with transfers	13
Homotopy invariant presheaves	17
Lecture 3. Motivic cohomology	21
Lecture 4. Weight one motivic cohomology	25
Lecture 5. Relation to Milnor K -Theory	29
Part 2. Étale Motivic Theory	35
Lecture 6. Étale sheaves with transfers	37
Lecture 7. The relative Picard group and Suslin's Rigidity Theorem	47
Lecture 8. Derived tensor products	55
Appendix 8A. Tensor triangulated categories	63
Lecture 9. \mathbb{A}^1 -weak equivalence	67
Étale \mathbb{A}^1 -local complexes	71
Lecture 10. Étale motivic cohomology and algebraic singular homology	75
Part 3. Nisnevich Sheaves with Transfers	81
Lecture 11. Standard triples	83
Lecture 12. Nisnevich sheaves	89
The cdh topology	94

Lecture 13. Nisnevich sheaves with transfers	99
Cdh sheaves with transfers	105
Part 4. The Triangulated Category of Motives	107
Lecture 14. The category of motives	109
Nisnevich \mathbb{A}^1 -local complexes	111
Motives with \mathbb{Q} -coefficients	116
Lecture 15. The complex $\mathbb{Z}(n)$ and \mathbb{P}^n	119
Lecture 16. Equidimensional cycles	125
Motives with compact support	128
Part 5. Higher Chow Groups	133
Lecture 17. Higher Chow groups	135
Appendix 17A. Cycle maps	143
Lecture 18. Higher Chow groups and equidimensional cycles	149
Appendix 18A. Generic equidimensionality	155
Lecture 19. Motivic cohomology and higher Chow groups	159
Lecture 20. Geometric motives	167
Part 6. Zariski Sheaves with Transfers	173
Lecture 21. Covering morphisms of triples	175
Lecture 22. Zariski sheaves with transfers	183
Lecture 23. Contractions	191
Lecture 24. Homotopy invariance of cohomology	197
Bibliography	203
Glossary	207
Index	211

Preface

This book was written by Carlo Mazza and Charles Weibel on the basis of the lectures on motivic cohomology which I gave at the Institute for Advanced Study in Princeton in 1999/2000.

From the point of view taken in these lectures, motivic cohomology with coefficients in an abelian group A is a family of contravariant functors

$$H^{p,q}(-, A) : Sm/k \rightarrow Ab$$

from smooth schemes over a given field k to abelian groups, indexed by integers p and q . The idea of motivic cohomology goes back to P. Deligne, A. Beilinson and S. Lichtenbaum.

Most of the known and expected properties of motivic cohomology (predicted in [ABS87] and [Lic84]) can be divided into two families. The first family concerns properties of motivic cohomology itself – there are theorems about homotopy invariance, Mayer-Vietoris and Gysin long exact sequences, projective bundles, etc. This family also contains conjectures such as the Beilinson-Soulé vanishing conjecture ($H^{p,q} = 0$ for $p < 0$) and the Beilinson-Lichtenbaum conjecture, which can be interpreted as a partial étale descent property for motivic cohomology. The second family of properties relates motivic cohomology to other known invariants of algebraic varieties and rings. The power of motivic cohomology as a tool for proving results in algebra and algebraic geometry lies in the interaction of the results in these two families; applying general theorems of motivic cohomology to the specific cases of classical invariants, one gets new results about these invariants.

The idea of these lectures was to define motivic cohomology and to give careful proofs for the elementary results in the second family. In this sense they are complementary to the study of [VSF00], where the emphasis is on the properties of motivic cohomology itself. The structure of the proofs forces us to deal with the main properties of motivic cohomology as well (such as homotopy invariance). As a result, these lectures cover a considerable portion of the material of [VSF00], but from a different point of view.

One can distinguish the following “elementary” comparison results for motivic cohomology. Unless otherwise specified, all schemes below are assumed to be smooth or (in the case of local or semilocal schemes) limits of smooth schemes.

(1) $H^{p,q}(X, A) = 0$ for $q < 0$, and for a connected X one has

$$H^{p,0}(X, A) = \begin{cases} A & \text{for } p = 0 \\ 0 & \text{for } p \neq 0; \end{cases}$$

(2) one has

$$H^{p,1}(X, \mathbb{Z}) = \begin{cases} \mathcal{O}^*(X) & \text{for } p = 1 \\ \text{Pic}(X) & \text{for } p = 2 \\ 0 & \text{for } p \neq 1, 2; \end{cases}$$

(3) for a field k , one has $H^{p,p}(\text{Spec}(k), A) = K_p^M(k) \otimes A$ where $K_p^M(k)$ is the p -th Milnor K -group of k (see [Mil70]);

(4) for a strictly Hensel local scheme S over k and an integer n prime to $\text{char}(k)$, one has

$$H^{p,q}(S, \mathbb{Z}/n) = \begin{cases} \mu_n^{\otimes q}(S) & \text{for } p = 0 \\ 0 & \text{for } p \neq 0 \end{cases}$$

where $\mu_n(S)$ is the groups of n -th roots of unity in S ;

(5) one has $H^{p,q}(X, A) = CH^q(X, 2q - p; A)$. Here $CH^i(X, j; A)$ denotes the higher Chow groups of X introduced by S. Bloch in [Blo86], [Blo94]. In particular,

$$H^{2q,q}(X, A) = CH^q(X) \otimes A,$$

where $CH^q(X)$ is the classical Chow group of cycles of codimension q modulo rational equivalence.

The isomorphism between motivic cohomology and higher Chow groups leads to connections between motivic cohomology and algebraic K -theory, but we do not discuss these connections in the present lectures. See [Blo94], [BL94], [FS02], [Lev98] and [SV00].

Deeper comparison results include the theorem of M. Levine comparing $CH^i(X, j; \mathbb{Q})$ with the graded pieces of the gamma filtration in $K_*(X) \otimes \mathbb{Q}$ [Lev94], and the construction of the spectral sequence relating motivic cohomology and algebraic K -theory for arbitrary coefficients in [BL94] and [FS02].

The lectures in this book may be divided into two parts, corresponding to the fall and spring terms. The fall term lectures contain the definition of motivic cohomology and the proofs for all of the comparison results listed above except the last one. The spring term lectures include more advanced results in the theory of sheaves with transfers and the proof of the final comparison result (5).

The definition of motivic cohomology used here goes back to the work of Andrei Suslin in about 1985. As I understand it, when he came up with this definition he was able to prove the first three of the comparison results stated above. In particular, the proof of comparison (3) between motivic cohomology and Milnor's K -groups given in these lectures is exactly Suslin's original proof. The proofs of the last two comparison results, (4) and (5), are also based on results of Suslin. Suslin's formulation of the Rigidity Theorem ([Sus83]; see theorem 7.20) is a key

result needed for the proof of (4), and Suslin’s moving lemma (theorem 18A.1 below) is a key result needed for the proof of (5).

It took ten years and two main new ideas to finish the proofs of the comparisons (4) and (5). The first one, which originated in the context of the *qfh*-topology and was later transferred to sheaves with transfers (definition 2.1), is that the sheaf of finite cycles $\mathbb{Z}_{tr}(X)$ is the *free* object generated by X . This idea led to a group of results, the most important of which is lemma 6.23. The second idea, which is the main result of [CohTh], is represented here by theorem 13.8. Taken together they allow one to efficiently do homotopy theory in the category of sheaves with transfers.

A considerable part of the first half of the lectures is occupied by the proof of (4). Instead of stating it in the form used above, we prove a more detailed theorem. For a given weight q , the motivic cohomology groups $H^{p,q}(X,A)$ are defined as the hypercohomology (in the Zariski topology) of X with coefficients in a complex of sheaves $A(q)|_{X_{Zar}}$. This complex is the restriction to the small Zariski site of X (i.e., the category of open subsets of X) of a complex $A(q)$ defined on the site of all smooth schemes over k with the Zariski and even the étale topology. Restricting $A(q)$ to the small étale site of X , we may consider the étale version of motivic cohomology,

$$H_L^{p,q}(X,A) := \mathbb{H}_{\acute{e}t}^p(X,A(q)|_{X_{\acute{e}t}}).$$

The subscript L is in honor of Steve Lichtenbaum, who first envisioned this construction in [Lic94].

Theorem 10.2 asserts that the étale motivic cohomology of any X with coefficients in $\mathbb{Z}/n(q)$ where n is prime to $\text{char}(k)$ are isomorphic to $H_{\acute{e}t}^p(X, \mu_n^{\otimes q})$. This implies comparison result (4), since the Zariski and the étale motivic cohomology of a strictly Hensel local scheme X agree. There should also be an analog of (4) for the case of \mathbb{Z}/ℓ^r coefficients where $\ell = \text{char}(k)$, involving the logarithmic de Rham-Witt sheaves $v_r^q[-q]$, but I do not know much about it. We refer the reader to [GL00] for more information.

Vladimir Voevodsky
 Institute for Advanced Study
 May 2001

Introduction

This book is divided into six main parts. The first part (lectures 1–5) presents the definitions and the first three comparison results. The second part (lectures 6–10) presents the étale version of the theory, focussing on coefficients, $1/m \in k$. As Suslin’s Rigidity Theorem 7.20 demonstrates, a key role is played by locally constant étale sheaves such as $\mu_m^{\otimes i}$, which are quasi-isomorphic to the motivic $\mathbb{Z}/m(i)$ by theorem 10.3. The tensor triangulated category $\mathbf{DM}_{\acute{e}t}^-(k, \mathbb{Z}/m)$ of étale motives is constructed in lecture 9 and shown to be equivalent to the derived category of discrete \mathbb{Z}/m -modules over the Galois group $G = \text{Gal}(k_{\text{sep}}/k)$ in theorem 9.35.

The first main goal of the lecture notes, carried out in lectures 11–16, is to introduce the tensor triangulated category $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k, R)$ of effective motives and its subcategory of effective geometric motives $\mathbf{DM}_{\text{gm}}^{\text{eff}}$. The motive $M(X)$ of a scheme X is an object of $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k, R)$, and belongs to $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ if X is smooth. This requires an understanding of the cohomological properties of sheaves associated with homotopy invariant presheaves with transfers for the Zariski, Nisnevich and cdh topologies. This is addressed in the third part (lectures 11–13). Lecture 11 introduces the technical notion of a standard triple, and uses it to prove that homotopy invariant presheaves with transfers satisfy a Zariski purity property. Lecture 12 introduces the Nisnevich and cdh topologies, and lecture 13 considers Nisnevich sheaves with transfers and their associated cdh sheaves.

A crucial role in this development is played by theorem 13.8: if F is a homotopy invariant presheaf with transfers, and k is a perfect field, then the associated Nisnevich sheaf F_{Nis} is homotopy invariant, and so is its cohomology. For reasons of exposition, the proof of this result is postponed and occupies lectures 21 to 24.

In the fourth part (lectures 14–16) we introduce the categories $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k, R)$ and $\mathbf{DM}_{\text{gm}}^{\text{eff}}$. The main properties of these categories — homotopy, Mayer-Vietoris, projective bundle decomposition, blow-up triangles, Gysin sequence, the Cancellation Theorem, and the connection with Chow motives — are summarized in 14.5. We also show (in 15.9) that the product on motivic cohomology (defined in 3.12) is graded-commutative and in agreement (for coefficients \mathbb{Q}) with the étale theory presented in lectures 9 and 10 (see 14.30).

Lecture 16 introduces equidimensional algebraic cycles. These are used to construct the Suslin-Friedlander motivic complexes $\mathbb{Z}^{SF}(i)$, which are quasi-isomorphic to the motivic complexes $\mathbb{Z}(i)$; this requires the field to be perfect (see 16.7). They are also used to define motives with compact support $M^c(X)$. The basic

theory with compact support complements the theory presented in lecture 14; this requires the field to admit resolution of singularities. This lecture concludes with the use of Friedlander-Voevodsky duality (see 16.24) to establish the Cancellation Theorem 16.25; this lets us embed effective motives into the triangulated category of all motives.

The second main goal of this book is to establish the final comparison (theorem 19.1) with Bloch’s higher Chow groups: for any smooth separated scheme X over a perfect field k , we have

$$H^{p,q}(X, \mathbb{Z}) \cong CH^q(X, 2q - p).$$

This is carried out in the fifth part (lectures 17–19). In lecture 17, we introduce Bloch’s higher Chow groups and show (in 17.21) that they are presheaves with transfers over any field. Suslin’s comparison (18.3) of higher Chow groups with equidimensional cycle groups over any affine scheme is given in lecture 18, and the link between equidimensional cycle groups and motivic cohomology is given in lecture 19.

We briefly revisit the triangulated category \mathbf{DM}_{gm} of geometric motives in lecture 20. We work over a perfect field which admits resolution of singularities. First we embed Grothendieck’s classic category of Chow motives as a full subcategory. We then construct the dual of any geometric motive and use it to define internal Hom objects $\underline{Hom}(X, Y)$. The lecture culminates in theorem 20.17, which states that this structure makes \mathbf{DM}_{gm} a rigid tensor category.

The final part (lectures 21–24) is dedicated to the proof of theorem 13.8. Using technical results from lecture 21, we prove (in 22.3) that F_{Nis} is homotopy invariant. The proof that its cohomology is homotopy invariant (24.1) is given in lecture 24. We conclude with a proof that the sheaf F_{Nis} admits a “Gersten” resolution.

During the writing of the book, we received many suggestions and comments from the mathematical community. One popular suggestion was that we include some of the more well known and useful properties of motives that were missing from the original lectures, in order to make the exposition of the theory more complete. For this reason, a substantial amount of material has been added to lectures 12–14, 16 and 20. Another suggestion was that we warn the reader that the exercises vary in difficulty and content, from the concrete to the abstract; some are learning exercises and some augment the ideas presented in the text.

In Figure 1 we give a rough bird’s eye view of the structure of the book and how the various lectures depend upon each other. Lectures 1 and 2 are missing from the figure because they are prerequisites for all other lectures. We split lecture 13 into two parts to clarify that the results in the second half of the lecture crucially depend on theorem 13.8, which is proven in lecture 24. The dependency chart (and this Introduction) should serve as a guide to the reader.




FIGURE 1. Dependency graph of the lectures

Acknowledgements

The authors are deeply indebted to the Institute for Advanced Study, the Clay Mathematics Institute and Rutgers University for the support provided by these institutions during the writing of this book. In addition, we are grateful to Thomas Geisser, who carefully proofread an earlier version and provided us with his lecture notes on this subject, which were used as an outline for the additions in lectures 12, 16 and 20.

In addition, the authors were supported on numerous grants during the writing phase (2000–2005) of this book. This includes support from the NSF, NSA, INDAM and the institutions named above.

Carlo Mazza
Vladimir Voevodsky
Charles A. Weibel
December 2005

Part 1

Presheaves with Transfers

LECTURE 1

The category of finite correspondences

In this lecture we shall define the additive category Cor_k of finite correspondences over a field k . The objects of Cor_k will be the smooth separated schemes (of finite type) over k . The morphisms in Cor_k from X to Y will be the finite correspondences, which are special kinds of cycles in $X \times Y$. Composition is defined so that Cor_k contains the category Sm/k of smooth separated schemes over k .

By convention, all schemes will be separated, and defined over k . Although smooth schemes always have finite type over k [EGA4, 17.3.1], we will sometimes refer to local and even semilocal schemes as being smooth; by this we mean that they are the local (resp., semilocal) schemes associated to points on a smooth scheme.

Our point of view will be that a cycle in a scheme T is a formal \mathbb{Z} -linear combination of irreducible closed subsets of T . Each irreducible closed subset W is the support of its associated integral subscheme \tilde{W} so W and \tilde{W} determine each other. Thus we can ascribe some algebraic properties to W . We say that W is **finite** along a morphism $T \rightarrow S$ if the restriction $\tilde{W} \rightarrow S$ is a finite morphism. A cycle $\sum n_i W_i$ is said to be finite along a morphism if each W_i is finite.

DEFINITION 1.1. If X is a smooth connected scheme over k , and Y is any (separated) scheme over k , an **elementary correspondence** from X to Y is an irreducible closed subset W of $X \times Y$ whose associated integral subscheme is finite and surjective over X . By an elementary correspondence from a non-connected scheme X to Y , we mean an elementary correspondence from a connected component of X to Y .

The group $Cor(X, Y)$ is the free abelian group generated by the elementary correspondences from X to Y . The elements of $Cor(X, Y)$ are called **finite correspondences**.

If X is not connected and $X = \coprod X_i$ is the decomposition into its connected components, our definition implies that $Cor(X, Y) = \bigoplus_i Cor(X_i, Y)$.

EXAMPLE 1.2. Let $f : X \rightarrow Y$ be a morphism in Sm/k . If X is connected, the graph Γ_f of f is an elementary correspondence from X to Y . If X is not connected, the sum of the components of Γ_f is a finite correspondence from X to Y . Indeed the projection $\Gamma_f \rightarrow X$ is an isomorphism, and Γ_f is closed because Y is separated over k .

The graph Γ_1 of the identity on X is the support of the diagonal $\Delta(X) \subset X \times X$. We write id_X for the finite correspondence Γ_1 from X to itself. It is the identity element of $Cor(X, X)$ for the composition product. Note that id_X is an elementary correspondence when X is integral.

If X is connected, Y is smooth and $f : X \rightarrow Y$ is finite and surjective, the transpose of Γ_f in $Y \times X$ is a finite correspondence from Y to X . This is a useful construction; see exercise 1.11 below for one application.

CONSTRUCTION 1.3. *Every subscheme Z of $X \times Y$ which is finite and surjective over X determines a finite correspondence $[Z]$ from X to Y .*

PROOF. If Z is integral then its support $[Z]$ is by definition an elementary correspondence. In general we associate to Z the finite correspondence $\sum n_i W_i$, where the W_i are the irreducible components of the support of Z which are surjective over a component of X and n_i is the geometric multiplicity of W_i in Z , i.e., the length of the local ring of Z at W_i (see [Ser65] or [Ful84]). \square

We now define an associative and bilinear composition for finite correspondences between smooth schemes. For this, it suffices to define the composition $W \circ V$ of elementary correspondences $V \in Cor(X, Y)$ and $W \in Cor(Y, Z)$. Our definition will use the push-forward of a finite cycle.

Let $p : T \rightarrow S$ be any morphism. If W is an irreducible closed subset of T finite along p , the image $V = p(W)$ is a closed irreducible subset of S and $d = [k(W) : k(V)]$ is finite. In this case we define the **push-forward** of the cycle W along p to be the cycle $p_* W = d \cdot V$; see [Ful84]. By additivity we may define the push-forward of any cycle which is finite along p .

LEMMA 1.4. *Suppose that $f : T \rightarrow T'$ is a morphism of separated schemes of finite type over a Noetherian base S . Let W be an irreducible closed subset of T which is finite over S . Then $f(W)$ is closed and irreducible in T' and finite over S . If W is finite and surjective over S , then so is $f(W)$.*

PROOF. By Ex.II.4.4 of [Har77], $f(W)$ is closed in T' and proper over S . Since $f(W)$ has finite fibers over S , it is finite over S by [EGA3, 4.4.2]. If $W \rightarrow S$ is surjective, so is $f(W) \rightarrow S$. \square

Given elementary correspondences $V \in Cor(X, Y)$ and $W \in Cor(Y, Z)$, form the intersection product $[T] = (V \times Z) \cdot (X \times W)$ of the corresponding cycles in $X \times Y \times Z$. (The intersection product is defined in [Ser65] and [Ful84]; see 17A.1.)

The composition $W \circ V$ of V and W is defined to be the push-forward of the finite correspondence $[T]$, along the projection $p : X \times Y \times Z \rightarrow X \times Z$; see [Ful84]. By lemma 1.7 below, the cycle $[T]$ is finite over $X \times Z$. Thus the push-forward $p_* [T]$ is defined; it is a finite correspondence from X to Z by lemma 1.4.

We can easily check that id_X is the identity of $Cor(X, X)$, and that the composition of finite correspondences is associative and bilinear (see [Man68] and [Ful84, 16.1]).

DEFINITION 1.5. Let Cor_k be the category whose objects are the smooth separated schemes of finite type over k and whose morphisms from X to Y are elements of $Cor(X, Y)$. It follows from the above remarks that Cor_k is an additive category with \emptyset as the zero object, and disjoint union as coproduct.

LEMMA 1.6. *Let Z be an integral scheme, finite and surjective over a normal scheme S . Then for every morphism $T \rightarrow S$ with T connected, every component of $T \times_S Z$ is finite and surjective over T .*

PROOF. See [EGA4, 14.4.4]. \square

Recall that two irreducible closed subsets Z_1 and Z_2 of a smooth scheme are said to intersect **properly** if $Z_1 \cap Z_2 = \emptyset$ or $\text{codim}(Z_1 \cap Z_2) = \text{codim} Z_1 + \text{codim} Z_2$.

LEMMA 1.7. *Let $V \subset X \times Y$ and $W \subset Y \times Z$ be irreducible closed subsets which are finite and surjective over X and Y respectively. Then $V \times Z$ and $X \times W$ intersect properly, and each component of the push-forward of the cycle $[T]$ of $T = (V \times Z) \cap (X \times W)$ is finite and surjective over X .*

PROOF. Let \tilde{V} and \tilde{W} be the underlying integral subschemes associated to V and W respectively. Without loss of generality we can suppose both X and Y connected. We form the pullback of \tilde{V} and \tilde{W} .

$$\begin{array}{ccccc}
 \tilde{V} \times_Y \tilde{W} & \longrightarrow & \tilde{W} & \longrightarrow & Z \\
 \downarrow & & \downarrow \text{f.surj.} & & \\
 \tilde{V} & \longrightarrow & Y & & \\
 \downarrow \text{f.surj.} & & & & \\
 X & & & &
 \end{array}$$

By 1.6, each component of $\tilde{V} \times_Y \tilde{W}$ is finite and surjective over \tilde{V} and therefore over X too. The image T of the evident map $\tilde{V} \times_Y \tilde{W} \rightarrow X \times Y \times Z$ is the intersection of $\tilde{V} \times Z$ and $X \times \tilde{W}$. Thus each irreducible component T_i of T is the image of an irreducible component of $\tilde{V} \times_Y \tilde{W}$. By 1.4, we know that each T_i is finite and surjective over X . Therefore $\dim T_i = \dim X$ for all i , i.e., $\tilde{V} \times Z$ and $X \times \tilde{W}$ intersect properly.

Let $p(T_i)$ denote the image of T_i under the map $p : X \times Y \times Z \rightarrow X \times Z$. By lemma 1.4, each $p(T_i)$ is an irreducible closed subscheme of $X \times Z$ which is finite and surjective over X . Since the components of $p_*[T]$ are the supports of the $p(T_i)$, we are done. \square

REMARK 1.8. It is possible to extend the definition of finite correspondences to correspondences between singular schemes. This uses the category Cor_S , where S is a Noetherian scheme; see [RelCh]. Since we use only smooth schemes in these lectures, we describe this more general definition in the appendix of this lecture.

The additive category Cor_k is closely related to the category Sm/k of smooth schemes over k . Indeed, these categories have the same objects, and it is a routine computation (exercise!) to check that $\Gamma_g \circ \Gamma_f$ equals $\Gamma_{g \circ f}$. That is, there is a faithful functor $Sm/k \rightarrow Cor_k$, defined by:

$$X \mapsto X \quad (f : X \rightarrow Y) \mapsto \Gamma_f.$$

The tensor product is another important feature of the category Cor_k .

DEFINITION 1.9. If X and Y are two objects in Cor_k , their **tensor product** $X \otimes Y$ is defined to be the product of the underlying schemes over k :

$$X \otimes Y = X \times Y.$$

If V and W are elementary correspondences from X to X' and from Y to Y' , then the cycle associated to the subscheme $V \times W$ by 1.3 gives a finite correspondence from $X \otimes Y$ to $X' \otimes Y'$.

It is easy to verify that \otimes makes Cor_k a symmetric monoidal category (see [Mac71]).

EXERCISE 1.10. If $S = \text{Spec } k$ then $Cor_k(S, X)$ is the group of zero-cycles in X . If W is a finite correspondence from \mathbb{A}^1 to X , and $s, t : \text{Spec } k \rightarrow \mathbb{A}^1$ are k -points, show that the zero-cycles $W \circ \Gamma_s$ and $W \circ \Gamma_t$ are rationally equivalent (cf. [Ful84, 1.6]).

EXERCISE 1.11. Let x be a closed point on X , considered as a correspondence from $S = \text{Spec}(k)$ to X . Show that the composition $S \rightarrow X \rightarrow S$ is multiplication by the degree $[k(x) : k]$, and that $X \rightarrow S \rightarrow X$ is given by $X \times x \subset X \times X$.

Let L/k be a finite Galois extension with Galois group G and $T = \text{Spec}(L)$. Prove that $Cor_k(T, T) \cong \mathbb{Z}[G]$ and that $T \rightarrow S \rightarrow T$ is $\sum_{g \in G} g \in \mathbb{Z}[G]$. Then show that $Cor_k(S, Y) \cong Cor_k(T, Y)^G$ for every Y .

EXERCISE 1.12. If $k \subset F$ is a field extension, there is an additive functor $Cor_k \rightarrow Cor_F$ sending X to X_F . If F is finite and separable over k , there is an additive functor $Cor_F \rightarrow Cor_k$ sending U to U . These are adjoint: if U is smooth over F and X is smooth over k , there is a canonical identification:

$$Cor_F(U, X_F) = Cor_k(U, X).$$

EXERCISE 1.13. (a) Let F be a field extension of k and X and Y two smooth schemes over k . Writing X_F for $X \times_{\text{Spec } k} \text{Spec } F$ and so on, show that $Cor_F(X_F, Y_F)$ is the limit of the $Cor_E(X_E, Y_E)$ as E ranges over all finitely generated field extensions of k contained in F .

(b) Let $X \rightarrow S \rightarrow \text{Spec}(k)$ be smooth morphisms, with S connected, and let F denote the function field of S . For every smooth scheme Y over k , show that $Cor_F(X \times_S \text{Spec } F, Y \times_k \text{Spec } F)$ is the direct limit of the $Cor_k(X \times_S U, Y)$ as U ranges over all nonempty open subschemes of S . In the special case $X = S$, this shows that $Cor_F(\text{Spec } F, Y \times_k \text{Spec } F) = \varinjlim Cor_k(U, Y)$.

(c) Show that (a) and (b) remain valid if Y is any scheme over k , using definition 1.1 of $Cor_k(X, Y)$.

APPENDIX 1A

The category Cor_S

It is possible to generalize the notion of finite correspondence to construct a category Cor_S , associated to any Noetherian scheme S ; see [RelCh]. The objects of this category are the schemes of finite type over S ; the morphisms are the elements of an abelian group $Cor_S(X, Y)$ whose elements are the cycles W on $X \times_S Y$ which are “universally integral relative to X ”, and each of whose components are finite and surjective over X .

Universally integral cycles are defined in 1A.9 below as those cycles for which the pullback is always defined, and has integer coefficients. This condition is needed because, in order to compose an elementary correspondence V in $Cor_S(X, Y)$ with a correspondence W in $Cor_S(Y, Z)$, we must form the pullback W_V of W along $V \rightarrow Y$ to get a cycle on $V \times_S Z \subset X \times_S Y \times_S Z$ (see 1A.11).

Relabeling, we are reduced to the following basic setup for pulling back cycles. We are given a cycle W on X , a structure map $X \rightarrow S$ and a map $V \rightarrow S$. The problem is to define a pullback cycle W_V on $X \times_S V$ in a natural way. This is easy if V is flat over S (see [Ful84, 1.7]), but in general the problem is quite difficult even for $V = \text{Spec } K$.

$$\begin{array}{ccccc}
 W_V \subset & X \times_S V & \longrightarrow & V & \\
 & \downarrow & & \downarrow & \\
 W \subset & X & \longrightarrow & S &
 \end{array}$$

The general pullback is modelled on the pullback of flat cycles. If W is an irreducible cycle on X which is flat over S , we define the pullback $s^*(W)$ along $s : \text{Spec}(K) \rightarrow S$ to be the cycle on X_s defined by $W_s = W \times_S \text{Spec}(K)$.

EXAMPLE 1A.1. Let W be an irreducible cycle on X . By “platification” [RG71, 5.2], there is a proper birational map $T \rightarrow S$ such that the proper transform \tilde{W} of W is flat over T . Given a point $s_0 : \text{Spec}(k_0) \rightarrow S$, choose a finite field extension k_1 of k_0 such that $s_1 : \text{Spec}(k_1) \rightarrow S$ has a lift $t : \text{Spec}(k_1) \rightarrow T$; then the flat pullback $t^*(\tilde{W})$ is a candidate for the pullback $s_1^*(W)$.

There are two problems with this candidate: it may depend upon the choice of T and t (as in example 1A.4), and if $k_0 \neq k_1$ we need to descend from the cycle

$t^*(\tilde{W})$ on X_{k_1} to a cycle on X_{k_0} (as in example 1A.7).

$$\begin{array}{ccccc}
 \text{Spec } k_1 & \xrightarrow{t} & T & \xleftarrow{\text{flat}} & \tilde{W} \\
 \downarrow & \nearrow s_1 & \uparrow & & \\
 \text{Spec } k_0 & \xrightarrow{s_0} & S & \xleftarrow{} & W
 \end{array}$$

One way to attack the first problem is to restrict our attention to “relative cycles,” defined in 1A.5 using the notion of pullback along a fat point of S . This approach was introduced in [RelCh], using discrete valuation rings (DVRs). Recall that if K is a field, a K -point of S (or point) is a morphism $\text{Spec } K \rightarrow S$.

DEFINITION 1A.2. A **fat point** \underline{s} of S is a DVR D , a field K and morphisms

$$\text{Spec } K \xrightarrow{s_0} \text{Spec } D \xrightarrow{s_1} S,$$

so that the closed point of $\text{Spec } K$ goes to the closed point of $\text{Spec } D$ and the generic point $\text{Spec } F$ of $\text{Spec } D$ goes to a generic point of S . We say that the fat point $\underline{s} = (s_0, s_1)$ lies over the underlying K -point $\text{Spec } K \rightarrow S$.

Every point s in a Noetherian scheme S has a fat point lying over it in the sense that there is a field extension $k(s) \subset K$ and a fat point over $\text{Spec } K \rightarrow S$. That is, if s lies over a generic point s' of S , then there is a DVR D and a map $\text{Spec } D \rightarrow S$ sending the closed point (resp., generic point) of $\text{Spec } D$ to s (resp., to s'); see [EGA1, 0_I.6.5.8] or [Har77, II.4.11]. The following trick now lets us take the pullback of cycles to $\text{Spec } D$.

THEOREM 1A.3. *Let D be a DVR with field of fractions F . If X is a scheme of finite type over D and W_F is closed in the generic fiber X_F then there exists a unique closed subscheme W_D of W_F in X which is flat over $\text{Spec } D$.*

PROOF. Locally X has coordinate ring A , X_F has coordinate ring $A \otimes_D F$, and W_F has coordinate ring $(A \otimes_D F)/(f_1, \dots, f_n)$, where $f_i \in A$ for every $i = 1, \dots, n$. Let R_0 be $A/(f_1, \dots, f_n)$ and let R be R_0/I where I is the torsion submodule of the D -module R_0 . It is easy to see that R is independent of the choice of the f_i 's. Locally W_D is $\text{Spec } R$. \square

We can now define the pullback along a fat point \underline{s} of S over a K -point s . Given a closed subscheme W in X , we may form the (classical) flat pullback W_F along $\text{Spec } F \rightarrow S$, and consider the closed subscheme W_D of W_F flat over $\text{Spec } D$ as in 1A.3. The pullback $\underline{s}^*(W)$ of W is defined to be the cycle $[W_K]$ on X_K associated to the fiber W_K of the scheme W_D over the closed point $\text{Spec } K$ of $\text{Spec } D$.

Since the pullback $\underline{s}^*(W)$ is a cycle on $X_s = X \times_S \text{Spec}(K)$, it is a candidate for the pullback of W along s . However, two fat points over the same K -point may give two distinct candidates, as the following example shows.

EXAMPLE 1A.4. Let S be the node over a field k and X its normalization. There are two fat points over the singular point $s \in S$, corresponding to the two k -points of $X_s = \{p_0, p_1\}$. The pullbacks of $W = X$ along these fat points are $[p_0]$ and $[p_1]$, respectively.

In order to have a useful pullback, we need to get rid of the dependence on the choice of the fat point. The following definition is taken from [RelCh, 3.1.3].

DEFINITION 1A.5. Let $W = \sum n_i W_i$ be a cycle on X . We say that W is **dominant** over S if each term W_i of W is dominant over a component of S . We say that a dominant cycle W is a **relative cycle** on X over S if its pullbacks $\underline{s}^*(W)$ and $\underline{t}^*(W)$ coincide for any pair $\underline{s}, \underline{t}$ of fat points over a common K -point s . We will write $s^*(W)$ for this pullback cycle on X_s .

For example, any dominant cycle W which is flat and equidimensional over S is a relative cycle, because the pullback $\underline{s}^*(W)$ coincides with the classical pullback of a cycle along the K -point. This follows easily from the observation that since W_D is $W \times_S \text{Spec } D$, we have $W_K = W \times_S \text{Spec } K$.

We write $Cycl(X/S, r)$ for the free abelian group of the relative cycles W on X over S such that each component has dimension r over S . It turns out that every effective relative cycle in $Cycl(X/S, r)$ must be equidimensional over S ; see [RelCh, 3.1.7]. If S is normal, the following result shows that this is also a sufficient condition for being a relative cycle; it is proven in [RelCh, 3.4.2].

THEOREM 1A.6. *If S is normal or geometrically unibranch, and W is a cycle on X which is dominant equidimensional over S , then W is a relative cycle.*

The use of relative cycles solves the first problem raised in the situation of example 1A.1. Given a relative cycle W , find a proper birational map $T \rightarrow S$ as in 1A.1 so that the components of W have flat proper transforms in X_T . By the Valuative Criterion for Properness, fat points \underline{s} of S are in 1–1 correspondence with fat points \underline{t} of T . Given any pair of liftings $t', t'' : \text{Spec}(k_1) \rightarrow T$ of a k_1 -point s_1 on S , we can find fat points \underline{s}' and \underline{s}'' of S over a field extension $\eta : k_1 \rightarrow K$ whose lifts to T factor through t' and t'' . Since W is a relative cycle, we see that $\eta^* t'^*(\tilde{W}) = (\underline{s}')^*(W)$ and $\eta^* t''^*(\tilde{W}) = (\underline{s}'')^*(W)$ agree as cycles on X_K . Since η^* is an injection, we get $t'^*(\tilde{W}) = t''^*(\tilde{W})$.

$$\begin{array}{ccccc}
 & & t' & & \\
 & & \xrightarrow{\quad} & & \\
 \text{Spec } k_1 & \xrightarrow{\quad} & T & \xleftarrow{\quad} & \tilde{W} \\
 & \searrow^{s_1} & \downarrow & & \\
 & & S & \xleftarrow{\quad} & W \\
 \uparrow \eta & & \xrightarrow{\underline{s}'} & & \\
 \text{Spec } K & \xrightarrow{\quad} & S & \xleftarrow{\quad} & \\
 & & \xrightarrow{\underline{s}''} & &
 \end{array}$$

Now that we have a good definition for the pullback of a relative cycle along a k_1 -point $s_1 : \text{Spec } k_1 \rightarrow S$ which lifts to a k_1 -point t of T , we need to solve the second

problem raised in 1A.1, descending from $s_1^*(W)$ to the pullback $s^*(W)$ along any Zariski point $s : \text{Spec}(k_0) \rightarrow S$.

If k_1 is separable over k_0 , elementary Galois theory allows us to descend from the cycle $t^*(\tilde{W})$ on X_{k_1} to a cycle on X_{k_0} , which is the desired pullback $s^*(W)$. More precisely, we may assume that $i : k_0 \rightarrow k_1$ is a Galois extension with Galois group G , in which case every G -invariant cycle on X_{k_1} comes from a unique cycle on X_{k_0} . Since $t^*(\tilde{W})$ is G -invariant, there is a unique cycle, which we call $s^*(W)$, such that $t^*(\tilde{W}) = t^*(s^*(W))$.

However, it may be that the fields $k(t)$ are inseparable over $k(s)$ for every point t of T over s . To fix this, it turns out that we need to invert the characteristic p of $k(s)$ (see 1A.7 below). Fixing s and t , let K denote the maximal purely inseparable extension of $k(s)$ in $k(t)$; by the preceding paragraph, $t^*(\tilde{W})$ descends to a unique cycle Z' on X_K . Since the index $[K : k(s)]$ is a power of p , and elementary field theory shows that $[K : k(s)]Z$ comes from a unique cycle Z on X_s , we write $s^*(W)$ for the cycle $Z/[K : k(s)]$. This completes the solution of the second problem referred to in 1A.1.

$$\begin{array}{ccccc}
 & & \text{Spec } k(t) & \xrightarrow{t} & T & \longleftarrow & \tilde{W} \\
 & \swarrow \text{sep.} & \downarrow & & \downarrow & & \\
 \text{Spec } K & \xrightarrow{\text{p.insep}} & \text{Spec } k(s) & \xrightarrow{s} & S & \longleftarrow & W
 \end{array}$$

EXAMPLE 1A.7. Let K be a purely inseparable extension of k with $[K : k] = p$ and set $W = X = \text{Spec}(K[t])$. Let $S = \text{Spec } A$, where $A \subset K[t]$ is the ring of polynomials $f(t)$ where $f(0) \in k$. If $s : \text{Spec}(k) \rightarrow S$ is the origin, and the K -point s_1 of S lies over s , then using the fat point with $D = K[[t]]$ we have $s_1^*(W) = [s_1]$ on X_K . It follows that $s^*(W) = [s]/p$ as a cycle on $X_k = \text{Spec}(K)$.

Even if X is smooth and S is normal, there can be a relative cycle W for which the coefficient $1/p$ occurs in its pullbacks $\underline{s}^*(W)$. An example, due to Merkurjev, is given in Example 3.5.10 in [RelCh].

THEOREM 1A.8. *Let W be a relative cycle on X over a Noetherian scheme S . For each map $f : T \rightarrow S$, there exists a unique and well-defined relative cycle W_T of $X \times_S T$ over T , whose coefficients may lie in $\mathbb{Z}[1/p]$ in characteristic p , satisfying the following condition: for every point t of T , the pullback $t^*(W)$ to X_t agrees with the pullback $f(t)^*(W)$. The relative cycle W_T is called the **pullback** of W .*

PROOF. For each generic point t of T , consider the pullback cycles $t^*(W) = \sum n_{t,i} Z'_{t,i}$ on X_t constructed above. Let $Z_{t,i}$ denote the closure of $Z'_{t,i}$ in $X \times_S T$. Then $W_T = \sum_{t,i} n_{t,i} Z_{t,i}$ is the desired cycle on $X \times_S T$ over T . The verification is straightforward but lengthy, and is given in [RelCh, 3.3]. \square

DEFINITION 1A.9. A relative cycle W is called **universally integral** when its pullbacks W_T always have integer coefficients; see [RelCh, 3.3.9].

We define $c(X/S, 0)$ to be the free abelian group on the universally integral relative cycles of X which are finite and surjective over S . Finally we set

$Cor_S(X, Y) = c(X \times_S Y/X, 0)$. That is, $Cor_S(X, Y)$ is the group of universally integral cycles on $X \times_S Y$ whose support is finite over X (i.e., proper over X of relative dimension 0).

In [RelCh] the notation $z(X/S, 0)$ was used for the subgroup of $Cycl(X/S, 0)$ generated by universally integral cycles, and the notation $c(X/S, 0)$ was introduced for the subgroup generated by the proper cycles in $z(X/S, 0)$.

The following theorem was proved in [RelCh, 3.3.15] and [RelCh, 3.4.8].

THEOREM 1A.10. *Any relative cycle of X over S is universally integral provided that either*

- (1) S is regular, or
- (2) X is a smooth curve over S .

DEFINITION 1A.11. The composition of relative cycles $V \in Cor_S(X, Y)$ and $W \in Cor_S(Y, Z)$ is defined as follows. Form the pullback W_V of W with respect to the map $V \rightarrow Y$, as in 1A.8. The composition $W \circ V$ is defined to be the push-forward of W_V along the projection $p : X \times Y \times Z \rightarrow X \times Z$. By [RelCh, 3.7.5], the composition will be a universally integral cycle which is finite and surjective over X .

In the special case when V is the graph of $f : X \rightarrow Y$, we see that $W \circ V$ is just the pullback W_X of 1A.8. That is, $Cor_S(Y, Z) \rightarrow Cor_S(X, Z)$ is $W \mapsto W_X$.

EXAMPLE 1A.12. By definition, $c(X/S, 0) = Cor_S(S, X)$. If S and X are smooth over a field k , then clearly $Cor_S(S, X) \subseteq Cor_k(S, X)$ via the embedding of X in $S \times X$. Hence, for every map $S' \rightarrow S$, there is a map $c(X/S, 0) \rightarrow c(X \times_S S'/S', 0)$ induced by composition in Cor_k .

$$\begin{array}{ccc} c(X/S, 0) & \hookrightarrow & Cor_k(S, X) \\ \vdots & & \downarrow \\ c(X \times_S S'/S', 0) & \hookrightarrow & Cor_k(S', X) \end{array}$$

EXAMPLE 1A.13. If $S = \text{Spec } k$ for a field k and X and Y are smooth over S , then the group $Cor_S(X, Y) = c(X \times Y/X, 0)$ agrees with the group $Cor_k(X, Y)$ of definition 1.1.

To see this, note that $c(X \times Y/X, 0) \subseteq Cor_k(X, Y)$ by definition. By 1A.6 and 1A.10, every cycle in $X \times Y$ which is finite and surjective over X is a universally integral relative cycle, so we have equality.

Since composition in Cor_S (as defined in 1A.11) evidently agrees with composition in Cor_k , we see that Cor_k is just the restriction of Cor_S to Sm/k .

EXAMPLE 1A.14. Suppose that $V \subset S$ is a closed immersion of regular schemes and let W be an equidimensional cycle on a scheme X of finite type over S . It is shown in [RelCh, 3.5.8] that the pullback cycle W_V coincides with the image of W under the pullback homomorphism for the map $V \times_S X \rightarrow X$ as defined in [Ser65] and [Ful84], using an alternating sum of Tor terms.

LECTURE 2

Presheaves with transfers

In order to define motivic cohomology we need to introduce the notion of a presheaf with transfers. In this lecture we develop the basic properties of presheaves with transfers.

DEFINITION 2.1. A **presheaf with transfers** is a contravariant additive functor $F : Cor_k \rightarrow \mathbf{Ab}$. We will write $PreSh(Cor_k)$, or $\mathbf{PST}(k)$ or \mathbf{PST} if the field is understood, for the functor category whose objects are presheaves with transfers and whose morphisms are natural transformations.

By additivity, there is a pairing $Cor_k(X, Y) \otimes F(Y) \rightarrow F(X)$ for all F, X and Y .

Restricting to the subcategory Sm/k of Cor_k , we see that a presheaf with transfers F may be regarded as a presheaf of abelian groups on Sm/k which is equipped with extra “transfer” maps $F(Y) \rightarrow F(X)$ indexed by finite correspondences from X to Y .

EXAMPLE 2.2. Every constant presheaf A on Sm/k may be regarded as a presheaf with transfers. If W is an elementary correspondence from X to Y (both connected), the homomorphism $A \rightarrow A$ defined by W is multiplication by the degree of W over X .

The following theorem is a special case of a well known result on functor categories, see [Wei94] 1.6.4 and Exercises 2.3.7 and 2.3.8.

THEOREM 2.3. *The category $\mathbf{PST}(k)$ is abelian and has enough injectives and projectives.*

EXAMPLE 2.4. The sheaf \mathcal{O}^* of global units and the sheaf \mathcal{O} of global functions are two examples of presheaves with transfers.

Recall first that if X is normal and $W \rightarrow X$ is finite and surjective then there is a norm map $N : \mathcal{O}^*(W) \rightarrow \mathcal{O}^*(X)$ induced from the usual norm map on the function fields, $k(W)^* \rightarrow k(X)^*$. Indeed, if $f \in \mathcal{O}^*(W)$ then Nf and Nf^{-1} are both in the integrally closed subring $\mathcal{O}(X)$ of $k(X)$.

Similarly, there is a trace map $\mathrm{Tr} : \mathcal{O}(W) \rightarrow \mathcal{O}(X)$ induced from the usual trace map on the function fields, $k(W) \rightarrow k(X)$. Indeed, if $f \in \mathcal{O}(W)$ then $\mathrm{Tr}f$ belongs to the integrally closed subring $\mathcal{O}(X)$ of $k(X)$.

If $W \subset X \times Y$ is an elementary correspondence from X to Y , we define the transfer map $\mathcal{O}^*(Y) \rightarrow \mathcal{O}^*(X)$ associated to W to be the composition

$$\mathcal{O}^*(Y) \longrightarrow \mathcal{O}^*(W) \xrightarrow{N} \mathcal{O}^*(X).$$

We define the transfer $\mathcal{O}(Y) \rightarrow \mathcal{O}(X)$ associated to W to be the composition

$$\mathcal{O}(Y) \longrightarrow \mathcal{O}(W) \xrightarrow{\text{Tr}} \mathcal{O}(X).$$

We omit the verification that these transfers are compatible with the composition in Cor_k . It is clear from the transfer formula that the subsheaf μ_n of n^{th} roots of unity in \mathcal{O}^* is also a presheaf with transfers, and that the subsheaf k of \mathcal{O} is just the constant sheaf with transfers described in 2.2.

EXAMPLE 2.5. The classical Chow groups $CH^i(-)$ are presheaves with transfers. To see this, we need to construct a map $\phi_W : CH^i(Y) \rightarrow CH^i(X)$ for each elementary finite correspondence W from a smooth scheme X to a smooth scheme Y , and check that this defines a functor from Cor_k to abelian groups.

The correspondence homomorphism ϕ_W is given by the formula $\phi_W(\alpha) = q_*(W \cdot p^*\alpha)$, where $\alpha \in CH^i(Y)$. Here $p^* : CH^i(Y) \rightarrow CH^i(X \times Y)$ is the flat pull-back along the projection $X \times Y \rightarrow Y$, the ‘ \cdot ’ is the intersection product (see 17A.1), and $q : X \times Y \rightarrow X$ is the projection. If Y were proper, this would be exactly the formula given in Chapter 16 of [Ful84]. For general Y , we need to observe that $W \cdot p^*\alpha$ has finite support over X , so that the push-forward $q_*(W \cdot p^*\alpha)$ is defined in $CH^i(X)$.

The verification that the definition of ϕ_W is compatible with the composition of correspondences is now a routine calculation using the projection formula; it is practically the same as the calculation in the proper case, which is given in [Ful84, 16.1.2].

EXAMPLE 2.6. We will see in 13.11 that the motivic cohomology groups $H^{p,q}(-, \mathbb{Z})$ of 3.4 are presheaves with transfers.

EXAMPLE 2.7. The functor K_0 , considered as a presheaf of abelian groups on Sm/k , has no extension to a presheaf with transfers. To see this, it suffices to find a finite étale cover $f : Y \rightarrow X$ of degree 2 and an element $x \in K_0(X)$ such that $f^*(x) = 0$ but $2x \neq 0$. Indeed, if $\Phi \in \text{Cor}(X, Y)$ is the canonical “transfer” morphism defined by f , then $f \circ \Phi = 2$ in $\text{Cor}(X, X)$ (cf. 1.11), so any presheaf with transfers F would have $F(\Phi)f^*(x) = 2x$ for all $x \in F(X)$.

Let \mathcal{L} be a line bundle on a smooth variety X satisfying $\mathcal{L}^2 \cong \mathcal{O}_X$ but $[\mathcal{L} \oplus \mathcal{L}] \neq [\mathcal{O}_X \oplus \mathcal{O}_X]$ in $K_0(X)$. It is well-known that such \mathcal{L} exists; see [Swa62]. It is also well-known that there is an étale cover $f : Y \rightarrow X$ of degree 2 with $Y = \text{Spec}(\mathcal{O}_X \oplus \mathcal{L})$; see [Har77, IV Ex.2.7]. Since $f^*\mathcal{L} \cong \mathcal{O}_Y$, the element $x = [\mathcal{L}] - [\mathcal{O}_X]$ of $K_0(X)$ satisfies $f^*(x) = 0$ but $2x \neq 0$, as required.

Representable functors provide another important class of presheaves with transfers. We will use the notation $\mathbb{Z}_{\text{tr}}(X)$, which was introduced in [SV00]; the

alternate terminology $L(X)$ was used in [TriCa], while $c_{\text{equi}}(X/\text{Spec } k, 0)$ was used in [RelCh] and [CohTh].

By the Yoneda lemma, representable functors provide embeddings of Sm/k and Cor_k into an abelian category, namely $\mathbf{PST}(k)$:

$$\begin{array}{ccccc} Sm/k & \longrightarrow & Cor_k & \longrightarrow & \mathbf{PST}(k) \\ X & \longmapsto & X & \longmapsto & \mathbb{Z}_{tr}(X). \end{array}$$

DEFINITION 2.8. If X is a smooth scheme over k we let $\mathbb{Z}_{tr}(X)$ denote the presheaf with transfers represented by X , so that $\mathbb{Z}_{tr}(X)(U) = Cor(U, X)$. By the Yoneda lemma,

$$\mathbf{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(X), F) \cong F(X).$$

It follows that $\mathbb{Z}_{tr}(X)$ is a projective object in $\mathbf{PST}(k)$.

For every X and U , $\mathbb{Z}_{tr}(X)(U)$ is the group of finite correspondences from U to X and the map $\mathbb{Z}_{tr}(X)(U) \rightarrow \mathbb{Z}_{tr}(X)(V)$ associated to a morphism $f : V \rightarrow U$ is defined to be composition with the correspondence associated to f .

We will write \mathbb{Z} for the presheaf with transfers $\mathbb{Z}_{tr}(\text{Spec } k)$; it is just the constant Zariski sheaf \mathbb{Z} on Sm/k , equipped with the transfer maps of 2.2. Thus the structure map $X \rightarrow \text{Spec } k$ induces a natural map $\mathbb{Z}_{tr}(X) \rightarrow \mathbb{Z}$.

Here are three exercises. Carefully writing up their solutions requires some knowledge about cycles, such as that found in [Ful84].

EXERCISE 2.9. If F is a presheaf with transfers and T is a smooth scheme, define $F^T(U) = F(U \times T)$. Show that F^T is a presheaf with transfers and that every morphism $S \rightarrow T$ induces a morphism $F^T \rightarrow F^S$ of presheaves with transfers. If F is constant and T is geometrically connected, then $F^T = F$.

EXERCISE 2.10. If $k \subset L$ is a separable field extension, every X in Sm/L is an inverse limit of schemes X_α in Sm/k . For every presheaf with transfers F over k , we set $F(X) = \varinjlim F(X_\alpha)$. Show that this makes F a presheaf with transfers over L .

EXERCISE 2.11. Let X be a (non-smooth) scheme of finite type over k . For each smooth U , define $\mathbb{Z}_{tr}(X)(U)$ to be the group $Cor(U, X)$ of 1.1. Show that the composition \circ defined after lemma 1.4 makes $\mathbb{Z}_{tr}(X)$ into a presheaf with transfers.

Given a pointed scheme (X, x) , we define $\mathbb{Z}_{tr}(X, x)$ to be the cokernel of the map $x_* : \mathbb{Z} \rightarrow \mathbb{Z}_{tr}(X)$ associated to the point $x : \text{Spec } k \rightarrow X$. Since x_* splits the structure map $\mathbb{Z}_{tr}(X) \rightarrow \mathbb{Z}$, we have a natural splitting $\mathbb{Z}_{tr}(X) \cong \mathbb{Z} \oplus \mathbb{Z}_{tr}(X, x)$.

The pointed scheme $\mathbb{G}_m = (\mathbb{A}^1 - 0, 1)$ and its associated presheaf with transfers $\mathbb{Z}_{tr}(\mathbb{G}_m) = \mathbb{Z}_{tr}(\mathbb{A}^1 - 0, 1)$ will be of particular interest to us.

DEFINITION 2.12. If (X_i, x_i) are pointed schemes for $i = 1, \dots, n$ we define $\mathbb{Z}_{tr}((X_1, x_1) \wedge \dots \wedge (X_n, x_n))$, or $\mathbb{Z}_{tr}(X_1 \wedge \dots \wedge X_n)$, to be:

$$\text{coker} \left(\bigoplus_i \mathbb{Z}_{tr}(X_1 \times \dots \times \hat{X}_i \times \dots \times X_n) \xrightarrow{id \times \dots \times x_i \times \dots \times id} \mathbb{Z}_{tr}(X_1 \times \dots \times X_n) \right).$$

By definition $\mathbb{Z}_{tr}((X,x)^{\wedge 1}) = \mathbb{Z}_{tr}(X,x)$ and $\mathbb{Z}_{tr}((X,x)^{\wedge q}) = \mathbb{Z}_{tr}((X,x) \wedge \cdots \wedge (X,x))$ for $q > 0$. By convention $\mathbb{Z}_{tr}((X,x)^{\wedge 0}) = \mathbb{Z}$ and $\mathbb{Z}_{tr}((X,x)^{\wedge q}) = 0$ when $q < 0$.

LEMMA 2.13. *The presheaf $\mathbb{Z}_{tr}((X_1,x_1) \wedge \cdots \wedge (X_n,x_n))$ is a direct summand of $\mathbb{Z}_{tr}(X_1 \times \cdots \times X_n)$. In particular, it is a projective object of **PST**.*

Moreover, the following sequence of presheaves with transfers is split-exact:

$$\begin{aligned} 0 \rightarrow \mathbb{Z} \xrightarrow{\{x_i\}} \bigoplus_i \mathbb{Z}_{tr}(X_i) \rightarrow \bigoplus_{i,j} \mathbb{Z}_{tr}(X_i \times X_j) \rightarrow \cdots \\ \cdots \rightarrow \bigoplus_{i,j} \mathbb{Z}_{tr}(X_1 \times \cdots \hat{X}_i \cdots \hat{X}_j \cdots \times X_n) \rightarrow \bigoplus_i \mathbb{Z}_{tr}(X \times \cdots \hat{X}_i \cdots \times X_n) \rightarrow \\ \rightarrow \mathbb{Z}_{tr}(X_1 \times \cdots \times X_n) \rightarrow \mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_n) \rightarrow 0. \end{aligned}$$

This lemma is illustrated by the formulas $\mathbb{Z}_{tr}(X) \cong \mathbb{Z} \oplus \mathbb{Z}_{tr}(X,x)$ and

$$\mathbb{Z}_{tr}(X_1 \times X_2) \cong \mathbb{Z} \oplus \mathbb{Z}_{tr}(X_1,x_1) \oplus \mathbb{Z}_{tr}(X_2,x_2) \oplus \mathbb{Z}_{tr}(X_1 \wedge X_2).$$

PROOF. The projections $[x_i] : X_i \rightarrow \{x_i\} \rightarrow X_i$ are idempotent, as are the correspondences $e_i = 1_{X_i} - [x_i]$. These idempotents induce a decomposition of $\mathbb{Z}_{tr}(X_1 \times \cdots \times X_n)$ into 2^n summands, and we see by inspection that $\mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_n)$ is the image of $e_1 \times \cdots \times e_n$. Since $\mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_n)$ is a summand of a projective object, it is projective. The individual terms in the indicated sequence decompose in a similar fashion, and each map is a projection followed by an inclusion; it is easy to see from this description that the sequence is split-exact (see [Wei94, 1.4.1]). \square

We shall also need a functorial construction of a chain complex associated to a presheaf with transfers. For this we use the cosimplicial scheme Δ^\bullet over k which is defined by:

$$\Delta^n = \text{Spec} k[x_0, \dots, x_n] / \left(\sum_{i=0}^n x_i = 1 \right).$$

The j^{th} face map $\partial_j : \Delta^n \rightarrow \Delta^{n+1}$ is given by the equation $x_j = 0$. Although this construction is clearly taken from topology, the use of Δ^\bullet in an algebraic setting originated with D. Rector in [Rec71].

DEFINITION 2.14. If F is a presheaf of abelian groups on Sm/k , $F(\Delta^\bullet)$ and $F(U \times \Delta^\bullet)$ are simplicial abelian groups. We will write $C_\bullet F$ for the simplicial presheaf $U \mapsto F(U \times \Delta^\bullet)$, i.e., $C_n(F)(U) = F(U \times \Delta^n)$. If F is a presheaf with transfers, $C_\bullet F$ is a simplicial presheaf with transfers by exercise 2.9.

As usual, we can take the alternating sum of the face maps to get a chain complex of presheaves (resp., presheaves with transfers) which (using $*$ in place of \bullet), we will call $C_* F$. It sends U to the complex of abelian groups:

$$\cdots \rightarrow F(U \times \Delta^2) \rightarrow F(U \times \Delta^1) \rightarrow F(U) \rightarrow 0.$$

Both $F \mapsto C_\bullet F$ and $F \mapsto C_* F$ are exact functors. Moreover, the Dold-Kan correspondence (see [Wei94, 8.4.1]), which describes an equivalence between simplicial objects and positive chain complexes, associates to $C_\bullet F$ a normalized subcomplex $C_*^{DK} F$ of the complex $C_* F$, which is quasi-isomorphic to the complex $C_* F$.

If A is the constant presheaf with transfers $A(U) = A$ then C_*A is the complex $\cdots \rightarrow A \xrightarrow{id} A \xrightarrow{0} A \rightarrow 0$; it is quasi-isomorphic to $C_*^{DK}(A)$, which is A regarded as a complex concentrated in degree zero.

Homotopy invariant presheaves

We now introduce a special class of presheaves which will play a major role in these notes.

DEFINITION 2.15. A presheaf F is **homotopy invariant** if for every X the map $p^* : F(X) \rightarrow F(X \times \mathbb{A}^1)$ is an isomorphism. As $p : X \times \mathbb{A}^1 \rightarrow X$ has a section, p^* is always split injective. Thus homotopy invariance of F is equivalent to p^* being onto.

The homotopy invariant presheaves of abelian groups form a Serre subcategory of presheaves, meaning that if $0 \rightarrow F_0 \rightarrow F_1 \rightarrow F_2 \rightarrow 0$ is an exact sequence of presheaves then F_1 is homotopy invariant if and only if both F_0 and F_2 are. In particular, if F and G are homotopy invariant presheaves with transfers then the kernel and the cokernel of every map $f : F \rightarrow G$ are homotopy invariant presheaves with transfers.

Let $i_\alpha : X \hookrightarrow X \times \mathbb{A}^1$ be the inclusion $x \mapsto (x, \alpha)$. We shall write i_α^* for $F(i_\alpha) : F(X \times \mathbb{A}^1) \rightarrow F(X)$.

LEMMA 2.16. F is homotopy invariant if and only if

$$i_0^* = i_1^* : F(X \times \mathbb{A}^1) \rightarrow F(X) \quad \text{for all } X.$$


PROOF. ([Swa72, 4.1]) One direction is clear, so suppose that $i_0^* = i_1^*$ for all X . Applying F to the multiplication map $m : \mathbb{A}^1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^1$, $(x, y) \mapsto xy$, yields the diagram

$$\begin{array}{ccccc} & & F(X \times \mathbb{A}^1) & \xrightarrow{i_0^*} & F(X) \\ & \nearrow 1_{X \times \mathbb{A}^1} & \downarrow (1_X \times m)^* & & \downarrow p^* \\ F(X \times \mathbb{A}^1) & \xleftarrow{(i_1 \times 1_{\mathbb{A}^1})^*} & F(X \times \mathbb{A}^1 \times \mathbb{A}^1) & \xrightarrow{(i_0 \times 1_{\mathbb{A}^1})^*} & F(X \times \mathbb{A}^1). \end{array}$$

Hence $p^*i_0^* = (1 \times i_0)^*m^* = (1 \times i_1)^*m^* = id$. Since $i_0^*p^* = id$, p^* is an isomorphism. \square

DEFINITION 2.17. For $i = 0, \dots, n$ we define $\theta_i : \Delta^{n+1} \rightarrow \Delta^n \times \mathbb{A}^1$ to be the map that sends the vertex v_j to $v_j \times \{0\}$ for $j \leq i$ and to $v_{j-1} \times \{1\}$ otherwise. (See figure 2.1.) These are the algebraic analogues of the top-dimensional simplices in the standard simplicial decomposition of the polyhedron $\Delta^n \times \Delta^1$.

LEMMA 2.18. Let F be a presheaf. Then the maps $i_0^*, i_1^* : C_*F(X \times \mathbb{A}^1) \rightarrow C_*F(X)$ are chain homotopic.

FIGURE 2.1. Simplicial decomposition of $\Delta^n \times \mathbb{A}^1$

PROOF. The maps θ_i defined in 2.17 induce maps

$$h_i = F(1_X \times \theta_i) : C_n F(X \times \mathbb{A}^1) \rightarrow C_{n+1} F(X).$$

The h_i form a simplicial homotopy ([Wei94, 8.3.11]) from $i_1^* = \partial_0 h_0$ to $i_0^* = \partial_{n+1} h_n$. By [Wei94, 8.3.13], the alternating sum $s_n = \sum (-1)^i h_i$ is a chain homotopy from i_1^* to i_0^* . \square

Combining lemmas 2.16 and 2.18, we obtain

COROLLARY 2.19. *If F is a presheaf then the homology presheaves*

$$H_n C_* F : X \mapsto H_n C_* F(X)$$

are homotopy invariant for all n .

EXAMPLE 2.20. ([Swa72, 4.2]) The surjection $F \rightarrow H_0 C_* F$ is the universal morphism from F to a homotopy invariant presheaf.

EXERCISE 2.21. Set $H_0^{\text{sing}}(X/k) = H_0 C_* \mathbb{Z}_{\text{tr}}(X)(\text{Speck})$. Show that there is a natural surjection from $H_0^{\text{sing}}(X/k)$ to $CH_0(X)$, the Chow group of zero cycles modulo rational equivalence (see exercise 1.10). If X is projective, $H_0^{\text{sing}}(X/k) \cong CH_0(X)$. If $X = \mathbb{A}^1$, show that $H_0^{\text{sing}}(\mathbb{A}^1/k) = \mathbb{Z}$. We will return to this point in 7.1.

LEMMA 2.22. *Let F be a presheaf of abelian groups. Suppose that for every smooth scheme X there is a natural homomorphism $h_X : F(X) \rightarrow F(X \times \mathbb{A}^1)$ which fits into the diagram*

$$\begin{array}{ccccc}
 & & F(X) & & \\
 & \swarrow 0 & \downarrow h_X & \searrow id & \\
 F(X) & \xleftarrow{F(i_0)} & F(X \times \mathbb{A}^1) & \xrightarrow{F(i_1)} & F(X).
 \end{array}$$

Then the complex $C_ F$ is chain contractible.*

The assertion that h_X is natural means that for every map $f : X \rightarrow Y$ we have a commutative diagram

$$\begin{array}{ccc} F(X) & \xrightarrow{h_X} & F(X \times \mathbb{A}^1) \\ \uparrow & & \uparrow \\ F(Y) & \xrightarrow{h_Y} & F(Y \times \mathbb{A}^1). \end{array}$$

PROOF. By naturality, h_X induces a map $C_*h : C_*F(X) \rightarrow C_*F(X \times \mathbb{A}^1)$. By lemma 2.18, the identity map $id = i_1^*(C_*h)$ is chain homotopic to $0 = i_0^*(C_*h)$. \square

EXAMPLE 2.23. The prototype for lemma 2.22 is the sheaf of global functions. The complex $C_*\mathcal{O}$ is chain contractible, because $\mathcal{O}(X \times \mathbb{A}^1) \cong \mathcal{O}(X)[t]$ and $h_X(f) = tf$ satisfies the conditions of lemma 2.22.

Here is a second application of lemma 2.22. Note that the projection $p : X \times \mathbb{A}^1 \rightarrow X$ induces a map $\mathbb{Z}_{tr}(X \times \mathbb{A}^1) \rightarrow \mathbb{Z}_{tr}(X)$.

COROLLARY 2.24. $C_*\mathbb{Z}_{tr}(X \times \mathbb{A}^1) \rightarrow C_*\mathbb{Z}_{tr}(X)$ is a chain homotopy equivalence.

PROOF. Let F denote the cokernel of $\mathbb{Z}_{tr}(i_0) : \mathbb{Z}_{tr}(X) \rightarrow \mathbb{Z}_{tr}(X \times \mathbb{A}^1)$ induced by $i_0 : X \rightarrow X \times \mathbb{A}^1$. That is, each $F(U)$ is the cokernel of $Cor(U, X) \rightarrow Cor(U, X \times \mathbb{A}^1)$. Let H_U denote the composition of the product with \mathbb{A}^1 and multiplication $\mathbb{A}^1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^1$:

$$Cor(U, X \times \mathbb{A}^1) \rightarrow Cor(U \times \mathbb{A}^1, (X \times \mathbb{A}^1) \times \mathbb{A}^1) \rightarrow Cor(U \times \mathbb{A}^1, X \times \mathbb{A}^1).$$

Since H_U sends $Cor(U, X \times \{0\})$ to $Cor(U \times \mathbb{A}^1, X \times \{0\})$, it induces a natural map $h_U : F(U) \rightarrow F(U \times \mathbb{A}^1)$. For $U = X \times \mathbb{A}^1$ it is easy to see that the composition of H_U with $i_0, i_1 : U \rightarrow U \times \mathbb{A}^1$ sends $1_U \in Cor(U, X \times \mathbb{A}^1)$ to the projection $i_0p : U \rightarrow X \rightarrow X \times \mathbb{A}^1$ and 1_U , respectively. Therefore $F(i_0)h_U(1_U) = 0$ and $F(i_1)h_U(1_U) = 1_U$ for $U = X \times \mathbb{A}^1$. For any other U , every element $\tilde{f} \in F(U)$ is the image of $1_{X \times \mathbb{A}^1}$ under some correspondence $f : U \rightarrow X \times \mathbb{A}^1$, so again $F(i_0)h_U(\tilde{f}) = 0$ and $F(i_1)h_U(\tilde{f}) = \tilde{f}$. Therefore lemma 2.22 applies to show that C_*F is chain contractible. Since $C_*\mathbb{Z}_{tr}(X \times \mathbb{A}^1) \cong C_*\mathbb{Z}_{tr}(X) \oplus C_*F$, we are done. \square

An elementary \mathbb{A}^1 -homotopy between two morphisms $f, g : X \rightarrow Y$ is a map $h : X \times \mathbb{A}^1 \rightarrow Y$ so that f and g are the restrictions of h along $X \times 0$ and $X \times 1$. This relation is not transitive (exercise!). To correct this, we pass to correspondences.

DEFINITION 2.25. We say that two finite correspondences from X to Y are \mathbb{A}^1 -homotopic if they are the restrictions along $X \times 0$ and $X \times 1$ of an element of $Cor(X \times \mathbb{A}^1, Y)$. This is an equivalence relation on $Cor(X, Y)$. The sum and composition of \mathbb{A}^1 -homotopic maps are \mathbb{A}^1 -homotopic, so the \mathbb{A}^1 -homotopy classes of finite correspondences form the morphisms of an additive category.

We say that $f : X \rightarrow Y$ is an \mathbb{A}^1 -homotopy equivalence if there exists a $g : Y \rightarrow X$ so that fg and gf are \mathbb{A}^1 -homotopic to the identity.

The projection $p : X \times \mathbb{A}^1 \rightarrow X$ is the prototype of an \mathbb{A}^1 -homotopy equivalence; its \mathbb{A}^1 -homotopy inverse is given by the zero-section.

LEMMA 2.26. *If $f : X \rightarrow Y$ is an \mathbb{A}^1 -homotopy equivalence with \mathbb{A}^1 -homotopy inverse g , then $f_* : C_*\mathbb{Z}_{tr}(X) \rightarrow C_*\mathbb{Z}_{tr}(Y)$ is a chain homotopy equivalence with chain homotopy inverse g_* .*

PROOF. Applying $C_*\mathbb{Z}_{tr}$ to the data gives a diagram

$$\begin{array}{ccccc}
 & & C_*\mathbb{Z}_{tr}(X) & & \\
 & \nearrow^{g_*f_*} & \uparrow^{h_*} & \nwarrow_{1_X} & \\
 C_*\mathbb{Z}_{tr}(X) & \xrightarrow[\cong]{(i_0)_*} & C_*\mathbb{Z}_{tr}(X \times \mathbb{A}^1) & \xleftarrow[\cong]{(i_1)_*} & C_*\mathbb{Z}_{tr}(X)
 \end{array}$$

and similarly for Y . The horizontal maps are chain homotopy equivalences by 2.24, and are homotopy inverses to p_* . From the right triangle, $h_* \simeq p_*$. From the left triangle, we get $g_*f_* \simeq 1_X$. Similarly, the diagram for Y gives $f_*g_* \simeq 1_Y$. Hence $f_* : C_*\mathbb{Z}_{tr}(X) \rightarrow C_*\mathbb{Z}_{tr}(Y)$ is a chain homotopy equivalence with inverse g_* . \square

EXERCISE 2.27. Show that there is a natural identification for every X and Y :

$$H_0C_*\mathbb{Z}_{tr}(Y)(X) = Cor(X, Y)/\mathbb{A}^1\text{-homotopy}.$$

We will return to the subject of \mathbb{A}^1 -homotopy in lectures 7, 9, 13, and 14; see 7.2, 9.9 and 14.14.

The motive associated to X will be the class $M(X)$ of $C_*\mathbb{Z}_{tr}(X)$ in an appropriate triangulated category $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ constructed in 14.1 from the derived category of $\mathbf{PST}(k)$. By 2.24, we have $M(X) \cong M(X \times \mathbb{A}^1)$ for all X . More generally, any \mathbb{A}^1 -homotopy equivalence $X \rightarrow Y$ induces an isomorphism $M(X) \cong M(Y)$ by 2.26.

EXERCISE 2.28. If $k \subset F$ is a finite separable field extension, exercise 1.12 implies that there are adjoint functors $i^* : \mathbf{PST}(k) \rightarrow \mathbf{PST}(F)$, $i_* : \mathbf{PST}(F) \rightarrow \mathbf{PST}(k)$. Show that there is a natural transformation $\pi : i^*i_*M \rightarrow M$ whose composition $\pi\eta$ with the adjunction map $\eta : M \rightarrow i^*i_*M$ is multiplication by $[F : k]$ on M . *Hint:* $X_F \rightarrow X$ is finite.

LECTURE 3

Motivic cohomology

Using the tools developed in the last lecture, we will define motivic cohomology. It will be hypercohomology with coefficients in the special cochain complexes $\mathbb{Z}(q)$, called motivic complexes.

DEFINITION 3.1. For every integer $q \geq 0$ the **motivic complex** $\mathbb{Z}(q)$ is defined as the following complex of presheaves with transfers:

$$\mathbb{Z}(q) = C_*\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge q})[-q].$$

We consider $\mathbb{Z}(q)$ to be a bounded above cochain complex; the shifting convention for $[-q]$ implies that the terms $\mathbb{Z}(q)^i = C_{q-i}\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge q})$ vanish whenever $i > q$, and the term with $i = q$ is $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge q})$.

If A is any other abelian group then $A(q) = \mathbb{Z}(q) \otimes A$ is another complex of presheaves with transfers.

When $q = 0$, we have $\mathbb{Z}(0) = C_*(\mathbb{Z})$. As observed after 2.14 above, $\mathbb{Z}(0)$ is quasi-isomorphic to \mathbb{Z} , regarded as a complex concentrated in degree 0.

When $q = 1$, we have $\mathbb{Z}(1) = C_*\mathbb{Z}_{tr}(\mathbb{G}_m)[-1]$. We will give another description of $\mathbb{Z}(1)$ in the next lecture.

By convention $\mathbb{Z}(q) = 0$ if $q < 0$.

We now show that these complexes of presheaves are actually complexes of sheaves with respect to the Zariski topology. Later on, in 6.2, we will show that the $\mathbb{Z}_{tr}(Y)$ are also sheaves in the étale topology.

LEMMA 3.2. *For every scheme Y over k , $\mathbb{Z}_{tr}(Y)$ is a sheaf in the Zariski topology, and $C_*\mathbb{Z}_{tr}(Y)$ is a chain complex of sheaves.*

Similarly, if A is any abelian group, the proof of 3.2 shows that $A \otimes \mathbb{Z}_{tr}(Y)$ is a sheaf in the Zariski topology, and $A \otimes C_*\mathbb{Z}_{tr}(Y)$ is a complex of sheaves.

PROOF. We have to prove that whenever U is covered by U_1 and U_2 the sequence

$$0 \rightarrow \text{Cor}(U, Y) \xrightarrow{\text{diag}} \text{Cor}(U_1, Y) \oplus \text{Cor}(U_2, Y) \xrightarrow{(+, -)} \text{Cor}(U_1 \cap U_2, Y)$$

is exact. We may suppose that U is connected and therefore (being smooth) irreducible. As every finite correspondence from U to Y is dominant over U , it is completely determined by the fiber at the generic point of U . Hence $\text{Cor}(U, Y)$ injects into each $\text{Cor}(U_i, Y)$.

To see that the sequence is exact at the other spot, take cycles $Z_1 = \sum_{i \in I} m_i Z_{1i} \subset U_1 \times Y$ and $Z_2 = \sum_{j \in J} n_j Z_{2j} \subset U_2 \times Y$ that coincide on $(U_1 \cap U_2) \times Y$. It is possible to pair up the Z_{1i} and Z_{2j} , since they are determined by their fibers at the common generic point of U , U_1 and U_2 . Hence there is a bijection between I and J such that, if $i \in I$ corresponds to $j \in J$ then $m_i = n_j$ and the restrictions of Z_{1i} and Z_{2j} agree in $(U_1 \cap U_2) \times Y$. Thus we may assume that Z_1 and Z_2 are elementary correspondences. But then their union $Z = Z_1 \cup Z_2$ in $U \times Y$ is a finite correspondence from U to Y , and its restriction to both $U_i \times Y$ is Z_i , i.e., Z is a preimage of the pair.

Now whenever F is a sheaf and X is smooth, each presheaf $U \mapsto F(U \times X)$ is also a sheaf for the Zariski topology. In particular each $C_n F$ is a sheaf and $C_* F$ is a complex of sheaves. Thus $C_* \mathbb{Z}_{tr}(Y)$ is a complex of Zariski sheaves. \square

We have already seen (in exercises 2.21 and 2.27 above) that the complex $C_* \mathbb{Z}_{tr}(Y)$ is not exact. There we showed that the last map may not be surjective, because its cokernel $H_0 C_* \mathbb{Z}_{tr}(Y)(S) = Cor(S, Y) / \mathbb{A}^1$ -homotopy can be non-zero. When $S = \text{Spec}(k)$, it is the group $H_0^{sing}(Y/k)$ described in exercise 2.21 above and 7.3 below.

Recall that the (small) Zariski site X_{Zar} over a scheme X is the category of open subschemes of X , equipped with the Zariski topology.

COROLLARY 3.3. *The restriction $\mathbb{Z}(q)_X$ of $\mathbb{Z}(q)$ to the Zariski site over X is a complex of sheaves in the Zariski topology.*

Similarly, if A is any abelian group, $A(q)$ is a complex of Zariski sheaves.

PROOF. Set $Y = (\mathbb{A}^1 - 0)^q$. By lemma 3.2 we know that $C_* \mathbb{Z}_{tr}(Y)$ is a complex of sheaves. The complex $\mathbb{Z}(q)[q]$ is a direct summand of $C_* \mathbb{Z}_{tr}(Y)$ by lemma 2.13, so it must be a complex of sheaves too. \square

Note that $A(q)$ represents the derived sheaf tensor product $\mathbb{Z}(q) \otimes^{\mathbf{L}} A$, since $\mathbb{Z}(q)$ is a flat complex of sheaves.

DEFINITION 3.4. The **motivic cohomology groups** $H^{p,q}(X, \mathbb{Z})$ are defined to be the hypercohomology of the motivic complexes $\mathbb{Z}(q)$ with respect to the Zariski topology:

$$H^{p,q}(X, \mathbb{Z}) = \mathbb{H}_{Zar}^p(X, \mathbb{Z}(q)).$$

If A is any abelian group, we define:

$$H^{p,q}(X, A) = \mathbb{H}_{Zar}^p(X, A(q)).$$

REMARK 3.5. Motivic cohomology is well-defined even if the $\mathbb{Z}(q)$ are unbounded complexes because the X in Sm/k are finite dimensional; see [Wei94, 10.6.8]. We will see in 13.11 and 14.16 that motivic cohomology is representable in several derived categories.

VANISHING THEOREM 3.6. *For every smooth scheme X and any abelian group A , we have $H^{p,q}(X, A) = 0$ when $p > q + \dim X$.*

PROOF. By definition, the complex $\mathbb{Z}(q)$ is zero in degrees greater than q . Since $H_{Zar}^i(X, F)$ vanishes for every sheaf F when $i > \dim X$, the result is now an immediate consequence of the hypercohomology spectral sequence. \square

We will prove in theorem 19.3 that, for every smooth variety X and any abelian group A , we have $H^{p,q}(X, A) = 0$ for $p > 2q$ as well.

REMARK 3.7. The groups $H^{p,q}(X, \mathbb{Z})$ are contravariantly functorial in X . To see this we need to check that for a morphism $f : X \rightarrow Y$ we can construct a natural map $\mathbb{Z}(q)_Y \rightarrow f_*\mathbb{Z}(q)_X$. But this is true for any complex C of presheaves on Sm/k : for each open $U \subset Y$, the restriction $f^{-1}U \rightarrow U$ induces the desired map from $C_Y(U) = C(U)$ to $f_*C_X(U) = C(f^{-1}U)$.

The groups $H^{p,q}(X, A)$ are also covariantly functorial in k . That is, if $i : k \subset F$ is a field extension, there is a natural map $H^{*,*}(X, A) \rightarrow H^{*,*}(X_F, A)$. It is induced by the sheaf map $\mathbb{Z}(q)_X \rightarrow i_*\mathbb{Z}(q)_{X_F}$ assembled from the natural maps $\mathbb{Z}_{tr}(Y)(U) \rightarrow i_*\mathbb{Z}_{tr}(Y_F)(U) = \mathbb{Z}_{tr}(Y_F)(U_F)$ of exercise 1.12.

PROPOSITION 3.8. *If $k \subset F$ is a finite and separable field extension and U is smooth over F , then the two motivic chain complexes $\mathbb{Z}(q)_U$ (defined using Cor_k and Cor_F , respectively) are isomorphic. Hence the motivic cohomology groups $H^{p,q}(U, A)$ are independent of the choice of the ground field.*

PROOF. Let T be any smooth scheme over k , and T_F its base change over F . By exercise 1.12 the groups $C_n\mathbb{Z}_{tr}(T_F)(U) = Cor_F(U \times_F \Delta_F^n, T_F)$ and $C_*\mathbb{Z}_{tr}(T)(U) = Cor_k(U \times_k \Delta_k^n, T)$ are isomorphic. That is, $C_*\mathbb{Z}_{tr}(T_F)(U) \cong C_*\mathbb{Z}_{tr}(T)(U)$. Letting T be $(\mathbb{A}_k^1 - 0)^q$, the result follows from lemma 2.13, which says that the complex $\mathbb{Z}(q)[q]$ is a direct summand of $C_*\mathbb{Z}_{tr}(T)$ over k , and of $C_*\mathbb{Z}_{tr}(T_F)$ over F . \square

The following colimit lemmas are elementary consequences of exercise 1.13. They will be useful later on.

LEMMA 3.9. (Colimits) *Let $k \subset F$ be a field extension and X smooth over k . Then:*

$$H^{*,*}(X_F, A) = \operatorname{colim}_{\substack{k \subset E \subset F \\ E \text{ of finite type}}} H^{*,*}(X_E, A).$$

If $f : X \rightarrow S$ is a smooth morphism of smooth schemes over k such that S is connected and $F = k(S)$, then:

$$H^{*,*}(X \times_S \operatorname{Spec} F, A) = \operatorname{colim}_{\substack{U \subset S \\ \text{nonempty}}} H^{*,*}(X \times_S U, A).$$

And now we want to introduce a multiplicative structure on the sheaves $\mathbb{Z}(n)$. We will need the following construction:

CONSTRUCTION 3.10. If (X_s, x_s) are pointed schemes for $s = 1, \dots, j$, then for every $i < j$ we define a morphism of presheaves with transfers:

$$\mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_i) \otimes \mathbb{Z}_{tr}(X_{i+1} \wedge \cdots \wedge X_j) \rightarrow \mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_j).$$

Indeed, definition 1.9 provides a map:

$$\begin{aligned} & \mathbb{Z}_{tr}(X_1 \times \cdots \times X_i)(U) \otimes \mathbb{Z}_{tr}(X_{i+1} \times \cdots \times X_j)(U) \\ &= \text{Cor}_k(U, X_1 \times \cdots \times X_i) \otimes \text{Cor}_k(U, X_{i+1} \times \cdots \times X_j) \rightarrow \\ & \rightarrow \text{Cor}_k(U \times U, X_1 \times \cdots \times X_j) = \mathbb{Z}_{tr}(X_1 \times \cdots \times X_j)(U \times U). \end{aligned}$$

Composing with the diagonal $U \rightarrow U \times U$, we have:

$$\mathbb{Z}_{tr}(X_1 \times \cdots \times X_i)(U) \otimes \mathbb{Z}_{tr}(X_{i+1} \times \cdots \times X_j)(U) \xrightarrow{\Delta} \mathbb{Z}_{tr}(X_1 \times \cdots \times X_j)(U).$$

Now recall that by definition $\mathbb{Z}_{tr}(X_1 \wedge \cdots \wedge X_n)$ is a quotient of $\mathbb{Z}_{tr}(X_1 \times \cdots \times X_n)$. It is easy to check that the map Δ factors through the quotient, giving the required morphism.

CONSTRUCTION 3.11. For each m and n we construct a map

$$\mathbb{Z}(m) \otimes \mathbb{Z}(n) \rightarrow \mathbb{Z}(m+n)$$

using the map $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge m}) \otimes \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n}) \rightarrow \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge m+n})$ of 3.10, as follows.

For any smooth U we need to build a map of complexes of abelian groups:

$$\mathbb{Z}(m)[m](U) \otimes \mathbb{Z}(n)[n](U) \rightarrow \mathbb{Z}(m+n)[m+n](U),$$

or equivalently, $\mathbb{Z}(m)(U) \otimes \mathbb{Z}(n)(U) \rightarrow \mathbb{Z}(m+n)(U)$. Recall that by definition 3.1, $\mathbb{Z}(n)[n](U)$ is the chain complex $C_* \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(U)$. Let us write the underlying simplicial object as $A_\bullet^n = \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(U \times \Delta^\bullet)$, and the associated unnormalized chain complex $\mathbb{Z}(n)[n]$ as A_*^n . Similarly, we write $(A_\bullet^m \otimes A_\bullet^n)_*$ for the chain complex associated to $\text{diag}(A_\bullet^m \otimes A_\bullet^n)$. The Eilenberg-Zilber theorem ([Wei94, 8.5.1]) yields a quasi-isomorphism $\nabla : A_*^m \otimes A_*^n \rightarrow (A_\bullet^m \otimes A_\bullet^n)_*$.

Therefore if we find a simplicial map $m : \text{diag} A_\bullet^m \otimes A_\bullet^n \rightarrow A_\bullet^{m+n}$ we have also a map $(A_\bullet^m \otimes A_\bullet^n)_* \rightarrow A_*^{m+n}$ which, composed with the previous one, gives the multiplicative structure. Unfolding the definitions again, we have:

$$A_i^n = \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(U \times \Delta^i).$$

We define the components of m to be the maps of 3.10:

$$\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge m})(U \times \Delta^i) \otimes \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(U \times \Delta^i) \rightarrow \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge (m+n)})(U \times \Delta^i).$$

The morphisms in 3.10 are associative and the map ∇ in the Eilenberg-Zilber theorem is homotopy associative ([Wei94, 8.5.4]). It follows that the pairing of construction 3.11 is homotopy associative.

COROLLARY 3.12. *For each smooth X , there are pairings:*

$$H^{p,q}(X, \mathbb{Z}) \otimes H^{p',q'}(X, \mathbb{Z}) \rightarrow H^{p+p',q+q'}(X, \mathbb{Z}).$$

In 15.9 we will show that this pairing is skew-commutative with respect to the first grading, so that $H^{*,*}(X, \mathbb{Z})$ is an associative graded-commutative ring.

LECTURE 4

Weight one motivic cohomology

In this lecture we describe $\mathbb{Z}(1)$ and $\mathbb{Z}/l(1)$ in terms of units and roots of unity.

THEOREM 4.1. *There is a quasi-isomorphism of complexes of presheaves with transfers:*

$$\mathbb{Z}(1) \xrightarrow{\simeq} \mathcal{O}^*[-1].$$

COROLLARY 4.2. *Let X be a smooth scheme over k . Then we have:*

$$H^{p,q}(X, \mathbb{Z}) = \begin{cases} 0 & q \leq 1 \text{ and } (p, q) \neq (0, 0), (1, 1), (2, 1) \\ \mathbb{Z}(X) & (p, q) = (0, 0) \\ \mathcal{O}^*(X) & (p, q) = (1, 1) \\ \text{Pic}(X) & (p, q) = (2, 1). \end{cases}$$




FIGURE 4.1. Weight q motivic cohomology

This theorem will follow from lemmas 4.3–4.6 below. An alternative proof is given in [SV96].

Consider the functor $\mathcal{M}^*(\mathbb{P}^1; 0, \infty) : \text{Sm}/k \rightarrow \mathbf{Ab}$ which sends a scheme X to the group of rational functions on $X \times \mathbb{P}^1$ which are regular in a neighborhood of $X \times \{0, \infty\}$ and equal to 1 on $X \times \{0, \infty\}$. Clearly $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)$ is a sheaf for the Zariski topology. Given a rational function f on $X \times \mathbb{P}^1$ let $D(f)$ denote its (Weil) divisor.

LEMMA 4.3. *For all f in $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)(X)$, the Weil divisor $D(f)$ belongs to the subgroup $\text{Cor}(X, \mathbb{A}^1 - 0)$ of the group of cycles on $X \times \mathbb{P}^1$.*

PROOF. Since the support of $D(f)$ is disjoint from $X \times \{0, \infty\}$, $D(f)$ is a cycle in $X \times \mathbb{A}^1 - 0$. To see that it is finite and surjective over X we may assume that $X = \text{Spec} A$ is an affine domain. We may write $f = f_+/f_-$ where $f_+ = a_m t^m + \cdots + a_0$ and $f_- = b_n t^n + \cdots + b_0$ are in $A[t]$ and a_m, a_0, b_n and b_0 are nonzero. Since f is regular near $X \times \{0\}$, f_- is relatively prime to t , and we may assume that b_0 is a unit of A . Similarly, we may assume that b_n is a unit of A . Since $f = 1$ on $X \times \infty$, we have $m = n$ and may assume that $a_n = b_n = 1$. But then the divisors $D(f_+)$ of $f_+ = t^n + \cdots + a_0$ and $D(f_-)$ of $f_- = t^n + \cdots + b_0$ are finite and surjective over X . Since they belong to $\text{Cor}(X, \mathbb{A}^1 - 0)$, so does $D(f) = D(f_+) - D(f_-)$. \square

From 4.3 we get a morphism of sheaves: $\mathcal{M}^*(\mathbb{P}^1; 0, \infty) \hookrightarrow \mathbb{Z}_{tr}(\mathbb{A}^1 - 0)$.

LEMMA 4.4. *For any connected X there is a short exact sequence in **Ab**:*

$$0 \longrightarrow \mathcal{M}^*(\mathbb{P}^1; 0, \infty)(X) \longrightarrow \mathbb{Z}_{tr}(\mathbb{A}^1 - \{0\})(X) \xrightarrow{\lambda} \mathbb{Z} \oplus \mathcal{O}^*(X) \longrightarrow 0.$$

PROOF. We know that $\text{Pic}(X \times \mathbb{P}^1) \cong \text{Pic}(X) \times \mathbb{Z}$, so for any Z in $\text{Cor}(X, \mathbb{A}^1) \subset \text{Cor}(X, \mathbb{P}^1)$ there is a unique rational function f on $X \times \mathbb{P}^1$ and an integer n so that $D(f) = Z$ and $f/t^n = 1$ on $X \times \{\infty\}$. If Z lies in $\text{Cor}(X, \mathbb{A}^1 - 0)$, then $f(0) \in \mathcal{O}^*(X)$. We define $\lambda : \mathbb{Z}_{tr}(\mathbb{A}^1 - 0) \rightarrow \mathbb{Z} \oplus \mathcal{O}^*$ by $\lambda(Z) = (n, (-1)^n f(0))$. If $u \in \mathcal{O}^*(X)$ and $Z_u = D(t - u)$ then $\lambda(Z_u) = (1, u)$. Since $\lambda(Z_u - Z_1) = (0, u)$ λ is onto. The kernel of λ consists of all Z whose f lies in $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)(X)$, so we are done. \square

LEMMA 4.5. *The map λ respects transfers. Hence $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)$ is a **PST**.*

PROOF. It is easy to see that the first component of λ is a morphism in **PST** because it is the map $\text{Cor}_k(X, \mathbb{A}^1 - 0) \rightarrow \text{Cor}_k(X, \text{Spec} k)$, induced by the structure map $\pi : \mathbb{A}^1 - 0 \rightarrow \text{Spec} k$. To check the second component of λ , we see from exercise 1.13 that it suffices to check that the following diagram commutes for every finite field extension $F \subset E$.

$$\begin{array}{ccc} \mathbb{Z}_{tr}(\mathbb{A}^1 - 0)(\text{Spec} E) & \longrightarrow & E^* \\ \downarrow & & \downarrow N_{E/F} \\ \mathbb{Z}_{tr}(\mathbb{A}^1 - 0)(\text{Spec} F) & \longrightarrow & F^* \end{array}$$

This is a straightforward verification using exercise 1.10. \square

Write \mathcal{M}^* for $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)$. By 2.14, $(C_i F)(U) = F(U \times \Delta^i)$, so 4.4 gives us:

$$0 \rightarrow C_*(\mathcal{M}^*) \rightarrow C_* \mathbb{Z}_{tr}(\mathbb{A}^1 - 0) \rightarrow C_*(\mathbb{Z} \oplus \mathcal{O}^*) \rightarrow 0.$$

Splitting off $0 \rightarrow C_* \mathbb{Z} = C_* \mathbb{Z} \rightarrow 0$ we get an exact sequence:

$$0 \rightarrow C_*(\mathcal{M}^*) \rightarrow \mathbb{Z}(1)[1] \rightarrow C_*(\mathcal{O}^*) \rightarrow 0.$$

But $C_*(\mathcal{O}^*) \simeq \mathcal{O}^*$ because $\mathcal{O}^*(U \times \Delta^n) = \mathcal{O}^*(U)$. We will prove in lemma 4.6 that the first term $C_*(\mathcal{M}^*)$ is acyclic. Therefore $\mathbb{Z}(1)[1]$ is quasi-isomorphic to \mathcal{O}^* . This is the statement of theorem 4.1, shifted once.

LEMMA 4.6. *If X is a smooth scheme over k , then $C_*(\mathcal{M}^*)(X)$ is an acyclic complex of abelian groups. Hence $C_*(\mathcal{M}^*)$ is an acyclic complex of sheaves.*

PROOF. Let $f \in C_i^{DK}(\mathcal{M}^*)(X)$ be a cycle, i.e., an element vanishing in $C_{i-1}^{DK}(\mathcal{M}^*)(X)$. Then f is a regular function on some neighborhood U of $Z = X \times \Delta^i \times \{0, \infty\}$ in $X \times \Delta^i \times \mathbb{P}^1$, and $f = 1$ on each face $X \times \Delta^{i-1} \times \mathbb{P}^1$, as well as on Z . Consider the regular function $h_X(f) = 1 - t(1 - f)$ on the neighborhood $\mathbb{A}^1 \times U$ of $\mathbb{A}^1 \times Z$ in $\mathbb{A}^1 \times X \times \Delta^i \times \mathbb{P}^1$, where t denotes the coordinate function of \mathbb{A}^1 . Then $h_X(f)$ is a cycle in $C_i^{DK}(\mathcal{M}^*)(\mathbb{A}^1 \times X)$, because it equals 1 where f equals 1. The restrictions along $t = 0, 1$, from $C_i^{DK}(\mathcal{M}^*)(\mathbb{A}^1 \times X)$ to $C_i^{DK}(\mathcal{M}^*)(X)$, send $h_X(f)$ to 1 and f , respectively. Since these restrictions are chain homotopy equivalent by 2.18, f is a boundary. \square

This completes the proof of theorem 4.1.

REMARK 4.7. We will revisit this in lecture 7 in 7.11.

Lemma 4.6 works more generally to show that $C_*\mathcal{M}^*(Y;Z)(X)$ is acyclic for every affine X , where $\mathcal{M}^*(Y;Z)(X)$ is the group of rational functions on $X \times Y$ which are regular in a neighborhood of $X \times Z$ and equal to 1 on $X \times Z$.

Now let us consider the complex $\mathbb{Z}/l(1)$. By theorem 4.1 $\mathbb{Z}(1)$ is quasi-isomorphic to $\mathcal{O}^*[-1]$. Tensoring with \mathbb{Z}/l we have $\mathbb{Z}/l(1) \simeq \mathcal{O}^*[-1] \otimes^L \mathbb{Z}/l$, which is just the complex $[\mathcal{O}^* \xrightarrow{l} \mathcal{O}^*]$ in degrees 0 and 1. Then we have the universal coefficients sequence:

$$0 \longrightarrow H^{p,q}(X, \mathbb{Z})/l \longrightarrow H^{p,q}(X, \mathbb{Z}/l) \longrightarrow {}_l H^{p+1,q}(X, \mathbb{Z}) \longrightarrow 0.$$

COROLLARY 4.8. *There is a quasi-isomorphism of complexes of étale sheaves*

$$\mathbb{Z}/l(1)_{\acute{e}t} \simeq \mu_l.$$

PROOF. Since sheafification is exact ([Mil80] p. 63), theorem 4.1 gives $\mathbb{Z}(1)_{\acute{e}t} \simeq \mathcal{O}_{\acute{e}t}^*[-1]$, and hence

$$\mathbb{Z}/l(1)_{\acute{e}t} \simeq \mathcal{O}_{\acute{e}t}^*[-1] \otimes^{\mathbb{L}} \mathbb{Z}/l \simeq \mu_l. \quad \square$$

COROLLARY 4.9. *If $1/l \in k$ and X is smooth, then $H^{p,1}(X, \mathbb{Z}/l) = 0$ for $p \neq 0, 1, 2$ while:*

$$H^{0,1}(X, \mathbb{Z}/l) = \mu_l(X), \quad H^{1,1}(X, \mathbb{Z}/l) = H_{\acute{e}t}^1(X, \mu_l),$$

$$H^{2,1}(X, \mathbb{Z}/l) = \text{Pic}(X)/l\text{Pic}(X).$$

PROOF. The calculation of $H^{p,1}$ for $p \neq 1$ follows from the universal coefficients sequence, since the only nonzero Zariski cohomology groups of \mathcal{O}^* on a smooth scheme are H^0 and $H^1(X, \mathcal{O}^*) = \text{Pic}(X)$. For $p = 1$ note that corollary 4.8 gives a natural map

$$\mathbb{H}_{\text{Zar}}^*(X, \mathbb{Z}/l(1)) \rightarrow \mathbb{H}_{\acute{e}t}^*(X, \mathbb{Z}/l(1)_{\acute{e}t}) = H_{\acute{e}t}^1(X, \mu_l)$$

fitting into the diagram:

$$\begin{array}{ccccc}
 \mathbb{H}_{\text{Zar}}^1(X, \mathbb{Z}(1))/l & \hookrightarrow & \mathbb{H}_{\text{Zar}}^1(X, \mathbb{Z}/l(1)) & \twoheadrightarrow & {}_l\mathbb{H}_{\text{Zar}}^2(X, \mathbb{Z}(1)) \\
 \cong \downarrow & & \downarrow & & \downarrow \cong \\
 \mathbb{H}_{\acute{e}t}^1(X, \mathbb{Z}(1))/l & \hookrightarrow & \mathbb{H}_{\acute{e}t}^1(X, \mathbb{Z}/l(1)) & \twoheadrightarrow & {}_l\mathbb{H}_{\acute{e}t}^2(X, \mathbb{Z}(1)).
 \end{array}$$

Since $H_{\acute{e}t}^1(X, \mathcal{O}^*) = H_{\text{Zar}}^1(X, \mathcal{O}^*)$ by Hilbert's Theorem 90 (see [Mil80, III 4.9]), the 5-lemma concludes the proof. \square

REMARK 4.10. (Deligne) If $\text{char } k = l$ then $H^{1,1}(X, \mathbb{Z}/l) \cong H_{\text{fpp}}^1(X, \mu_l)$. In fact, the proof of 4.9 is valid in this setting.

LECTURE 5

Relation to Milnor K -Theory

The Milnor K -theory $K_*^M(F)$ of a field F is defined to be the quotient of the tensor algebra $T(F^*)$ over \mathbb{Z} by the ideal generated by the elements of the form $x \otimes (1-x)$ where $x \in F^*$. In particular, $K_0^M(F) = \mathbb{Z}$ and $K_1^M(F) = F^*$.

The goal of this lecture is to prove the following:

THEOREM 5.1. *For any field F and any n we have:*

$$H^{n,n}(\mathrm{Spec} F, \mathbb{Z}) \cong K_n^M(F).$$

We have already seen that this holds for $n = 0, 1$, because by definition 3.4 $H^{0,0}(\mathrm{Spec} F, \mathbb{Z}) = H_{\mathrm{Zar}}^0(\mathrm{Spec} F, \mathbb{Z}) = \mathbb{Z}$ and by theorem 4.1,

$$H^{1,1}(\mathrm{Spec} F, \mathbb{Z}) = H_{\mathrm{Zar}}^1(\mathrm{Spec} F, \mathcal{O}^*[-1]) = H_{\mathrm{Zar}}^0(\mathrm{Spec} F, \mathcal{O}^*) = F^*.$$

The proof of theorem 5.1 will follow [SV00, 3.4] which is based on [NS89]. It will consist of three steps:

- (1) Construction of $\theta : H^{n,n}(\mathrm{Spec} F, \mathbb{Z}) \rightarrow K_n^M(F)$. This will use 5.5.
- (2) Construction of $\lambda_F : K_n^M(F) \rightarrow H^{n,n}(\mathrm{Spec} F, \mathbb{Z})$. This will be done using proposition 5.9 and lemma 5.6. The proof of 5.9 will need 5.8.

$$5.6 + (5.8 \Rightarrow 5.9) \Rightarrow \exists \lambda_F$$

- (3) Proof that these two maps are inverse to each other. For this we will need lemma 5.10 (proved using lemma 5.11).

Before starting the proof of the theorem we need some additional properties of motivic cohomology and Milnor K -theory.

Recall that $\mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge n})(\mathrm{Spec} F)$ is a quotient of $\mathbb{Z}_{\mathrm{tr}}((\mathbb{A}^1 - 0)^n)(\mathrm{Spec} F)$, which by 1.10 is the group of zero cycles of $(\mathbb{A}^1 - 0)^n$.

LEMMA 5.2. *We have $H^{p,q}(\mathrm{Spec} F, \mathbb{Z}) = H_{q-p}(C_* \mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge q})(\mathrm{Spec} F))$ for all p and q . In particular we have*

$$\begin{aligned} H^{n,n}(\mathrm{Spec} F, \mathbb{Z}) &= H_0(C_* \mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge n})(\mathrm{Spec} F)) \\ &= \mathrm{coker} \left(\mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge n})(\mathbb{A}^1) \xrightarrow{\partial_0 - \partial_1} \mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge n})(\mathrm{Spec} F) \right). \end{aligned}$$

PROOF. Write A_* for $C_* \mathbb{Z}_{\mathrm{tr}}(\mathbb{G}_m^{\wedge q})(\mathrm{Spec} F)$ so the right side is $H_{q-p} A_* = H^{p-q} A_*$. By definition 3.1, the restriction of $\mathbb{Z}(q)$ to $\mathrm{Spec} F$ is the chain complex $A_*[-q]$. Since Zariski cohomology on $\mathrm{Spec} F$ is just ordinary cohomology, we have

$$H^{p,q}(\mathrm{Spec} F, \mathbb{Z}) = H^p(A_*[-q]) = H^{p-q}(A_*) = H_{q-p}(A_*). \quad \square$$

LEMMA 5.3. *If $F \subset E$ is a finite field extension, then the proper push-forward of cycles induces a map $N_{E/F} : H^{*,*}(\text{Spec } E, \mathbb{Z}) \rightarrow H^{*,*}(\text{Spec } F, \mathbb{Z})$. Moreover, if $x \in H^{*,*}(\text{Spec } E, \mathbb{Z})$ and $y \in H^{*,*}(\text{Spec } F, \mathbb{Z})$ then:*

- (1) $N_{E/F} : H^{0,0}(\text{Spec } E, \mathbb{Z}) = \mathbb{Z} \rightarrow \mathbb{Z} = H^{0,0}(\text{Spec } F, \mathbb{Z})$ is multiplication by the degree of E/F .
- (2) $N_{E/F} : H^{1,1}(\text{Spec } E, \mathbb{Z}) = E^* \rightarrow F^* = H^{1,1}(\text{Spec } F, \mathbb{Z})$ is the classical norm map $E^* \rightarrow F^*$.
- (3) $N_{E/F}(y_E \cdot x) = y \cdot N_{E/F}(x)$ and $N_{E/F}(x \cdot y_E) = N_{E/F}(x) \cdot y$.
- (4) If $F \subset E \subset K$, and K is normal over F , we have:

$$N_{E/F}(x)_K = [E : F]_{\text{insep}} \sum_{j:E \xrightarrow{\quad} K} j^*(x) \quad \text{in } H^{*,*}(\text{Spec } K, \mathbb{Z}).$$

- (5) If $F \subset E' \subset E$ then $N_{E/F}(x) = N_{E'/F}(N_{E/E'}(x))$.

PROOF. All but property 2 follow immediately from the corresponding properties of proper push-forward. Property 2 follows from property 4 since this formula also holds for the classical norm map $N_{E/F} : E^* \rightarrow F^*$. \square

If $F \subset E$ is a finite field extension, there is a “norm map” $N_{E/F} : K_n^M(E) \rightarrow K_n^M(F)$ satisfying the analogue of lemma 5.3. In addition, it satisfies the following condition (see [Sus82]).

THEOREM 5.4 (Weil Reciprocity). *Suppose that L is an algebraic function field over k . For each discrete valuation w on L there is a map*

$$\partial_w : K_{n+1}^M(L) \rightarrow K_n^M(k(w)),$$

and for all $x \in K_{n+1}^M(L)$:

$$\sum_w N_{k(w)/k} \partial_w(x) = 0.$$

COROLLARY 5.5. *Let $p : Z \rightarrow \mathbb{A}_F^1$ be a finite surjective morphism and suppose that Z is integral. Let $f_1, \dots, f_n \in \mathcal{O}^*(Z)$ and:*

$$p^{-1}(\{0\}) = \coprod n_i^0 z_i^0 \quad p^{-1}(\{1\}) = \coprod n_i^1 z_i^1$$

where n_i^ε are the multiplicities of the points $z_i^\varepsilon = \text{Spec } E_i^\varepsilon$ ($\varepsilon = 0, 1$). Define:

$$\varphi_0 = \sum n_i^0 N_{E_i^0/F}(\{f_1, \dots, f_n\}_{E_i^0}) \quad \varphi_1 = \sum n_i^1 N_{E_i^1/F}(\{f_1, \dots, f_n\}_{E_i^1}).$$

Then we have:

$$\varphi_0 = \varphi_1 \in K_n^M(F).$$

PROOF. Let L be the function field of Z and consider $x = \{t/t-1, f_1, \dots, f_n\}$. At every infinite place, $t/t-1$ equals 1 and $\partial_w(x) = 0$. Similarly, $\partial_w(x) = 0$ at all finite places except those over 0 and 1. If w_i lies over $t = 0$ then $\partial_{w_i}(x) = n_i^0 \{f_1, \dots, f_n\}$ in $K_n^M(E_i^0)$; if w_i lies over $t = 1$ then $\partial_{w_i}(x) = -n_i^1 \{f_1, \dots, f_n\}$ in $K_n^M(E_i^1)$. By Weil Reciprocity 5.4, $\sum N \partial_{w_i}(x) = \varphi_0 - \varphi_1$ vanishes in $K_n^M(F)$. \square

We are now ready to define the map θ . By lemma 5.2 it is enough to find a map f from $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(\text{Spec } F)$ to $K_n^M(F)$ which composed with the difference of the face operators is zero. Such a map must induce a unique map θ on the cokernel:

$$\begin{array}{ccc} \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(\mathbb{A}^1) & \xrightarrow{\partial_0 - \partial_1} & \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(\text{Spec } F) \twoheadrightarrow H^{n,n}(\text{Spec } F, \mathbb{Z}) \\ & & \searrow f \qquad \qquad \downarrow \theta \\ & & K_n^M(F). \end{array}$$

But now $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(\text{Spec } F)$ is a quotient of the free abelian group generated by the closed points of $(\mathbb{A}_F^1 - \{0\})^n$ (by exercise 1.10), modulo the subgroup generated by all points of the form $(x_1, \dots, 1, \dots, x_n)$ where the 1's can be in any position. If x is a closed point of $(\mathbb{A}_F^1 - \{0\})^n$ with residue field E then x is defined by a canonical sequence (x_1, \dots, x_n) of nonzero elements of E . Now E is a finite field extension of F , and $\{x_1, \dots, x_n\} \in K_n^M(E)$. Using the norm map for Milnor K -theory $N_{E/F} : K_n^M(E) \rightarrow K_n^M(F)$, we define

$$f(x) = N_{E/F}(\{x_1, \dots, x_n\}).$$

Since $\{x_1, \dots, 1, \dots, x_n\} = 0$ in $K_n^M(E)$, this induces a well-defined map $f : \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(\text{Spec } F) \rightarrow K_n^M(F)$. By 5.5 the composition of f with the face operators is zero. We define θ to be the map induced on the cokernel.

If x is an F -point of $(\mathbb{A}_F^1 - 0)^n$ then its coordinates x_1, \dots, x_n are nonzero elements of F . We shall write $[x_1 : \dots : x_n]$ for the class of x in $H^{n,n}(\text{Spec } F, \mathbb{Z})$. The map θ is obviously surjective since $\theta([x_1 : \dots : x_n]) = \{x_1, \dots, x_n\}$ for x_1, \dots, x_n in F .

Now let us build the opposite map, λ_F . For this, we will use the multiplicative structure (3.12) on $H^{*,*}(X, \mathbb{Z})$. The following lemma is immediate from construction 3.11 and lemma 5.2.

LEMMA 5.6. *For $a_1, \dots, a_n \in F$ we have $[a_1 : \dots : a_n] = [a_1] \cdots [a_n]$.*

By definition $K_n^M(F) = T(F^*)/(x \otimes (1-x))$. Therefore we define a map:

$$T(F^*) \rightarrow \bigoplus_n H^{n,n}(\text{Spec } F, \mathbb{Z}), \quad a_1 \otimes \cdots \otimes a_n \mapsto [a_1] \cdots [a_n].$$

We will prove that this maps factors through $K_n^M(F)$. By 5.6, it is enough to prove that $[a : 1-a]$ is zero, which is the statement of proposition 5.9 below.

EXAMPLE 5.7. We can use a special cycle to show that $[a : -a] = 0$. Consider the correspondence Z from \mathbb{A}^1 (parametrized by t) to $X = \mathbb{A}^1 - \{0\}$ (parametrized by x) defined by

$$x^2 - t(a+b)x - (1-t)(1+ab)x + ab = 0.$$

Restricting along $t = 0, 1$ yields correspondences $[ab] + [1]$ and $[a] + [b]$ in $\text{Cor}(\text{Spec } F, X)$. Setting these equal recovers the identity $[ab] = [a] + [b]$ in $H^{1,1}(\text{Spec } F, \mathbb{Z}) \cong F^*$, because $[1] = 0$.

Let Y denote the composition of Z with the diagonal embedding $X \hookrightarrow X^2$. Since $[1 : 1] = [1][1] = 0$, equating the restrictions along $t = 0, 1$ yields the identity $[ab : ab] = [ab : ab] + [1 : 1] = [a : a] + [b : b]$ in $H^{2,2}(\text{Spec } F, \mathbb{Z})$. Bilinearity (5.6) yields skew-commutativity: $[a : b] + [b : a] = 0$. In particular, $2[a : a] = 0$.

Passing to $E = F(\sqrt{a})$, we see that $0 = 2[\sqrt{a} : \sqrt{a}] = [a : \sqrt{a}]$ in $H^{2,2}(\text{Spec } E, \mathbb{Z})$. By 5.3, applying $N_{E/F}$ yields $0 = [a : -a]$ in $H^{2,2}(\text{Spec } F, \mathbb{Z})$.

LEMMA 5.8. *Suppose that there exists an $n > 0$ so that $n[x : 1 - x] = 0$ for all finite extensions of F and $x \neq 0, 1$ in F . Then $[x : 1 - x] = 0$ in $H^{2,2}(\text{Spec } F, \mathbb{Z})$ for every $x \neq 0, 1$.*

PROOF. Suppose $n = m \cdot p$ where p is a prime; we want to prove $m[x : 1 - x] = 0$. Let us consider $y = \sqrt[p]{x}$ and $E = F(y)$. Then $0 = mp[y : 1 - y] = m[x : 1 - y]$, and $1 - x = N_{E/F}(1 - y)$. Hence

$$0 = N_{E/F}(m[x : 1 - x]) = m \cdot [x : N_{E/F}(1 - y)] = m[x : 1 - x].$$

The formula $[x : 1 - x] = 0$ follows by induction on n . \square

PROPOSITION 5.9. *The element $[x : 1 - x]$ in $H^{2,2}(\text{Spec } F, \mathbb{Z})$ is the zero element.*

PROOF. Let Z be the finite correspondence from \mathbb{A}^1 (parametrized by t) to $X = \mathbb{A}^1 - 0$ (parametrized by x) defined by:

$$x^3 - t(a^3 + 1)x^2 + t(a^3 + 1)x - a^3 = 0.$$

Let ω be a root of $x^2 + x + 1$, so $\omega^3 = 1$, and $E = F(\omega)$. The fiber over $t = 0$ consists of $a, \omega a$, and $\omega^2 a$ and the fiber over $t = 1$ consists of a^3 and two sixth roots of 1. Using the embedding $x \mapsto (x, 1 - x)$ of $\mathbb{A}^1 - \{0, 1\}$ into X^2 , Z yields a correspondence Z' from \mathbb{A}^1 to X^2 . Then in $H^{2,2}(\text{Spec } E, \mathbb{Z})$

$$\begin{aligned} \partial_0(Z') &= [a : 1 - a] + [\omega a : 1 - \omega a] + [\omega^2 a : 1 - \omega^2 a] \\ &= [a : 1 - a^3] + [\omega : (1 - \omega a)(1 - \omega^2 a)^2] \end{aligned}$$

is equal to

$$\partial_1(Z') = [a^3 : 1 - a^3] + [-\omega : 1 + \omega] + [-\omega^2 : 1 + \omega^2].$$

Multiplying by 3 eliminates terms $[\omega : b]$, noting that $[-1 : 1 + \omega] + [-1 : 1 + \omega^2] = 0$ as $(1 + \omega)(1 + \omega^2) = 1$. Therefore $0 = 2[a^3 : 1 - a^3]$ over E . Applying the norm yields $0 = 4[a^3 : 1 - a^3]$ over F . Passing to the extension $F(\sqrt[3]{a})$ and norming yields $0 = 12[a : 1 - a]$ over F . Applying lemma 5.8 with $n = 12$, we see that $0 = [a : 1 - a]$ as well. \square

Proposition 5.9 shows that the algebra map of lemma 5.6 induces a map on the quotient $\lambda_F : K_n^M(F) \rightarrow H^{n,n}(\text{Spec } F, \mathbb{Z})$. Now we need to check that λ_F and θ are inverse to each other. Since $\theta \circ \lambda_F$ is the identity by construction, it is enough to prove that λ_F is surjective.

LEMMA 5.10. *The map λ_F is surjective.*

PROOF. By 5.2, it suffices to show that if x is a closed point of $X = (\mathbb{A}_F^1 - 0)^n$ then $[x] \in H^{n,n}(\text{Spec } F, \mathbb{Z})$ belongs to the image of λ_F . Set $E = k(x)$, and choose a lift $\tilde{x} \in X_E$ of x . Since x is the proper push-forward of \tilde{x} , the definition of the norm map (see 5.3) implies that:

$$[x] = N_{E/F}([\tilde{x}]) \quad \tilde{x} = (a_1, \dots, a_n) \in (\mathbb{A}^1 - 0)^n(E).$$

Since \tilde{x} is a rational point of X_E , $[\tilde{x}]$ is the image under λ_E of its coordinates. So $[x] = N_{E/F} \lambda_E \{a_1, \dots, a_n\}$. The lemma now follows from the assertion, proven in 5.11 below, that the diagram (5.10.1) commutes. \square

$$(5.10.1) \quad \begin{array}{ccc} K_n^M(E) & \xrightarrow{\lambda_E} & H^{n,n}(\text{Spec } E, \mathbb{Z}) \\ \downarrow N_{E/F} & & \downarrow N_{E/F} \\ K_n^M(F) & \xrightarrow{\lambda_F} & H^{n,n}(\text{Spec } F, \mathbb{Z}). \end{array}$$

LEMMA 5.11. *If $F \subset E$ is any finite field extension, then the diagram (5.10.1) commutes.*

PROOF. By 5.3 (3) we may assume that $[E : F] = l$ for some prime number l . Assume first that F has no extensions of degree prime to l and $[E : F] = l$. The Bass-Tate lemma (5.3) in [BT73] states that in this case $K_n^M(E)$ is generated by the symbols $a = \{a_1, \dots, a_{n-1}, b\}$ where $a_i \in F$ and $b \in E$. The properties of the norm on K_*^M and 5.6 yield:

$$\lambda_F N\{a_1, \dots, a_{n-1}, b\} = \lambda_F \{a_1, \dots, a_{n-1}, N(b)\} = [a_1 : \dots : a_{n-1}] \cdot [Nb].$$

But using the assertions of lemma 5.3 we have:

$$N \lambda_E(a) = N[a_1 : \dots : a_{n-1} : b] \stackrel{(2)}{=} [a_1 : \dots : a_{n-1}] \cdot N[b] \stackrel{(4)}{=} [a_1 : \dots : a_{n-1}] \cdot [Nb].$$

This concludes the proof in this case.

Now we use a standard reduction. For simplicity, we will write $H^{p,q}(F)$ for $H^{p,q}(\text{Spec } F, \mathbb{Z})$. If F' is a maximal prime-to- l extension of F then the kernel of $H^{n,n}(F) \rightarrow H^{n,n}(F')$ is a torsion group of exponent prime to l by (1) and (3) of 5.3. Fix $a \in K_n^M(E)$. By the above case, $t = N \lambda_E(a) - \lambda_F N(a)$ is a torsion element of $H^{n,n}(F)$, of exponent prime to l .

Since the kernel of $H^{n,n}(F) \rightarrow H^{n,n}(E)$ has exponent l , $t_E \neq 0$ if and only if $t = 0$. If E is an inseparable extension of F then by 5.3 (4) we have $t_E = l \lambda_E(a) - \lambda_E(la) = 0$. If E is separable over F then $E \otimes_F E$ is a finite product of fields E_i

with $[E_i : E] < l$. Moreover, Weil Reciprocity implies that the diagrams

$$\begin{array}{ccc}
 K_n^M(E) & \xrightarrow{\text{diag}} & \oplus K_n^M(E_i) \\
 \downarrow N_{E/F} & & \downarrow \oplus N_{E_i/E} \\
 K_n^M(F) & \longrightarrow & K_n^M(E)
 \end{array}
 \qquad
 \begin{array}{ccc}
 H^{n,n}(E) & \xrightarrow{\text{diag}} & \oplus H^{n,n}(E_i) \\
 \downarrow N_{E/F} & & \downarrow \oplus N_{E_i/E} \\
 H^{n,n}(F) & \longrightarrow & H^{n,n}(E)
 \end{array}$$

commute (see p. 387 of [BT73]). By induction on l , we have

$$t_E = \oplus N_{E_i/E} \lambda_{E_i}(a_{E_i}) - \oplus \lambda_E N_{E_i/E}(a_{E_i}) = 0.$$

Since $t_E = 0$ we also have $t = 0$. □

This completes the proof of theorem 5.1.

Part 2

Étale Motivic Theory

LECTURE 6

Étale sheaves with transfers

The goal of this lecture is to study the relations between presheaves with transfers and étale sheaves. The main result (theorem 6.17) will be that sheafification preserves transfers.

DEFINITION 6.1. A presheaf F of abelian groups on Sm/k is an **étale sheaf** if it restricts to an étale sheaf on each X in Sm/k . That is, if:

- (1) the sequence $0 \rightarrow F(X) \xrightarrow{\text{diag}} F(U) \xrightarrow{(+,-)} F(U \times_X U)$ is exact for every surjective étale morphism of smooth schemes $U \rightarrow X$;
- (2) $F(X \amalg Y) = F(X) \oplus F(Y)$ for all X and Y .

We will write $Sh_{\acute{e}t}(Sm/k)$ for the category of étale sheaves, which is a full subcategory of the category of presheaves of abelian groups.

A presheaf with transfers F is an **étale sheaf with transfers** if its underlying presheaf is an étale sheaf on Sm/k . We will write $Sh_{\acute{e}t}(Cor_k)$ for the full subcategory of **PST**(k) whose objects are the étale sheaves with transfers.

For example, we saw in lecture 2 that the étale sheaves \mathbb{Z} and \mathcal{O}^* have transfers, so they are étale sheaves with transfers. Lemma 6.2 shows that $\mathbb{Z}_{tr}(T)$ is an étale sheaf with transfers, even if T is singular (see 2.11).

LEMMA 6.2. *For any scheme T over k , $\mathbb{Z}_{tr}(T)$ is an étale sheaf.*

PROOF. Since **PST**(k) is an additive category, we have the required decomposition of $\mathbb{Z}_{tr}(T)(X \amalg Y) = \text{Hom}_{Cor_k}(X \amalg Y, T)$. To check the sheaf axiom for surjective étale maps $U \rightarrow X$, we proceed as in the proof of 3.2.

As $U \times T \rightarrow X \times T$ is flat, the pullback of cycles is well-defined and is an injection. Hence the subgroup $\mathbb{Z}_{tr}(T)(X) = Cor_k(X, T)$ of cycles on $X \times T$ injects into the subgroup $\mathbb{Z}_{tr}(T)(U) = Cor_k(U, T)$ of cycles on $U \times T$.

To see that the sequence 6.1 (1) is exact at $\mathbb{Z}_{tr}(T)(U)$, take Z_U in $Cor_k(U, T)$ whose images in $Cor_k(U \times_X U, T)$ coincide. We may assume that X and U are integral, with function fields F and L , respectively. Since $Cor_F(F, T_F)$ is the equalizer of $Cor_F(L, T_F) \rightrightarrows Cor_F(L \otimes_F L, T_F)$ by 1.11, $Z_L \in Cor_F(L, T_F)$ comes from a cycle Z_F in $Cor_F(F, T_F)$. Thus by 1.13 there is a Zariski open $V \subset X$ and a cycle Z_V in $Cor_k(V, T)$ agreeing with Z_U in $Cor(U \times_X V, T)$. Writing $Z_V = \sum n_i Z_i$, we see that we can decompose $Z_U = \sum n_i Z_i'$ so that Z_i and Z_i' agree in $Cor(U \times_X V, T)$. By restricting attention to Z_i and Z_i' , we may assume that Z_V is an elementary correspondence.

Let Z be the closure of Z_V in $X \times T$; it is irreducible and dominant over X since $Z \times_X V$ is. Since the group of cycles on $U \times T$ meeting $(U \times_X V) \times T$ injects into the group of cycles on $(U \times_X V) \times T$, we see that the lift of Z to a cycle on $U \times T$ must be Z_U . Hence the components of $Z \times_X U$ are finite over U . But by faithfully flat descent, this implies that Z is finite over X , i.e., a finite correspondence in $Cor_k(X, T)$. \square

COROLLARY 6.3. *Let F be an étale sheaf with transfers. If X is smooth, then*

$$\mathrm{Hom}_{Sh_{\acute{e}t}(Cor_k)}(\mathbb{Z}_{tr}(X), F) = \mathrm{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(X), F) = F(X).$$

COROLLARY 6.4. *For any abelian group A , the $A(n)$ are complexes of étale sheaves. If $1/n \in k$, the motivic complex of étale sheaves $\mathbb{Z}/n(1)$ is quasi-isomorphic to the étale sheaf μ_n .*

PROOF. The $\mathbb{Z}(n)$ are étale sheaves with transfers by lemmas 2.13 and 6.2, as in 3.3. We know that the $\mathbb{Z}_{tr}(T)$ are sheaves of free abelian groups. Hence $A \otimes \mathbb{Z}_{tr}(T)$ are étale sheaves. We conclude that the $A(n)$ are étale sheaves by the same argument we used for the $\mathbb{Z}(n)$. The last assertion is just a restatement of corollary 4.8 using 6.2. \square

EXERCISE 6.5. Let $\pi : X \rightarrow S$ be a finite étale map, and π_t the induced finite correspondence from S to X . If F is any étale sheaf with transfers, show that $\pi_t^* : F(X) \rightarrow F(S)$ is the étale trace map of [Mil80, V.1.12]. *Hint:* If $Y \rightarrow S$ is Galois with group G , and factors through X , then $Cor(S, X) = Cor(Y, X)^G$ by 6.2. Show that the image of π in $Cor(Y, X)$ is the sum $\sum f$ of all S -maps from $f : Y \rightarrow X$, and hence determines $\pi_t \in Cor(S, X)$.

Locally constant étale sheaves form a second important class of étale sheaves with transfers.

DEFINITION 6.6. The full subcategory Et/k of Sm/k consists of all the schemes of finite type over k which are smooth of dimension zero. Every S in Et/k is a finite disjoint union of spectra of separable field extensions of k .

It is well known (see [Mil80] and [SGA4, VIII 2.2]) that the category of étale sheaves on Et/k is equivalent to the category of discrete modules over the profinite group $\mathrm{Gal}(k_{sep}/k)$. If F corresponds to the Galois module M and $S = \mathrm{Spec}(\ell)$ then $F(S) = M^H$, where $H = \mathrm{Gal}(k_{sep}/\ell)$.

We have the following functors:

$$Sh_{\acute{e}t}(Et/k) \begin{array}{c} \xleftarrow{\pi_*} \\ \xrightarrow{\pi^*} \end{array} Sh_{\acute{e}t}(Sm/k),$$

where the restriction π_* is the right adjoint of π^* ; they are both exact functors.

DEFINITION 6.7. An étale sheaf is **locally constant** if $\pi^* \pi_* F \rightarrow F$ is an isomorphism. We will write $Sh_{\acute{e}t}^c$ for the full subcategory of $Sh_{\acute{e}t}(Sm/k)$ consisting of all locally constant sheaves.

EXERCISE 6.8. Let F be the locally constant sheaf π^*M corresponding to the G -module M . If X is connected, and l is the separable closure of k in $H^0(X, \mathcal{O}_X)$, show that $F(X) = M^H$ where $H = \text{Gal}(k_{\text{sep}}/l)$. Conclude that π_*F is the Galois module M . Note that $F(X) = M^H$ is also defined if X is normal.

LEMMA 6.9. *The functors π^* and π_* induce an equivalence between the category $Sh_{\acute{e}t}^c$ and the category of discrete modules over the profinite group $\text{Gal}(k_{\text{sep}}/k)$.*

PROOF. If M is in $Sh_{\acute{e}t}(Et/k)$, then $M \rightarrow \pi_*\pi^*M$ is an isomorphism by exercise 6.8. Thus π^* is faithful. By category theory, $\pi^*\pi_*\pi^* \cong \pi^*$, so for F locally constant we have a natural isomorphism $\pi^*\pi_*F \cong F$. \square

EXERCISE 6.10. Let L be a Galois extension of k , and let $G = \text{Gal}(L/k)$. Show that $\mathbb{Z}_{\text{tr}}(L)$ is the locally constant étale sheaf corresponding to the G -module \mathbb{Z}^G of maps $G \rightarrow \mathbb{Z}$. *Hint: $(\mathbb{Z}^G)^H = \mathbb{Z}^{G/H}$.*

LEMMA 6.11. *Any locally constant étale sheaf has a unique underlying étale sheaf with transfers.*

PROOF. Let $Z' \subset X \times Y$ be an elementary correspondence and let Z be the normalization of Z' in a normal field extension L of $F = k(X)$ containing $K = k(Z')$. If $G = \text{Gal}(L/F)$ then we also have $G = \text{Aut}_X(Z)$, and it is well known that the set $\text{Hom}_X(Z, Z')$ of maps $q : Z \rightarrow Z'$ over X is in one-to-one correspondence with the set of field maps $\text{Hom}_F(K, L)$. The cardinality of this set is the separable degree of K over F .

Let M be a Galois module, considered as a locally constant étale sheaf. It is easy to check using exercise 6.8 that $M(X)$ is isomorphic to $M(Z')^G$.

Write i for the inseparable degree of K over F . Then the transfer map $M(Y) \rightarrow M(X)$ is defined to be the composite of $M(Y) \rightarrow M(Z')$, multiplication by i , and the sum over all maps $q : Z \rightarrow Z'$ over X of $q^* : M(Z') \rightarrow M(Z)$.

The verification that this gives M the structure of a presheaf with transfers is now straightforward, and we refer the reader to [SV96, 5.17] for details. \square

It is clear that the locally constant étale sheaves form an abelian subcategory of $Sh_{\acute{e}t}(Cor_k)$, i.e., the inclusion is an exact functor.

In order to describe the relation between presheaves and étale sheaves with transfers (see 6.18), we need two preliminary results.

If $p : U \rightarrow X$ is an étale cover, we define $\mathbb{Z}_{\text{tr}}(\check{U})$ to be the Čech complex

$$\dots \xrightarrow{p_0 - p_1 + p_2} \mathbb{Z}_{\text{tr}}(U \times_X U) \xrightarrow{p_0 - p_1} \mathbb{Z}_{\text{tr}}(U) \longrightarrow 0.$$

PROPOSITION 6.12. *Let $p : U \rightarrow X$ be an étale covering of a scheme X . Then $\mathbb{Z}_{\text{tr}}(\check{U})$ is an étale resolution of the sheaf $\mathbb{Z}_{\text{tr}}(X)$, i.e., the following complex is exact as a complex of étale sheaves.*

$$\dots \xrightarrow{p_0 - p_1 + p_2} \mathbb{Z}_{\text{tr}}(U \times_X U) \xrightarrow{p_0 - p_1} \mathbb{Z}_{\text{tr}}(U) \xrightarrow{p} \mathbb{Z}_{\text{tr}}(X) \rightarrow 0$$

PROOF. As this is a complex of sheaves it suffices to verify the exactness of the sequence at every étale point. Since points in the étale topology are strictly Hensel local schemes, it is enough to prove that, for every Hensel local scheme S over k , the following sequence of abelian groups is exact.

$$(6.12.1) \quad \cdots \rightarrow \mathbb{Z}_{tr}(U)(S) \rightarrow \mathbb{Z}_{tr}(X)(S) \rightarrow 0.$$

Here S is an inverse limit of smooth schemes S_i , and by abuse of notation $\mathbb{Z}_{tr}(T)(S)$ denotes $\lim \mathbb{Z}_{tr}(T)(S_i)$.

To prove that (6.12.1) is exact we need another reduction step. Let Z be a closed subscheme of $X \times S$ which is quasi-finite over S . We write $L(Z/S)$ for the free abelian group generated by the irreducible connected components of Z which are finite and surjective over S . $L(Z/S)$ is covariantly functorial on Z with respect to morphisms of quasi-finite schemes over S . Clearly, the sequence (6.12.1) is the colimit of complexes of the form:

$$(6.12.2) \quad \cdots \rightarrow L(Z_U \times_Z Z_U/S) \rightarrow L(Z_U/S) \rightarrow L(Z/S) \rightarrow 0$$

where $Z_U = Z \times_X U$ and the limit is taken over all Z closed subschemes of $X \times S$ which are finite and surjective over S . Therefore the proof of proposition 6.12 will be completed once we show that the sequence (6.12.2) is exact for every subscheme Z of $X \times S$ which is finite and surjective over S .

Since S is Hensel local and Z is finite over S , Z is also Hensel. Therefore the covering $Z_U \rightarrow Z$ splits. Let $s_1 : Z \rightarrow Z_U$ be a splitting. We set $(Z_U)_Z^k = Z_U \times_Z \cdots \times_Z Z_U$. It is enough to check that the maps $s_k : L((Z_U)_Z^k/S) \rightarrow L((Z_U)_Z^{k+1}/S)$ are contracting homotopies where $s_k = L\left(s_1 \times_Z id_{(Z_U)_Z^k}\right)$.

This is the end of the proof of proposition 6.12. □

The proof shows that $\mathbb{Z}_{tr}(\check{U})$ is also a Nisnevich resolution of $\mathbb{Z}_{tr}(X)$, i.e., the sequence of 6.12 is also exact as a complex of Nisnevich sheaves. We can pinpoint why this proof holds in the étale topology and in the Nisnevich topology, but does not hold in the Zariski topology. This is because:

- if S is strictly Hensel local (i.e., a point in the étale topology) and Z is finite over S then Z is strictly Hensel;
- if S is Hensel local (i.e., a point in the Nisnevich topology) and Z is finite over S then Z is Hensel;
- if S is local (i.e., a point in the Zariski topology) and Z is finite over S then Z need *not* be local but will be semilocal.

EXAMPLE 6.13. Let X be a connected semilocal scheme finite over a local scheme S . X is covered by its local subschemes U_i . If X is not local, its graph Γ defines an element of $\mathbb{Z}_{tr}(X)(S)$ that cannot come from $\bigoplus \mathbb{Z}_{tr}(U_i)(S)$, because Γ does not lie in any $S \times U_i$. (By 1.4, every elementary correspondence from $\amalg U_i$ to S is an elementary correspondence from X to S , and they form a basis for the image of $\bigoplus \mathbb{Z}_{tr}(U_i)(S) \rightarrow \mathbb{Z}_{tr}(X)(S)$.) Hence $\bigoplus \mathbb{Z}_{tr}(U_i) \rightarrow \mathbb{Z}_{tr}(X)$ is not a surjection of Zariski sheaves.

We will see in 13.14 that $\text{Tot}(C_*\mathbb{Z}_{\text{tr}}(\check{U}))$ is a Zariski resolution of $C_*\mathbb{Z}_{\text{tr}}(X)$.

If $\mathcal{U} = \{U_i \rightarrow X\}$ is a Zariski covering, we can replace the infinite complex $\mathbb{Z}_{\text{tr}}(\check{U})$ of 6.12 by the bounded complex

$$\mathbb{Z}_{\text{tr}}(\check{\mathcal{U}}) : 0 \rightarrow \mathbb{Z}_{\text{tr}}(U_1 \cap \cdots \cap U_n) \rightarrow \cdots \rightarrow \bigoplus_i \mathbb{Z}_{\text{tr}}(U_i) \rightarrow 0.$$

PROPOSITION 6.14. *Let $\mathcal{U} = \{U_i \rightarrow X\}$ be a Zariski open covering of X . Then $\mathbb{Z}_{\text{tr}}(\check{\mathcal{U}})$ is an étale resolution of $\mathbb{Z}_{\text{tr}}(X)$, i.e., the following sequence is exact as a complex of étale sheaves:*

$$0 \rightarrow \mathbb{Z}_{\text{tr}}(U_1 \cap \cdots \cap U_n) \rightarrow \cdots \rightarrow \bigoplus_i \mathbb{Z}_{\text{tr}}(U_i) \rightarrow \mathbb{Z}_{\text{tr}}(X) \rightarrow 0.$$

PROOF. If $n = 2$, we apply 6.12 to $U = U_1 \amalg U_2$. Since $U \times_X U = U_1 \amalg U_2 \amalg (U_1 \cap U_2)$, we see that the image of $\mathbb{Z}_{\text{tr}}(U^{\times 3})$ in $\mathbb{Z}_{\text{tr}}(U \times_X U)$ is $\mathbb{Z}_{\text{tr}}(U_1) \oplus \mathbb{Z}_{\text{tr}}(U_2)$ in the exact complex of 6.12. It follows that $\mathbb{Z}_{\text{tr}}(\check{\mathcal{U}}) \rightarrow \mathbb{Z}_{\text{tr}}(X)$ is exact for $n = 2$. For $n > 2$, the exactness follows by induction on n . \square

EXAMPLE 6.15. If \mathcal{U} is the cover of \mathbb{P}^1 by $\mathbb{A}^1 = \text{Spec} k[t]$ and $\text{Spec} k[t^{-1}]$, and we mod out by the basepoint $t = 1$, we obtain the exact sequence

$$0 \rightarrow \mathbb{Z}_{\text{tr}}(\mathbb{G}_m) \rightarrow 2\mathbb{Z}_{\text{tr}}(\mathbb{A}^1, 1) \rightarrow \mathbb{Z}_{\text{tr}}(\mathbb{P}^1, 1) \rightarrow 0.$$

Applying C_* yields an exact sequence of complexes (see 2.14). Recalling that $C_*\mathbb{Z}_{\text{tr}}(\mathbb{A}^1, 1) \simeq 0$, we obtain quasi-isomorphisms of étale complexes (or even Nisnevich complexes)

$$C_*\mathbb{Z}_{\text{tr}}(\mathbb{P}^1, 1) \simeq C_*\mathbb{Z}_{\text{tr}}(\mathbb{G}_m)[1] = \mathbb{Z}(1).$$

LEMMA 6.16. *Let $p : U \rightarrow Y$ be an étale covering and $f : X \rightarrow Y$ a finite correspondence. Then there is an étale covering $p' : V \rightarrow X$ and a finite correspondence $f' : V \rightarrow U$ so that the following diagram commutes in Cor_k .*

$$\begin{array}{ccc} V & \xrightarrow{f'} & U \\ p' \downarrow & & \downarrow p \\ X & \xrightarrow{f} & Y \end{array}$$

PROOF. We may suppose that f is defined by the elementary correspondence $Z \subset X \times Y$. (For a general correspondence f , we take a disjoint union of such V 's.) Form the pullback $Z_U = Z \times_Y U$ inside $X \times U$. Since the projection $Z_U \rightarrow Z$ is étale and $Z \rightarrow X$ is finite, the projection splits étale-locally on X . That is, there is an étale cover $V \rightarrow X$ so that $V \times_X Z_U \rightarrow V \times_X Z$ has a section s . But then $s(V \times_X Z) \subset V \times U$ is finite over V and defines the required finite correspondence $V \rightarrow U$. \square

$$\begin{array}{ccccccc}
V \times_X Z_U & \longrightarrow & Z_U & \hookrightarrow & X \times U & \longrightarrow & U \\
\downarrow \text{split} & & \downarrow & & \downarrow & & \downarrow \\
V \times_X Z & \longrightarrow & Z & \hookrightarrow & X \times Y & \longrightarrow & Y \\
\downarrow & \searrow & \downarrow & \swarrow & \downarrow & & \downarrow \\
V & \longrightarrow & X & & & &
\end{array}$$

As in [Mil80] pp. 61–65, the inclusion $i : Sh_{\acute{e}t}(Sm/k) \rightarrow PreSh(Sm/k)$ has a left adjoint $a_{\acute{e}t}$, and $i \circ a_{\acute{e}t}$ is left exact. Hence the category of étale sheaves on Sm/k is abelian, and the functor $a_{\acute{e}t}$ is exact.

If F is a presheaf with transfers, the following theorem shows that its étale sheafification admits transfers. The same holds in the Nisnevich topology but *not* in the Zariski topology. However, we will prove later (in 22.15) that if F is a *homotopy invariant* presheaf with transfers, its Zariski sheafification admits transfers.

Recall that there is a forgetful functor $\varphi : \mathbf{PST}(k) \rightarrow PreSh(Sm/k)$.

THEOREM 6.17. *Let F be a presheaf with transfers, and write $F_{\acute{e}t}$ for $a_{\acute{e}t}\varphi F$. Then $F_{\acute{e}t}$ has a unique structure of presheaf with transfers such that $F \rightarrow F_{\acute{e}t}$ is a morphism of presheaves with transfers.*

COROLLARY 6.18. *The inclusion functor $Sh_{\acute{e}t}(Cor_k) \xrightarrow{i} \mathbf{PST}(k)$ has a left adjoint $a_{\acute{e}t}$. The category $Sh_{\acute{e}t}(Cor_k)$ is abelian, $a_{\acute{e}t}$ is exact and commutes with the forgetful functor φ to (pre)sheaves on Sm/k .*

The connections between these abelian categories, given by 6.17 and 6.18, are described by the following diagram, where the φ are (exact) forgetful functors and both functors $a_{\acute{e}t}$ are exact.

$$\begin{array}{ccc}
PreSh(Sm/k) & \xleftarrow{\varphi} & \mathbf{PST}(k) \\
\begin{array}{c} \uparrow i \\ \downarrow a_{\acute{e}t} \end{array} & & \begin{array}{c} \uparrow i \\ \downarrow a_{\acute{e}t} \end{array} \\
Sh_{\acute{e}t}(Sm/k) & \xleftarrow{\varphi} & Sh_{\acute{e}t}(Cor_k)
\end{array}$$

PROOF OF 6.17. Uniqueness. Suppose that two étale sheaves with transfers F_1 and F_2 satisfy the conditions of the theorem. We already know that $F_1(X) = F_2(X) = F_{\acute{e}t}(X)$ for all X and we just need to check that $F_1(f) = F_2(f)$ holds when $f : X \rightarrow Y$ is a morphism in Cor_k . This is given if f comes from Sm/k .

Let $y \in F_1(Y) = F_2(Y) = F_{\acute{e}t}(Y)$. Choose an étale covering $p : U \rightarrow Y$ so that $y|_U \in F_{\acute{e}t}(U)$ is the image of some $u \in F(U)$. Lemma 6.16 yields the following

diagram.

$$\begin{array}{ccc} V & \xrightarrow{f'} & U \\ p' \downarrow & & \downarrow p \\ X & \xrightarrow{f} & Y \end{array}$$

Because $y|_U$ comes from $F(U)$, we have $F_1(f')(y|_U) = F_2(f')(y|_U)$.

$$\begin{aligned} F_1(p')F_1(f)(y) &= F_1(f')F_1(p)(y) && \text{as the diagram commutes,} \\ &= F_1(f')(y|_U) && \text{as } p \text{ comes from } Sm/k, \\ &= F_2(f')(y|_U) && \text{as } y|_U \text{ comes from } F(U), \\ &= F_2(p')F_2(f)(y) && \text{as the diagram commutes,} \\ &= F_1(p')F_2(f)(y) && \text{as } p' \text{ comes from } Sm/k. \end{aligned}$$

This implies that $F_1(f) = F_2(f)$ as p' is a covering and F_1 is an étale sheaf.

Existence. We need to define a morphism $F_{\acute{e}t}(Y) \rightarrow F_{\acute{e}t}(X)$ for each finite correspondence from X to Y . We first produce a map

$$F_{\acute{e}t}(Y) \rightarrow \text{Hom}_{Sh}(\mathbb{Z}_{tr}(Y), F_{\acute{e}t})$$

natural in Cor_k and compatible with $F(Y) \rightarrow \text{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(Y), F)$.

For all $y \in F_{\acute{e}t}(Y)$ there is an étale covering $p : U \rightarrow Y$ and an element $u \in F(U)$ so that y and u agree in $F_{\acute{e}t}(U)$. By representability (see 2.8), u determines a morphism $\mathbb{Z}_{tr}(U) \rightarrow F$ of presheaves with transfers. By shrinking U , we may arrange that the difference map sends u to zero in $F(U \times_Y U)$. A chase in the commutative diagram below (where U_Y^2 denotes $U \times_Y U$) will produce the map of sheaves $[y] : \mathbb{Z}_{tr}(Y) \rightarrow F_{\acute{e}t}$. The top row is exact by 6.12.

$$\begin{array}{ccccc} 0 \rightarrow \text{Hom}_{Sh}(\mathbb{Z}_{tr}(Y), F_{\acute{e}t}) & \longrightarrow & \text{Hom}_{Sh}(\mathbb{Z}_{tr}(U), F_{\acute{e}t}) & \longrightarrow & \text{Hom}_{Sh}(\mathbb{Z}_{tr}(U_Y^2), F_{\acute{e}t}) \\ & & \uparrow & & \uparrow \\ & & \text{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(U), F) & \longrightarrow & \text{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(U_Y^2), F) \end{array}$$

It is easy to see that $[y]$ is independent of the choice of U and u . We can now define a pairing $Cor(X, Y) \otimes F_{\acute{e}t}(Y) \rightarrow F_{\acute{e}t}(X)$. Let f be a correspondence from X to Y and $y \in F_{\acute{e}t}(Y)$. By the map just described, y induces a morphism of sheaves $[y] : \mathbb{Z}_{tr}(Y) \rightarrow F_{\acute{e}t}$. Consider the composition:

$$\mathbb{Z}_{tr}(X) \xrightarrow{f} \mathbb{Z}_{tr}(Y) \xrightarrow{[y]} F_{\acute{e}t}.$$

Hence there is a map $\mathbb{Z}_{tr}(X)(X) \rightarrow F_{\acute{e}t}(X)$. The image of the identity map will be the pairing of f and y . \square

We conclude with an application of these ideas to homological algebra.

PROPOSITION 6.19. *The abelian category $Sh_{\acute{e}t}(Cor_k)$ has enough injectives.*

PROOF. The category $\mathcal{S} = Sh_{\acute{e}t}(Cor_k)$ has products and filtered direct limits are exact, because this is separately true for presheaves with transfers and for étale sheaves. That is, \mathcal{S} satisfies axioms (AB5) and (AB3^{*}). By 6.3, the family of sheaves $\mathbb{Z}_{tr}(X)$ is a family of generators of \mathcal{S} . It is well known (see [Gro57, 1.10.1]) that this implies that \mathcal{S} has enough injectives. \square

EXAMPLE 6.20. Let F be an étale sheaf with transfers. We claim that the terms $E^n(F)$ in its canonical flasque resolution (as an étale sheaf, see [Mil80] p. 90) are actually étale sheaves with transfers. For this it suffices to consider $E = E^0(F)$. Fix an algebraic closure \bar{k} of k . For every X we define:

$$E(X) = \prod_{\bar{x} \in X(\bar{k})} F_{\bar{x}},$$

where $X(\bar{k})$ is the set of \bar{k} -points of X , and $F_{\bar{x}}$ denotes the fiber of F at \bar{x} . If $U \rightarrow X$ is étale, $E(U)$ is the product $\prod F_{\bar{x}}$ over $\bar{x} \in U(\bar{k})$. From this it follows that E is an étale sheaf, not only on X but on the big étale site of Sm/k . It is also easy to see that $F(X) \rightarrow E(X)$ is an injection.

In addition, E is a presheaf with transfers and $F \rightarrow E$ is a morphism in **PST**. For if $Z \subset X \times Y$ is an elementary correspondence from X to Y , we define the transfer $E(Y) \rightarrow E(X)$

$$E(Y) = \prod_{\bar{y} \in Y(\bar{k})} F_{\bar{y}} \rightarrow \prod_{\bar{x} \in X(\bar{k})} F_{\bar{x}} = E(X)$$

by stating that the component for $\bar{x} \in X(\bar{k})$ is the sum of the induced transfers $F_{\bar{y}} \rightarrow F_{\bar{x}}$, taken over all $\bar{y} \in Y(\bar{k})$ such that $z = (\bar{x}, \bar{y}) \in Z(\bar{k})$. To see that $F \rightarrow E$ is a morphism in **PST**, we may take X to be strictly Hensel local, so $F(X) = E(X)$. Since this forces Y to also be strictly Hensel semilocal, so $F(Y) = E(Y)$, this is a tautology.

The same construction works in the Nisnevich topology, letting $E(X)$ be the product over all closed points $x \in X$ of $F(\text{Spec } \mathcal{O}_{X,x}^h)$ (see 13.3). However, example 6.13 shows that it does not work in the Zariski topology, because the transfer $E(X) \rightarrow E(S)$ need not factor through the sum of the $E(U_i)$.

LEMMA 6.21. *If F is any étale sheaf with transfers, then its cohomology presheaves $H_{\acute{e}t}^n(-, F)$ are presheaves with transfers.*

PROOF. The canonical flasque resolution $F \rightarrow E^*(F)$ of 6.20 is a resolution of sheaves with transfers. Since the forgetful functor from **PST**(k) to presheaves is exact, and $H^n(-, F)$ is the cohomology $E^*(F)$ as a presheaf, we see that $H^n(-, F)$ is also the cohomology of $E^*(F)$ in the abelian category **PST**(k). \square

EXAMPLE 6.22. By 2.4, $F = \mathbb{G}_m$ is an étale sheaf with transfers. By 6.21, both the Picard group $\text{Pic}(X) = H_{\acute{e}t}^1(X, \mathbb{G}_m)$ and the cohomological Brauer group $Br^l(X) = H_{\acute{e}t}^2(X, \mathbb{G}_m)_{tors}$ are presheaves with transfers.

LEMMA 6.23. *For any $F \in Sh_{\acute{e}t}(Cor_k)$ and any smooth X and $i \in \mathbb{Z}$ we have:*

$$\mathrm{Ext}_{Sh_{\acute{e}t}(Cor_k)}^i(\mathbb{Z}_{tr}(X), F) = H_{\acute{e}t}^i(X, F).$$

PROOF. The case $i = 0$ is $\mathrm{Hom}(\mathbb{Z}_{tr}(X), F) = F(X)$; this is 6.3. For $i > 0$ it suffices to show that if F is an injective étale sheaf with transfers then $H^i(X, F)$ is zero. Consider the canonical flasque resolution $E^*(F)$ of example 6.20. Since F is injective, the canonical inclusion $F \rightarrow E^0$ must split, i.e., F is a direct factor of E^0 in $Sh_{\acute{e}t}(Cor_k)$. Since $H_{\acute{e}t}^i(X, F)$ is a direct summand of $H^i(X, E^0)$, it must vanish for $i > 0$. \square

If we restrict to the category $Sh_{\acute{e}t}(Cor_k, R)$ of étale sheaves of R -modules with transfers, $E^0(F)$ is a flasque sheaf of R -modules with transfers by 6.20. The proof of 6.23 goes through word for word to prove the following variation.

PORISM 6.24. *For any $F \in Sh_{\acute{e}t}(Cor_k, R)$ and any smooth X and $i \in \mathbb{Z}$:*

$$\mathrm{Ext}_{Sh_{\acute{e}t}(Cor_k, R)}^i(R_{tr}(X), F) = H_{\acute{e}t}^i(X, F).$$

The same proof also shows that lemma 6.23 and porism 6.24 hold for the Nisnevich topology (see 13.4). See [TriCa, 3.1.8] for an alternative proof.

EXERCISE 6.25. Let K be any complex of étale sheaves of R -modules with transfers. Show that its hyperext and hypercohomology agree in the sense that for any smooth X and $i \in \mathbb{Z}$:

$$\mathrm{Ext}^i(R_{tr}(X), K) \cong \mathbb{H}_{\acute{e}t}^i(X, K).$$

(For simplicity, the reader may assume that $\mathrm{cd}_R(k) < \infty$.)

If one is interested in extending the constructions of this lecture to possibly singular schemes, it would be useful to assume that k admits resolution of singularities and use the cdh topology, which we will introduce in lecture 12.

LECTURE 7

The relative Picard group and Suslin's Rigidity Theorem

In this lecture we introduce the relative Picard group $\text{Pic}(\bar{X}, X_\infty)$. When \bar{X} is a good compactification of X over S , its elements determine maps $F(X) \rightarrow F(S)$ for every homotopy invariant F . This pairing will be used to prove Suslin's Rigidity Theorem 7.20.

Recall from 1A.9 and 1A.10 that if S is a smooth connected scheme and $p : X \rightarrow S$ a smooth morphism then we write $c(X/S, 0)$ for the free abelian group generated by the irreducible closed subsets of X which are finite and surjective over S . In this lecture we will write $C_0(X/S)$ for $c(X/S, 0)$.

By 1A.12, given a map $S' \rightarrow S$, there is a map $C_0(X/S) \rightarrow C_0(X \times_S S'/S')$, induced from

$$C_0(X/S) \hookrightarrow \mathbb{Z}_{tr}(X)(S) = \text{Cor}_k(S, X).$$

DEFINITION 7.1. We define $H_0^{sing}(X/S)$ to be the cokernel of the map

$$C_0(X \times \mathbb{A}^1/S \times \mathbb{A}^1) \xrightarrow{\partial_0 - \partial_1} C_0(X/S)$$

where ∂_i is induced by “ $t = i$ ” : $\text{Spec } k \rightarrow \mathbb{A}_k^1$.

EXAMPLE 7.2. If $X = Y \times_k S$ then $C_0(X/S) = \text{Cor}(S, Y) = \mathbb{Z}_{tr}(Y)(S)$. In addition, $X \times \mathbb{A}^1 = Y \times_k (S \times \mathbb{A}^1)$ and the following diagram commutes:

$$\begin{array}{ccc} C_0(X \times \mathbb{A}^1/S \times \mathbb{A}^1) & \longrightarrow & C_0(X/S) \\ \downarrow = & & \downarrow = \\ \mathbb{Z}_{tr}(Y)(S \times \mathbb{A}^1) & \longrightarrow & \mathbb{Z}_{tr}(Y)(S). \end{array}$$

Taking cokernels, we conclude (using 2.27) that:

$$H_0^{sing}(Y \times S/S) = H_0 C_* \mathbb{Z}_{tr}(Y)(S) = \text{Cor}(S, Y)/\mathbb{A}^1\text{-homotopy}.$$

In particular, this implies that two elements of $\text{Cor}(S, Y)$ are \mathbb{A}^1 -homotopic exactly when they agree in $H_0^{sing}(Y \times S/S)$.

If $S = \text{Spec } k$ then $H_0^{sing}(X/S)$ is the cokernel $H_0^{sing}(X/k)$ of $\mathbb{Z}_{tr}(X)(\mathbb{A}^1) \rightarrow \mathbb{Z}_{tr}(X)(S)$ discussed in exercise 2.21, because $C_0(X/S) = \mathbb{Z}_{tr}(X)(\text{Spec } k)$. Also by 2.21, there is a natural surjection $H_0^{sing}(X/S) \rightarrow CH_0(X)$. If X is projective, this surjection is an isomorphism.

EXAMPLE 7.3. If $S = \text{Spec } k$, then 7.16 below shows that $H_0^{sing}(\mathbb{P}^1/S) = H_0^{sing}(\mathbb{A}^1/S) = \mathbb{Z}$ but $H_0^{sing}(\mathbb{A}^1 - 0/S) = \mathbb{Z} \oplus k^*$.

REMARK 7.4. In [SV96] the groups $H_*^{sing}(X/S)$ are defined to be the homology of the evident chain complex $C_*(X/S)$ with

$$C_n(X/S) = C_0(X \times \Delta^n / S \times \Delta^n).$$

We will consider the singular homology $H_*^{sing}(X/S)$ in lecture 10 below when $S = \text{Spec } k$, and $C_*(X/S) = C_*\mathbb{Z}_{tr}(X)(S)$.

Let F be a **PST**. The map $\text{Tr} : C_0(X/S) \otimes F(X) \rightarrow F(S)$ is defined to be the inclusion $C_0(X/S) \subset \text{Cor}_k(S, X)$ (see 1A.12) followed by evaluation on $F(X)$.

$$\begin{array}{ccc} C_0(X/S) \otimes F(X) & \xrightarrow{\text{Tr}} & F(S) \\ \downarrow & \nearrow \text{evaluate} & \\ \text{Cor}_k(S, X) \otimes F(X) & & \end{array}$$

LEMMA 7.5. *If F is a homotopy invariant presheaf with transfers then the map Tr factors through $H_0^{sing}(X/S) \otimes F(X) \rightarrow F(S)$.*

PROOF. Since $F(X) = F(X \times \mathbb{A}^1)$, we have a diagram

$$\begin{array}{ccc} C_0(X \times \mathbb{A}^1 / S \times \mathbb{A}^1) \otimes F(X) & \xrightarrow{\text{Tr}} & F(S \times \mathbb{A}^1) \\ \downarrow \partial_0 - \partial_1 & & \downarrow i_0 - i_1 = 0 \\ C_0(X/S) \otimes F(X) & \xrightarrow{\text{Tr}} & F(S). \quad \square \end{array}$$

EXAMPLE 7.6. If $\sigma : S \rightarrow X$ is a section of p , regarded as an element of $H_0^{sing}(X/S)$, then $\text{Tr}(\sigma, -)$ is the usual map $\sigma^* : F(X) \rightarrow F(S)$.

REMARK 7.7. The pairing $H_0^{sing}(X/S) \otimes F(X) \rightarrow F(S)$ is fundamental. It can be defined more generally for homotopy invariant presheaves equipped only with transfer maps $\text{Tr}_D : F(X) \rightarrow F(S)$ for any relative smooth curve X/S and any effective divisor $D \subset X$ which is finite and surjective over S , such that the transfer maps form a “pseudo pretheory”. This construction applies to the K -theory presheaves $K_n(X)$, equipped with the transfer maps of exercise 2.7, even though these are not presheaves with transfers.

In order to compute $H_0^{sing}(X/S)$, it is useful to embed X in a slightly larger scheme \bar{X} .

DEFINITION 7.8. A smooth curve $p : X \rightarrow S$ admits a **good compactification** \bar{X} if it factors as:

$$\begin{array}{ccc} X & \xrightarrow{j} & \bar{X} \\ & \searrow p & \downarrow \bar{p} \\ & & S \end{array}$$

where j is an open embedding, \bar{X} is a proper normal but not necessarily smooth curve over S and $Y = \bar{X} - X$ has an affine open neighborhood in \bar{X} .

If S is affine, for example, then $X = \mathbb{A}^1 \times S$ admits $\mathbb{P}^1 \times S$ as a good compactification. Similarly, if C is any smooth affine curve over k then $C \times S \rightarrow S$ admits $\bar{C} \times S$ as a good compactification. The following result implies that every point x of every X has an open neighborhood U which has a good compactification over a generic projection $X \rightarrow \mathbb{A}^{l-1}$.

LEMMA 7.9. *Let $p : X \rightarrow \mathbb{A}^l$ be an étale map. If k is infinite, there exists a linear projection $\mathbb{A}^l \rightarrow \mathbb{A}^{l-1}$ so that the composition $X \rightarrow \mathbb{A}^{l-1}$ is a curve with a good compactification.*

PROOF. There is an open $U \subset \mathbb{A}^l$ so that X is quasi-finite and surjective over U . Choose a linear projection $\mathbb{A}^l \rightarrow \mathbb{A}^{l-1}$ so that the restriction to $\mathbb{A}^l - U$ is finite; \mathbb{A}^l has good compactification $Y = \mathbb{P}^1 \times \mathbb{A}^{l-1}$. By Zariski's Main Theorem (as formulated in [EGA4, 8.12.6]), the map $X \rightarrow Y$ may be factored as an open immersion $X \hookrightarrow \bar{X}$ followed by a finite map $\bar{p} : \bar{X} \rightarrow Y$. Replacing \bar{X} by its normalization, we may assume that \bar{X} is normal. Note that \bar{p} is an affine map. Since Y is a good compactification of U , \bar{X} is a good compactification of X . \square

DEFINITION 7.10. If $Y \hookrightarrow \bar{X}$ is closed we set $\mathbb{G}_{\bar{X}, Y} = \text{Ker}(\mathcal{O}_{\bar{X}}^* \rightarrow i_* \mathcal{O}_Y^*)$. The **relative Picard group** is defined to be:

$$\text{Pic}(\bar{X}, Y) = H_{\text{Zar}}^1(\bar{X}, \mathbb{G}_{\bar{X}, Y}).$$

By [Mil80] p. 124, we also have $\text{Pic}(\bar{X}, Y) = H_{\text{ét}}^1(\bar{X}, \mathbb{G}_{\bar{X}, Y})$.

By [SV96, 2.1], the elements of $\text{Pic}(\bar{X}, Y)$ are the isomorphism classes (\mathcal{L}, t) of line bundles \mathcal{L} on \bar{X} with a trivialization t on Y . The group operation is \otimes , i.e., $(\mathcal{L}, t) \otimes (\mathcal{L}', t') = (\mathcal{L} \otimes \mathcal{L}', t \otimes t')$.

REMARK 7.11. For $\bar{X} = S \times \mathbb{P}^1$ and $Y = S \times \{0, \infty\}$, the “stalk” $(i^* \mathbb{G}_{\bar{X}, Y})(Y)$ of $\mathbb{G}_{\bar{X}, Y}$ at Y is the group $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)(S)$ of lecture 4.

The cohomology of $\mathcal{O}^* \rightarrow i_* \mathcal{O}_Y^*$ yields the exact sequence

$$\mathcal{O}^*(\bar{X}) \rightarrow \mathcal{O}^*(Y) \rightarrow \text{Pic}(\bar{X}, Y) \rightarrow \text{Pic}(\bar{X}) \rightarrow \text{Pic}(Y).$$

Comparing this exact sequence for \bar{X} and $\bar{X} \times \mathbb{A}^1$ yields:

COROLLARY 7.12. *If \bar{X} is a normal scheme and Y is reduced, we have:*

$$\text{Pic}(\bar{X}, Y) = \text{Pic}(\bar{X} \times \mathbb{A}^1, Y \times \mathbb{A}^1).$$

Let us write j for the open embedding of $X = \bar{X} - Y$ into \bar{X} .

LEMMA 7.13. *If $1/n \in k$, there is a natural injection*

$$\text{Pic}(\bar{X}, Y)/n \hookrightarrow H_{\text{ét}}^2(\bar{X}, j_! \mu_n).$$

PROOF. By Kummer Theory we have an exact sequence of étale sheaves:

$$(7.13.1) \quad 0 \longrightarrow j_! \mu_n \longrightarrow \mathbb{G}_{\bar{X}, Y} \xrightarrow{n} \mathbb{G}_{\bar{X}, Y} \longrightarrow 0.$$

Applying étale cohomology yields:

$$H^1(\bar{X}, j_! \mu_n) \longrightarrow H^1(\bar{X}, \mathbb{G}_{\bar{X}, Y}) \xrightarrow{n} H^1(\bar{X}, \mathbb{G}_{\bar{X}, Y}) \longrightarrow H_{\text{ét}}^2(\bar{X}, j_! \mu_n).$$

But the middle groups are both $\text{Pic}(\bar{X}, Y)$. □

EXAMPLE 7.14. Suppose that $S = \text{Spec } k$ and k is algebraically closed. If \bar{X} is a smooth connected curve, then $\text{Pic}(\bar{X}, Y)$ is an extension of $\text{Pic}(\bar{X})$ by a finite product of $|Y| - 1$ copies of k^* . Hence $\text{Pic}(\bar{X}, Y)/n \cong H_{\text{ét}}^2(\bar{X}, \mu_n) \cong \mathbb{Z}/n$.

Recall that $C_0(X/S)$ is generated by closed subsets Z of X which are finite and surjective over S . Because X is smooth, each such subset is an effective Cartier divisor on \bar{X} , and has an associated line bundle \mathcal{L} equipped with a canonical map $\mathcal{O} \rightarrow \mathcal{L}$. This map gives a trivialization of \mathcal{L} on $\bar{X} - Z$, which is a neighborhood of Y . Thus a good compactification \bar{X} induces a homomorphism

$$C_0(X/S) \rightarrow \text{Pic}(\bar{X}, Y).$$

When Y lies in an affine open neighborhood, this map is onto because every trivialization on Y extends to a neighborhood of Y .

EXERCISE 7.15. In this exercise we make the lifting to $C_0(X/S)$ explicit. Suppose that \mathcal{L} is a line bundle on \bar{X} with a fixed trivialization t on an open neighborhood U of Y . Show that t gives a canonical isomorphism of \mathcal{L} with a Cartier divisor $\mathcal{L}(D)$, i.e., an invertible subsheaf of the sheaf \mathcal{K} of total quotient rings of \mathcal{O} . (See [Har77, II.6].) Show that $\mathcal{L}(D)$ comes from a Weil divisor $D = \sum n_i Z_i$ on \bar{X} with the Z_i supported on $\bar{X} - U$. Then show that the map $C_0(X/S) \rightarrow \text{Pic}(\bar{X}, Y)$ sends $\sum n_i Z_i$ to (\mathcal{L}, t) .

Because $C_1(X/S) \rightarrow \text{Pic}(\bar{X}, Y)$ factors through $\text{Pic}(\bar{X} \times \mathbb{A}^1, Y \times \mathbb{A}^1)$, corollary 7.12 shows that $C_0(X/S) \rightarrow \text{Pic}(\bar{X}, Y)$ induces a homomorphism

$$H_0^{\text{sing}}(X/S) \rightarrow \text{Pic}(\bar{X}, Y).$$

THEOREM 7.16. *Let S be a smooth scheme. If $p : X \rightarrow S$ is a smooth quasi-affine curve with a good compactification (\bar{X}, Y) , then:*

$$H_0^{\text{sing}}(X/S) \xrightarrow{\cong} \text{Pic}(\bar{X}, Y).$$

PROOF. The kernel of $C_0(X/S) \rightarrow \text{Pic}(\bar{X}, Y)$ consists of $f \in K(\bar{X})$ which are defined and equal to 1 on Y . Since X is quasi-affine over S , Y contains at least one point in every irreducible component of every fiber of \bar{X} over S . Therefore the divisor D of $tf + (1-t)$ defines an element of $C_0(X \times \mathbb{A}^1/S \times \mathbb{A}^1)$ with $\partial_0 D = 0$ and $\partial_1 D = (f)$. Hence (f) represents 0 in $H_0^{\text{sing}}(X/S)$. This proves that the map $H_0^{\text{sing}}(X/S) \rightarrow \text{Pic}(\bar{X}, Y)$ is an injection, hence an isomorphism. □

Theorem 7.16 also holds when X is not quasi-affine over S , but the proof is more involved.

COROLLARY 7.17. *If F is a homotopy invariant presheaf with transfers, there is a pairing*

$$\mathrm{Pic}(\bar{X}, Y) \otimes F(X) \rightarrow F(S).$$

EXAMPLE 7.18. If X is a smooth curve over k and $1/n \in k$, then any two geometric points $x, x' : \mathrm{Spec} \bar{k} \rightarrow X$ induce the same map $F(X) \rightarrow F(\mathrm{Spec} \bar{k})$. Here F is any homotopy invariant presheaf with transfers satisfying $nF = 0$. Indeed, $[x] = [x']$ in $\mathrm{Pic}(\bar{X}, Y)/n$ by example 7.14. This phenomenon is known as ‘‘rigidity,’’ and is a simple case of theorem 7.20 below.

COROLLARY 7.19. *Let $p : X \rightarrow S$ be a smooth curve with a good compactification. Assume that S is Hensel local and let $X_0 \rightarrow S_0$ be the closed fiber of p . Then for every n prime to $\mathrm{char} k$ the following map is injective:*

$$H_0^{\mathrm{sing}}(X/S)/n \rightarrow H_0^{\mathrm{sing}}(X_0/S_0)/n.$$

PROOF. Kummer Theory yields the exact sequence 7.13.1 of étale sheaves, and similarly for (\bar{X}_0, Y_0) . Applying étale cohomology yields:

$$\begin{array}{ccccccc} H^1(\bar{X}, j_! \mu_n) & \longrightarrow & H^1(\bar{X}, \mathbb{G}_{\bar{X}, Y}) & \xrightarrow{n} & H^1(\bar{X}, \mathbb{G}_{\bar{X}, Y}) & \longrightarrow & H_{\acute{e}t}^2(\bar{X}, j_! \mu_n) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \cong \\ H^1(\bar{X}_0, j_! \mu_n) & \longrightarrow & H^1(\bar{X}_0, \mathbb{G}_{\bar{X}, Y}) & \xrightarrow{n} & H^1(\bar{X}_0, \mathbb{G}_{\bar{X}, Y}) & \longrightarrow & H_{\acute{e}t}^2(\bar{X}_0, j_! \mu_n). \end{array}$$

Since $H^2(\bar{X}, j_! \mu_n) = H_c^2(X, \mu_n)$, the right vertical map is an isomorphism by proper base change with compact supports (see [Mil80, VI.3.2]). We have a diagram:

$$\begin{array}{ccc} \mathrm{Pic}(\bar{X}, Y)/n & \hookrightarrow & H_{\acute{e}t}^2(\bar{X}, j_! \mu_n) \\ \downarrow & & \downarrow \cong \\ \mathrm{Pic}(X_0, Y_0)/n & \hookrightarrow & H_{\acute{e}t}^2(\bar{X}_0, j_! \mu_n). \end{array}$$

Corollary 7.19 now follows from theorem 7.16. \square

It follows from 6.8 that every locally constant étale sheaf F is homotopy invariant, because $H^0(X \times \mathbb{A}^1, \mathcal{O}) \cong H^0(X, \mathcal{O}) \otimes_k k[t]$. The following result shows that the converse is true for torsion sheaves. (Cf. [SV96, 4.5].)

THEOREM 7.20. (*Suslin’s ‘‘Rigidity Theorem’’*) *Let F be a homotopy invariant presheaf with transfers, such that the groups $F(X)$ are torsion of exponent prime to $\mathrm{char} k$. Then $F_{\acute{e}t}$ is locally constant.*

PROOF. Let $F_0 = \pi_* \pi^*(F)$ be the locally constant sheaf for the group $M = F(k_{\mathrm{sep}})$. We want to show that the adjunction $F_0 \rightarrow F$ is an isomorphism of étale sheaves. It suffices to check this on stalks. Since $\mathcal{O}_{X, x}^{\mathrm{sh}}$ contains a separable closure of k , we may assume that k is separably closed. In this case 7.20 asserts that $F_{\acute{e}t}$

is the constant sheaf for the group $M = F(\text{Spec } k)$. Since X is smooth at x , $\mathcal{O}_{X,x}^{sh}$ is isomorphic to the Henselization of \mathbb{A}^l at $\{0\}$. Thus the Rigidity Theorem is a consequence of proposition 7.21 below. \square

PROPOSITION 7.21. *Let S_l be the Henselization at $\{0\}$ in \mathbb{A}^l over a separably closed field k . Assume that F is as in 7.20. Then $F(S_l) = F(\text{Spec } k)$.*

PROOF. The hypothesis on F is inherited by $F(X)_n = \{x \in F(X) : nx = 0\}$. Therefore we may assume that F has exponent n for some prime n .

We use the following sequence of inclusions:

$$\text{Spec } k = S_0 \subset \cdots \subset S_{l-1} \xrightarrow{i} S_l.$$

By induction on l , it is enough to prove that the map $F(i) : F(S_l) \rightarrow F(S_{l-1})$ is an isomorphism. For this it suffices to prove that $F(i)$ is an injection, because it is split by the projection π

$$S_{l-1} \xrightleftharpoons[i]{\pi} S_l.$$

But $F(S_l) = \text{colim}_{(X,x_0) \rightarrow (\mathbb{A}^l, 0)} F(X)$ where the colimit is taken over all diagrams:

$$S_l \xrightarrow{\pi} X \xrightarrow{p} \mathbb{A}^l.$$

It suffices to show for every X that if $\varphi \in F(X)$ has $i_l^* \pi^* \varphi = 0$ then $\pi^* \varphi = 0$. By lemma 7.9 there is a curve $X \rightarrow \mathbb{A}_{l-1}$ with a good compactification. Let X' be the pullback in the following diagram.

$$\begin{array}{ccccc} S_{l-1} & \xrightarrow{i_l} & S_l & & \\ & & \searrow^{s_1} & \searrow^{\pi} & \\ & & X' & \xrightarrow{q} & X \\ & \searrow^{Id} & \downarrow & & \downarrow \\ & & S_l & \xrightarrow{\pi_l} & S_{l-1} & \longrightarrow & \mathbb{A}^{l-1} \end{array}$$

The maps π and $\pi i_l \pi_l : S_l \rightarrow X$ induce two sections $s_1, s_2 : S_l \rightarrow X'$ of $X' \rightarrow S_l$ which agree on the closed fiber $X_0 = X \times_S S_0$. Given $\varphi \in F(X)$ we need to show that $\pi_l^* i_l^* \pi^* \varphi = \pi^* \varphi$. But $\pi^* \varphi = s_1^* q^*(\varphi)$ and $\pi_l^* i_l^* \pi^* \varphi = s_2^* q^*(\varphi)$. The s_i coincide on the closed point of S_l by construction. So we are left to prove that $s^*(\psi) = (s')^*(\psi)$ for all $\psi \in F(X')$ and any $s, s' : S_l \rightarrow X'$ with $s_0 = s'_0$. Consider the following

diagram:

$$\begin{array}{ccccc}
 (\Gamma_s - \Gamma_{s'}) \otimes \psi & \xrightarrow{\hspace{10em}} & s^*(\psi) - s'^*(\psi) & & \\
 C_0(X'/S_l) \otimes F(X') & \longrightarrow & H_0(X'/S_l) \otimes F(X') & \xrightarrow{\text{Tr}} & F(S_l) \\
 & & \downarrow & & \downarrow \\
 & & H_0(X'_0/S_0) \otimes F(X') & \xrightarrow{\text{Tr}} & F(S_0).
 \end{array}$$

By assumption, the element $(\Gamma_s - \Gamma_{s'}) \otimes \psi$ in the top left group goes to zero in $H_0(X'_0/S_0) \otimes F(X')$. Hence it vanishes in $H_0(X'/S_l) \otimes F(X')$ by the immersion of $H_0(X'/S)/n$ in $H_0(X'_0/S_0)/n$ of 7.19. Therefore $s^*(\psi) - s'^*(\psi)$ vanishes in $F(S_l)$. \square

We conclude this lecture with a description of the behavior of the relative Picard group for finite morphisms. We will need this description in the proof of 21.9.

DEFINITION 7.22. Let (\bar{Y}, Y_∞) and (\bar{X}, X_∞) be two good compactifications, say of Y and X , respectively. Any finite map $f : \bar{Y} \rightarrow \bar{X}$ which restricts to a map $f : Y \rightarrow X$, yields a map $f_* : \mathcal{O}^*(Y_\infty) \rightarrow \mathcal{O}^*(X_\infty)$ constructed as follows.

Consider $\alpha \in \mathcal{O}^*(Y_\infty)$. We may extend α to $\tilde{\alpha} \in \mathcal{O}^*(U)$ where U is an affine open neighborhood of Y_∞ . Since f is finite, we may assume that $U = f^{-1}(V)$, where V is an open neighborhood of X_∞ . Since V is normal, there is a norm map $N : \mathcal{O}^*(U) \rightarrow \mathcal{O}^*(V)$ (see example 2.4). We define $f_*(\alpha) = N(\tilde{\alpha})|_{X_\infty}$. By 7.23 below, $f_*(\alpha)$ is independent of the choice of the extension $\tilde{\alpha}$.

EXERCISE 7.23. Let $f : U \rightarrow V$ be a finite morphism of normal schemes and let $Z \subset V$ be a reduced closed subscheme. If $\alpha \in \mathcal{O}^*(U)$ and $\alpha = 1$ on the reduced closed subscheme $f^{-1}(Z)$, show that $N(\alpha) = 1$ on Z .

LEMMA 7.24. Let (\bar{Y}, Y_∞) and (\bar{X}, X_∞) be good compactifications of Y and X , respectively. Let f be a finite map $f : \bar{Y} \rightarrow \bar{X}$ which restricts to a map $f : Y \rightarrow X$. Then the following diagram is commutative:

$$\begin{array}{ccccc}
 \mathcal{O}^*(Y_\infty) & \longrightarrow & \text{Pic}(\bar{Y}, Y_\infty) & \xrightarrow{\cong} & H_0(Y/S) \\
 \downarrow f_* & & & & \downarrow \\
 \mathcal{O}^*(X_\infty) & \longrightarrow & \text{Pic}(\bar{X}, X_\infty) & \xrightarrow{\cong} & H_0(X/S),
 \end{array}$$

where f_* was defined in 7.22 and the right vertical map is induced by the push-forward of cycles.

PROOF. Choose $\alpha \in \mathcal{O}^*(Y_\infty)$ and extend it to a rational function t on \bar{Y} which is regular in a neighborhood of the form $f^{-1}(V)$. By definition, $f_*(\alpha)$ extends to the regular function $N(t)$ on V . The horizontal maps send α and $f_*(\alpha)$ to $(\mathcal{O}_{\bar{Y}}, \alpha)$ and $(\mathcal{O}_{\bar{X}}, f_*\alpha)$. Let D and D' be the Weil divisors on \bar{Y} and \bar{X} associated to t and $N(t)$, respectively. We may regard D and D' as classes in $C_0(Y/S)$ and $C_0(X/S)$. By 7.15, D and D' represent the images of $(\mathcal{O}_{\bar{Y}}, \alpha)$ and $(\mathcal{O}_{\bar{X}}, f_*\alpha)$ in $H_0(Y/S)$ and

$H_0(X/S)$, respectively. The right vertical map sends D to D' because $D' = \text{div}(Nt)$ is the push-forward of $D = \text{div}(t)$ (see [Ful84, 1.4]). \square

LECTURE 8

Derived tensor products

The goal of this lecture is to define a tensor product on the derived category of étale sheaves with transfers, starting with the tensor product $X \otimes Y = X \times Y$ on Cor_k defined in 1.9. For this we first need to build a total tensor product on the category $\mathbf{PST}(k)$, and this construction makes sense in somewhat greater generality.

Let \mathcal{A} be a small additive category. We define $\mathbb{Z}(\mathcal{A})$ to be the category of all additive presheaves on \mathcal{A} , i.e., all contravariant additive functors $F : \mathcal{A} \rightarrow \mathbf{Ab}$. It is an abelian category. The Yoneda embedding $h : \mathcal{A} \rightarrow \mathbb{Z}(\mathcal{A})$ allows us to define the additive category \mathcal{A}^\oplus as the closure of \mathcal{A} under infinite direct sums in $\mathbb{Z}(\mathcal{A})$. If X_i are in \mathcal{A} , we will consider $X = \bigoplus X_i$ to be the object of \mathcal{A}^\oplus corresponding to the presheaf $h_X = \bigoplus h_{X_i}$ in $\mathbb{Z}(\mathcal{A})$.

More generally, if R is a ring, we define $R(\mathcal{A})$ to be the (abelian) category of all additive functors $F : \mathcal{A} \rightarrow R\text{-mod}$. By abuse of notation, we will write h_X for the functor $A \mapsto R \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{A}}(A, X)$ and call it “representable”.

LEMMA 8.1. *Every representable presheaf h_X is a projective object of $R(\mathcal{A})$, every projective object of $R(\mathcal{A})$ is a direct summand of a direct sum of representable functors, and every F in $R(\mathcal{A})$ has a projective resolution.*

PROOF. Since $\text{Hom}_{R(\mathcal{A})}(h_X, F) \cong F(X)$, each h_X is a projective object in $R(\mathcal{A})$. Moreover every F in $R(\mathcal{A})$ is a quotient of some h_X , $X \in \mathcal{A}^\oplus$, because of the natural surjection

$$\bigoplus_{X \text{ in } \mathcal{A}} \bigoplus_{\substack{x \in F(X) \\ x \neq 0}} h_X \xrightarrow{x} F. \quad \square$$

Now suppose that \mathcal{A} has an additive symmetric monoidal structure \otimes , such as $\mathcal{A} = Cor_k$. (By this, we mean that \otimes commutes with direct sums; see 8A.3.) We may extend \otimes to a tensor product on \mathcal{A}^\oplus in the obvious way, and this extends to tensor product of projectives. We now extend \otimes to a tensor product on all of $R(\mathcal{A})$.

If F and G are in $R(\mathcal{A})$, we can form the presheaf tensor product $(F \otimes_R G)(X) = F(X) \otimes_R G(X)$. However, it does not belong to $R(\mathcal{A})$, since $F \otimes_R G$ is not additive. In order to get a tensor product on $R(\mathcal{A})$, we need a more complicated construction.

Our construction of \otimes is dictated by the requirement that if X and Y are in \mathcal{A} , then the tensor product $h_X \otimes h_Y$ of their representable presheaves should be represented by $X \otimes Y$. As a first step, note that we can extend \otimes to a tensor product $\otimes : \mathcal{A}^\oplus \times \mathcal{A}^\oplus \rightarrow \mathcal{A}^\oplus$ commuting with \bigoplus . Thus if L_1 and L_2 are in the category

$\mathbf{Ch}^-(\mathcal{A}^\oplus)$ of bounded above cochain complexes ($\cdots \rightarrow F^n \rightarrow 0 \rightarrow \cdots$), the chain complex $L_1 \otimes L_2$ is defined as the total complex of the double complex $L_1^* \otimes L_2^*$.

DEFINITION 8.2. If F and G are objects of $R(\mathcal{A})$, choose projective resolutions $P_* \rightarrow F$ and $Q_* \rightarrow G$ and define $F \otimes^{\mathbb{L}} G$ to be $P \otimes Q$, i.e., $\text{Tot}(P_* \otimes Q_*)$. We define the tensor product and Hom presheaves to be:

$$F \otimes G = H_0(F \otimes^{\mathbb{L}} G),$$

$$\underline{\text{Hom}}(F, G) : X \mapsto \text{Hom}_{R(\mathcal{A})}(F \otimes h_X, G).$$

Since any two projective resolutions of F are chain homotopy equivalent, the chain complex $F \otimes^{\mathbb{L}} G$ is well-defined up to chain homotopy equivalence, and similarly for $\underline{\text{Hom}}(F, G)$. In particular, since h_X and h_Y are projective, we have $h_X \otimes^{\mathbb{L}} h_Y = h_X \otimes h_Y = h_{X \otimes Y}$ for all X and Y in \mathcal{A}^\oplus .

The following result implies that $R(\mathcal{A})$ is an additive symmetric monoidal category (see 8A.3).

LEMMA 8.3. *The functor $\underline{\text{Hom}}(F, -)$ is right adjoint to $F \otimes -$. In particular, $\underline{\text{Hom}}(F, -)$ is left exact and $F \otimes -$ is right exact.*

PROOF. Because $R(\mathcal{A})$ has enough projectives, it suffices to observe that

$$\text{Hom}_{R(\mathcal{A})}(h_X, \underline{\text{Hom}}(h_Y, G)) = G(X \otimes Y) = \text{Hom}_{R(\mathcal{A})}(h_X \otimes h_Y, G). \quad \square$$

EXAMPLE 8.4. If \mathcal{A} is the category of free R -modules over a commutative ring R , $R(\mathcal{A})$ is equivalent to the category of all R -modules; the presheaf associated to M is $M \otimes_R$, and Hom and \otimes are the familiar Hom_R and \otimes_R .

EXERCISE 8.5. If F_i and G_i are in $R(\mathcal{A})$, show that there is a natural map

$$\underline{\text{Hom}}(F_1, G_1) \otimes \underline{\text{Hom}}(F_2, G_2) \rightarrow \underline{\text{Hom}}(F_1 \otimes F_2, G_1 \otimes G_2),$$

compatible with the monoidal pairing $\text{Hom}_{\mathcal{A}}(U \times A_1, X_1) \otimes \text{Hom}_{\mathcal{A}}(U \times A_2, X_2) \rightarrow \text{Hom}_{\mathcal{A}}(U \times U \times A_1 \times A_2, X_1 \times X_2) \rightarrow \text{Hom}_{\mathcal{A}}(U \times A_1 \times A_2, X_1 \times X_2)$.

REMARK 8.6. If the (projective) objects h_X are flat, i.e., $h_X \otimes -$ is an exact functor, then \otimes is called a balanced functor ([Wei94, 2.7.7]). In this case $F \otimes^{\mathbb{L}} G$ agrees (up to chain equivalence) with the usual left derived functor $\mathbb{L}(F \otimes -)G$. But we do not know when the h_X are flat. It is true in example 8.4, but probably not true in $\mathbf{PST} = \mathbb{Z}(\text{Cor}_k)$.

We can now extend $\otimes^{\mathbb{L}}$ to a total tensor product on the category $\mathbf{Ch}^-R(\mathcal{A})$ of bounded above cochain complexes ($\cdots \rightarrow F^n \rightarrow 0 \rightarrow \cdots$). This would be the usual derived functor if \otimes were balanced (see [Wei94, 10.6]), and our construction is parallel. If C is a complex in $\mathbf{Ch}^-R(\mathcal{A})$, there is a quasi-isomorphism $P \xrightarrow{\simeq} C$ with P a complex of projective objects. Any such complex P is called a projective resolution of C , and any other projective resolution of C is chain homotopic to P ; see [Wei94, 5.7]. If D is any other complex in $\mathbf{Ch}^-R(\mathcal{A})$, and $Q \xrightarrow{\simeq} D$ is a projective resolution, we define

$$C \otimes^{\mathbb{L}} D = P \otimes Q.$$

Because P and Q are bounded above, each $(P \otimes Q)^n = \bigoplus_{i+j=n} P^i \otimes Q^j$ is a finite sum, and $C \otimes^{\mathbb{L}} D$ is bounded above. Because P and Q are defined up to chain homotopy, the complex $C \otimes^{\mathbb{L}} D$ is independent (up to chain homotopy equivalence) of the choice of P and Q . There is a natural map $C \otimes^{\mathbb{L}} D \rightarrow C \otimes D$, which extends the map $F \otimes^{\mathbb{L}} G \rightarrow F \otimes G$ of definition 8.2.

LEMMA 8.7. *Let C, C' and D be bounded above complexes of presheaves.*

- (1) *If C and D are complexes over \mathcal{A}^\oplus , or complexes of projectives, then $C \otimes^{\mathbb{L}} D \xrightarrow{\simeq} C \otimes D$ is a chain homotopy equivalence.*
- (2) *If $f : C \xrightarrow{\simeq} C'$ is a quasi-isomorphism of complexes, then $C \otimes^{\mathbb{L}} D \rightarrow C' \otimes^{\mathbb{L}} D$ is a chain homotopy equivalence.*

PROOF. If C is a complex over \mathcal{A}^\oplus , it is a complex of projectives. We may take $P = C$ in the definition of $\otimes^{\mathbb{L}}$: $C \otimes^{\mathbb{L}} D = C \otimes Q$. If D is also a complex of projectives, we may take $Q = D$ as well. Part 1 is now immediate. In part 2, we may take P to be a projective resolution of both C and C' , so that $C \otimes^{\mathbb{L}} D = C' \otimes^{\mathbb{L}} D = P \otimes Q$. \square

PROPOSITION 8.8. *The derived category $\mathbf{D}^-R(\mathcal{A})$, equipped with $\otimes^{\mathbb{L}}$, is a tensor triangulated category.*

PROOF. The category \mathcal{P} of projective objects in $R(\mathcal{A})$ is additive symmetric monoidal, and $\mathbf{D}^-R(\mathcal{A})$ is equivalent to the chain homotopy category $\mathbf{K}^-(\mathcal{P})$ by [Wei94, 10.4.8]. By 8A.4, this is a tensor triangulated category under \otimes . The result now follows from the natural isomorphism $\otimes \cong \otimes^{\mathbb{L}}$ in \mathcal{P} of 8.7. \square

DEFINITION 8.9. If C and D are bounded above complexes of presheaves, there is a canonical map from the presheaf tensor product $C \otimes_R D$ to the tensor product $C \otimes D$ of 8.2. By right exactness of \otimes_R and \otimes (see 8.3), it suffices to construct a natural map of presheaves $\eta : h_X \otimes_R h_Y \rightarrow h_{X \otimes Y}$. For U in \mathcal{A} , η_U is just the monoidal product in \mathcal{A} , followed by the diagonal $\Delta : U \rightarrow U \otimes U$:

$$\begin{aligned} h_X(U) \otimes_R h_Y(U) &= \mathrm{Hom}_{\mathcal{A}}(U, X) \otimes_R \mathrm{Hom}_{\mathcal{A}}(U, Y) \\ &\xrightarrow{\otimes} \mathrm{Hom}_{\mathcal{A}}(U \otimes U, X \otimes Y) \xrightarrow{\Delta^*} \mathrm{Hom}_{\mathcal{A}}(U, X \otimes Y) = h_{X \otimes Y}(U). \end{aligned}$$

Having disposed with these generalities, we now specialize to the case where \mathcal{A} is Cor_k and \otimes is the tensor product $X \otimes Y = X \times Y$ of 1.9. We have the Yoneda embedding

$$\mathrm{Cor}_k \subset \mathrm{Cor}_k^\oplus \subset \mathbf{PST}(k).$$

We will write \otimes^{tr} for the tensor product on $\mathbf{PST} = \mathbb{Z}(\mathrm{Cor}_k)$, or on $\mathbf{PST}(k, R) = R(\mathrm{Cor}_k)$, and \otimes_L^{tr} for $\otimes^{\mathbb{L}}$. Thus there are natural maps $C \otimes_L^{tr} D \rightarrow C \otimes^{tr} D$.

EXAMPLE 8.10. By lemma 8.1, $h_X = R_{tr}(X)$ is projective and

$$R_{tr}(X) \otimes^{tr} R_{tr}(Y) = R_{tr}(X \times Y).$$

Similarly if (X_i, x_i) are pointed schemes then the $R_{tr}(X_i, x_i)$ are projective and from 2.13 we see that

$$R_{tr}(X_1, x_1) \otimes^{tr} \cdots \otimes^{tr} R_{tr}(X_n, x_n) = R_{tr}((X_1, x_1) \wedge \cdots \wedge (X_n, x_n)).$$

In particular, $R_{tr}(\mathbb{G}_m)^{\otimes^{tr} n} = R_{tr}(\mathbb{G}_m^{\wedge n})$.

The next example, in which $R = \mathbb{Z}$, shows that \otimes^{tr} does not behave well on locally constant sheaves.

EXAMPLE 8.11. The complex $\mathbb{Z} \xrightarrow{n} \mathbb{Z}$ is a projective resolution of \mathbb{Z}/n , so we have $\mathbb{Z}/n \otimes^{tr} \mathbb{Z}_{tr}(X) = \mathbb{Z}/n \otimes_{\mathbb{Z}} \mathbb{Z}_{tr}(X) = (\mathbb{Z}/n)_{tr}(X)$ by 8.7.

If $\sqrt{-1} \notin k$ and $l = k(\sqrt{-1})$, let $\mathbb{Z}_{\varepsilon} = \mathbb{Z}_{tr}(l)/\mathbb{Z}$ denote the locally constant sheaf corresponding to the sign representation of $G = \text{Gal}(l/k)$. We see from 8.7 that $\mathbb{Z}/n \otimes_L^{tr} \mathbb{Z}_{\varepsilon}$ is quasi-isomorphic to the complex $(\mathbb{Z}/n) \otimes^{\mathbb{L}} (\mathbb{Z} \rightarrow \mathbb{Z}_{tr}(l))$, i.e.,

$$0 \rightarrow \mathbb{Z}/n \rightarrow (\mathbb{Z}/n)_{tr}(l) \rightarrow 0.$$

Hence the presheaf $(\mathbb{Z}/n) \otimes^{tr} \mathbb{Z}_{\varepsilon}$ sends $\text{Spec } k$ to 0 and $\text{Spec } l$ to \mathbb{Z}/n . If $n = 4$, this is not an étale sheaf because $(\mathbb{Z}_{\varepsilon}/4\mathbb{Z}_{\varepsilon})^G \neq 0$. It is easy to see, however, that its sheafification is the locally constant étale sheaf:

$$((\mathbb{Z}/4) \otimes^{tr} \mathbb{Z}_{\varepsilon})_{\acute{e}t} \cong \mu_4.$$

The étale sheaf μ_4 is the tensor product $(\mathbb{Z}/4) \otimes_{\acute{e}t} \mathbb{Z}_{\varepsilon}$ of the two underlying étale sheaves.

DEFINITION 8.12. If F and G are presheaves of R -modules with transfers, we write $F \otimes_{\acute{e}t}^{tr} G$ for $(F \otimes^{tr} G)_{\acute{e}t}$, the étale sheaf associated to $F \otimes^{tr} G$. If C and D are bounded above complexes of presheaves with transfers, we shall write $C \otimes_{\acute{e}t}^{tr} D$ for $(C \otimes^{tr} D)_{\acute{e}t}$, and $C \otimes_{L, \acute{e}t}^{tr} D$ for $(C \otimes_L^{tr} D)_{\acute{e}t} \simeq P \otimes_{\acute{e}t}^{tr} Q$, where P and Q are complexes of representable sheaves with transfers, and $P \simeq C$ and $Q \simeq D$. There is a natural map $C \otimes_{L, \acute{e}t}^{tr} D \rightarrow C \otimes_{\acute{e}t}^{tr} D$, induced by $C \otimes_L^{tr} D \rightarrow C \otimes^{tr} D$.

LEMMA 8.13. *If F, F' are étale sheaves of R -modules with transfers, and F is locally constant, then the map of 8.9 induces an isomorphism*

$$F \otimes_{\acute{e}t} F' \xrightarrow{\cong} F \otimes_{\acute{e}t}^{tr} F'.$$

PROOF. Let F correspond to the discrete Galois module M . As $M = \cup M^H$ and \otimes^{tr} commutes with colimits, we may assume that $M = M^H$ for some open normal H of $\text{Gal}(k_{sep}/k)$. Thus M is a G -module, where $G = \text{Gal}(k_{sep}/k)/H$. Choose a presentation over $R[G]$:

$$\oplus R[G]^{\alpha} \rightarrow \oplus R[G]^{\beta} \rightarrow M \rightarrow 0.$$

As $\otimes_{\acute{e}t}$ and $\otimes_{\acute{e}t}^{tr}$ are both right exact, we may assume $M = R[G]$ and $F' = R_{tr}(X)$. If $L = (k_{sep})^H$ and $T = \text{Spec}(L)$ then $F = R_{tr}(T)$ by exercise 6.10. But then $F \otimes^{tr} F' = R_{tr}(T \times X)$, so it suffices to observe that $R_{tr}(T) \otimes_{\acute{e}t} R_{tr}(X) \rightarrow R_{tr}(T \times X)$ is an

isomorphism. Since $T \times Y \rightarrow Y$ is an étale cover, it suffices to observe that for Y over T

$$\begin{aligned} R_{tr}(T) \otimes_{\acute{e}t} R_{tr}(X)(Y) &\cong R[G] \otimes \text{Cor}(Y, X) \\ &\cong R \otimes_{\mathbb{Z}} \text{Cor}(Y, T \times X) = R_{tr}(T \times X)(Y). \quad \square \end{aligned}$$

We are now going to show (in 8.16) that the tensor product $\otimes_{L, \acute{e}t}^{tr}$ induces a tensor triangulated structure on the derived category of étale sheaves of R -modules with transfers. Using proposition 8.8, we have $C \otimes_{L, \acute{e}t}^{tr} D \cong D \otimes_{L, \acute{e}t}^{tr} C$, and it suffices to show that $\otimes_{L, \acute{e}t}^{tr}$ preserves quasi-isomorphisms.

As a first step, fix Y and consider the right exact functor $\Phi(F) = R_{tr}(Y) \otimes_{\acute{e}t}^{tr} F$, from the category $\mathbf{PST}(k, R)$ of presheaves of R -modules with transfers to the category of étale sheaves of R -modules with transfers. Its left derived functors $L_p \Phi(F)$ are the homology sheaves of the total left derived functor $R_{tr}(Y) \otimes_{L, \acute{e}t}^{tr} F$. If C is a chain complex (bounded below in homological notation), the hyperhomology spectral sequence (see [Wei94, 5.7.6]) is

$$E_{p,q}^2 = L_p \Phi(H_q C) \Rightarrow \mathbb{L}_{p+q} \Phi(C).$$

EXAMPLE 8.14. If $U \rightarrow X$ is an étale cover, consider the augmented Čech complex

$$\check{C}: \quad \cdots \rightarrow R_{tr}(U \times_X U) \rightarrow R_{tr}(U) \rightarrow R_{tr}(X) \rightarrow 0.$$

Since $\check{C}_{\acute{e}t}$ is exact by 6.12, each homology presheaf $H_q(U/X) = H_q(\check{C})$ satisfies $H_q(U/X)_{\acute{e}t} = 0$. By definition, $R_{tr}(Y) \otimes_{\acute{e}t}^{tr} \check{C}$ is the augmented Čech complex

$$\cdots \rightarrow R_{tr}(U \times_X U \times Y) \rightarrow R_{tr}(U \times Y) \rightarrow R_{tr}(X \times Y) \rightarrow 0$$

for the étale cover $U \times Y \rightarrow X \times Y$, so $R_{tr}(Y) \otimes_{\acute{e}t}^{tr} \check{C}$ is again exact by 6.12. Thus $\mathbb{L}_n \Phi(\check{C}) = 0$ for all n . In particular, the 0th homology presheaf $H_0(U/X)$ satisfies

$$\Phi H_0(U/X) = R_{tr}(Y) \otimes_{\acute{e}t}^{tr} H_0(U/X) = H_0(R_{tr}(Y) \otimes_{\acute{e}t}^{tr} \check{C}) = 0.$$

The following lemma shows that in fact every derived functor $L_n \Phi$ vanishes on $H_0(U/X)$.

LEMMA 8.15. *Fix Y and set $\Phi = R_{tr}(Y) \otimes_{\acute{e}t}^{tr}$. If F is a presheaf of R -modules with transfers such that $F_{\acute{e}t} = 0$, then $L_n \Phi(F) = 0$ for all n .*

PROOF. Suppose that $F_{\acute{e}t} = 0$. Each map $R_{tr}(X) \rightarrow F$ is defined by an $x \in F(X)$, and there is an étale cover $U_x \rightarrow X$ such that x vanishes in $F(U_x)$. Thus the composition $R_{tr}(U_x) \rightarrow R_{tr}(X) \rightarrow F$ is zero, i.e., the given map factors through the cokernel $H_0(U_x/X)$ of $R_{tr}(U_x) \rightarrow R_{tr}(X)$. It follows that the canonical surjection $\oplus_{X,x} R_{tr}(X) \rightarrow F$ factors through a surjection $\oplus_{X,x} H_0(U_x/X) \rightarrow F$. If K denotes the kernel of this surjection then $K_{\acute{e}t} = 0$.

We now proceed by induction on n , noting that $L_n \Phi = 0$ for $n < 0$. For $n = 0$, we know that $\Phi H_0(U_x/X) = 0$ by example 8.14. Since Φ is right exact, this yields

$\Phi(F) = 0$. For $n > 0$, we may assume that the lemma holds for $L_p \Phi$ when $p < n$. From the exact sequence

$$\oplus_{X,x} (L_n \Phi) H_0(U_x/X) \rightarrow L_n \Phi(F) \rightarrow L_{n-1} \Phi(K)$$

we see that it suffices to prove that $(L_n \Phi) H_0(U/X) = 0$. We saw in 8.14 that $H_q(U/X)_{\acute{e}t} = 0$, so $L_p \Phi H_q(U/X) = 0$ by the inductive assumption. Hence the hypercohomology sequence for the complex \check{C} collapses to yield

$$\mathbb{L}_n \Phi(\check{C}) \cong (L_n \Phi) H_0(\check{C}) = (L_n \Phi) H_0(U/X).$$

But we saw in example 8.14 that $L_n \Phi(\check{C}) = 0$, whence the result. \square

Now we prove that $\otimes_{L,\acute{e}t}^{tr}$ preserves quasi-isomorphisms.

PROPOSITION 8.16. *Let $f : C \rightarrow C'$ be a morphism of bounded above complexes of presheaves of R -modules with transfers. If f induces a quasi-isomorphism $C_{\acute{e}t} \rightarrow C'_{\acute{e}t}$ between the associated complexes of étale sheaves, then $C \otimes_{L,\acute{e}t}^{tr} D \rightarrow C' \otimes_{L,\acute{e}t}^{tr} D$ is a quasi-isomorphism for every D .*

PROOF. If $P \xrightarrow{\simeq} C$ is a projective resolution of presheaves, then $P_{\acute{e}t} \rightarrow C_{\acute{e}t}$ is a quasi-isomorphism of complexes of étale sheaves. Thus we may assume that C, C' and D are complexes of representable presheaves. If A denotes the mapping cone of $C \rightarrow C'$, it suffices to show that $A \otimes_{L,\acute{e}t}^{tr} D = A \otimes_{\acute{e}t}^{tr} D$ is acyclic. As each row of the double complex underlying $A \otimes_{\acute{e}t}^{tr} D$ is a sum of terms $A \otimes_{\acute{e}t}^{tr} R_{tr}(Y)$, it suffices to show that $A \otimes_{\acute{e}t}^{tr} R_{tr}(Y)$ is acyclic. As in the proof of 8.15, its homology sheaves are the hyper-derived functors $\mathbb{L}_n \Phi(A)$, $\Phi = \otimes_{\acute{e}t}^{tr} R_{tr}(Y)$. In the hypercohomology spectral sequence

$$E_{p,q}^2 = L_p \Phi(H_q A) \Rightarrow \mathbb{L}_{p+q} \Phi(A)$$

the presheaves $H_q A$ have $(H_q A)_{\acute{e}t} = 0$ because $A_{\acute{e}t}$ is acyclic. By lemma 8.15 we have $L_q \Phi(H_q A) = 0$ for all p and q . Hence the spectral sequence collapses to yield $\mathbb{L}_n \Phi(A) = 0$ for all n , i.e., $\mathbb{L} \Phi(A) \simeq R_{tr}(Y) \otimes_{\acute{e}t}^{tr} A$ is acyclic. \square

COROLLARY 8.17. *The derived category of bounded above complexes of étale sheaves of R -modules with transfers is a tensor triangulated category.*

PROOF. By 8.8, $\mathbf{D}^- \mathbf{PST}(k, R)$ is tensor triangulated. Now combine 8.16 and 8A.7, letting W be the system of morphisms inducing quasi-isomorphisms on the associated complexes of étale sheaves. \square

LEMMA 8.18. *Let F be a locally constant étale sheaf of flat R -modules. Then the map $E \otimes_{L,\acute{e}t}^{tr} F \rightarrow E \otimes_{\acute{e}t}^{tr} F$ is a quasi-isomorphism for every étale sheaf with transfers E .*

PROOF. Suppose first that $E = R_{tr}(Y)$. Choose a resolution $C \rightarrow F$ in the category of locally constant sheaves in which each C_n is a sum of representables $R_{tr}(L_{n,\alpha})$ for finite Galois field extensions $L_{n,\alpha}$ of k . (This is equivalent to resolving the Galois module M corresponding to F by Galois modules $R[G_{n,\alpha}]$,

and the existence of such a resolution of M is well known.) By proposition 8.16, $E \otimes_{\acute{e}t}^{tr} C = E \otimes_{L, \acute{e}t}^{tr} C$ is quasi-isomorphic to $E \otimes_{L, \acute{e}t}^{tr} F$. By lemma 8.13,

$$E \otimes_{\acute{e}t}^{tr} C = E \otimes_{\acute{e}t} C \xrightarrow{\simeq} E \otimes_{\acute{e}t} F \xleftarrow{\simeq} E \otimes_{\acute{e}t}^{tr} F.$$

Hence the result is true for $E = R_{tr}(Y)$.

In the general case, choose a projective resolution $P \rightarrow E$ in the category of presheaves of R -modules with transfers. Then we have quasi-isomorphisms

$$E \otimes_{L, \acute{e}t}^{tr} F = P \otimes_{L, \acute{e}t}^{tr} F \xrightarrow{\simeq} P \otimes_{\acute{e}t}^{tr} F \xrightarrow{\simeq} P \otimes_{\acute{e}t} F.$$

Because sheafification is exact, $P \rightarrow E$ is also a resolution in the category of étale sheaves of R -modules. Since F is flat in this category, we have the final quasi-isomorphism:

$$P \otimes_{\acute{e}t} F \xrightarrow{\simeq} E \otimes_{\acute{e}t} F \xleftarrow{\simeq} E \otimes_{\acute{e}t}^{tr} F. \quad \square$$

It is clear that 8.18 also holds if E is a bounded above complex of étale sheaves with transfers.

COROLLARY 8.19. *In the derived category of étale sheaves of \mathbb{Z}/m -modules with transfers, the operation $M \mapsto M(1) = M \otimes_{L, \acute{e}t}^{tr} \mathbb{Z}/m(1)$ is invertible.*

PROOF. Indeed, if μ_m^* is the Pontrjagin dual of μ_m , then combining 8.18, 8.13, and 4.8 yields:

$$\mu_m^* \otimes_{L, \acute{e}t}^{tr} \mathbb{Z}/m(1) \stackrel{8.18}{\simeq} \mu_m^* \otimes_{\acute{e}t}^{tr} \mathbb{Z}/m(1) \stackrel{8.13}{\simeq} \mu_m^* \otimes_{\acute{e}t} \mathbb{Z}/m(1) \stackrel{4.8}{\simeq} \mu_m^* \otimes_{\acute{e}t} \mu_m \cong \mathbb{Z}/m. \quad \square$$

EXERCISE 8.20. If E and F are bounded above complexes of locally constant étale sheaves of R -modules, show that $E \otimes_{L, \acute{e}t}^{tr} F$ is quasi-isomorphic to $E \otimes_R^{\mathbb{L}} F$, their total tensor product as complexes of étale sheaves of R -modules. (Hint: Use 8.13, 8.16, and 8.18.)

REMARK 8.21. If $B \rightarrow I$ is a flasque resolution of B as a sheaf with transfers, we define $\underline{RHom}(R_{tr}X, B)$ to be $\underline{Hom}(R_{tr}X, I)$, so that $\underline{RHom}(R_{tr}X, B)(U) = \underline{RHom}(U \times X, B)$ for all U . If $cd(k) < \infty$ and X is proper then $\underline{RHom}(R_{tr}X, B)$ is bounded above by proper base change (citing 9.26); this construction extends to bounded above complexes B in the usual way. If A and B are both bounded above, a short calculation shows that in the derived category $\mathbf{D}^-(Sh(Cor_k, R))$ of sheaves with transfers we have the adjunction:

$$\mathrm{Hom}_{\mathbf{D}^-}(A \otimes_{L, \acute{e}t}^{tr} R_{tr}(X), B) \cong \mathrm{Hom}_{\mathbf{D}^-}(A, \underline{RHom}(R_{tr}X, B)).$$

Tensor triangulated categories

The notion of a tensor triangulated category is a generalization of the tensor product structure on the derived category of modules over a scheme, which played a central role in the development of the subject.

DEFINITION 8A.1. A **tensor triangulated category** is an additive category with two structures: that of a triangulated category and that of a symmetric monoidal category. In addition, we are given natural isomorphisms r and l of the form

$$C[1] \otimes D \xrightarrow[l_{C,D}]{\cong} (C \otimes D)[1] \xleftarrow[r_{C,D}]{\cong} C \otimes D[1],$$

which commute in the obvious sense with the associativity, commutativity and unity isomorphisms. There are two additional axioms:

(TTC1) For any distinguished triangle $C_0 \longrightarrow C_1 \longrightarrow C_2 \xrightarrow{\partial} C_0[1]$ and any D , the following triangles are distinguished:

$$C_0 \otimes D \longrightarrow C_1 \otimes D \longrightarrow C_2 \otimes D \xrightarrow{l(\partial \otimes D)} (C_0 \otimes D)[1],$$

$$D \otimes C_0 \longrightarrow D \otimes C_1 \longrightarrow D \otimes C_2 \xrightarrow{r(D \otimes \partial)} (D \otimes C_0)[1].$$

(TTC2) For any C and D , the following diagram commutes up to multiplication by -1 , i.e., $rl = -lr$:

$$\begin{array}{ccc} C[1] \otimes D[1] & \xrightarrow{r} & (C[1] \otimes D)[1] \\ \downarrow l & & \downarrow l \\ & -1 & \\ (C \otimes D[1])[1] & \xrightarrow{r} & (C \otimes D)[2]. \end{array}$$

This description is not minimal. For example the commutativity isomorphism $\tau : C \otimes D \cong D \otimes C$ allows us to recover r from l and vice versa using the formula $\tau l \tau = r$. In addition, $l_{C,D}$ can be recovered from $l_{1,D} : \mathbf{1}[1] \otimes D \cong D[1]$, where $\mathbf{1}$ is the identity object for \otimes . Moreover, if either of the two triangles in (TTC1) is distinguished, then both are distinguished.

The definition of tensor triangulated category that we have given is sufficient for our purposes. However, it is possible to add extra axioms in order to work

with a richer structure. For example, many more axioms are postulated by May in [May01].

EXERCISE 8A.2. Show that the canonical isomorphisms $l^i r^j, r^j l^i : C[i] \otimes D[j] \cong (C \otimes D)[i+j]$ differ by $(-1)^{ij}$, and are interchanged by the twist isomorphism τ on $C \otimes D$ and $C[i] \otimes D[j]$.

DEFINITION 8A.3. Let \mathcal{A} be an additive category with a symmetric monoidal structure \otimes . We say that \mathcal{A} is an **additive symmetric monoidal category** if $(\coprod A_i) \otimes B \cong \coprod (A_i \otimes B)$ for every direct sum $\coprod A_i$ in \mathcal{A} .

If C and D are bounded above complexes in \mathcal{A} , the tensor product $C \otimes D$ has $(C \otimes D)^n = \bigoplus_{p+q=n} C^p \otimes D^q$ and differential $d \otimes 1 + (-1)^p \otimes d$ on $C^p \otimes D^q$. It is associative.

We define the twist isomorphism $\tau : C \otimes D \rightarrow D \otimes C$ componentwise, as $(-1)^{pq}$ times the natural isomorphism $C^p \otimes D^q \rightarrow D^q \otimes C^p$ in \mathcal{A} . It is a straightforward exercise to verify that the category $\mathbf{Ch}^-(\mathcal{A})$ is an additive symmetric monoidal category.

The degree n part of each of $C \otimes D[1]$, $(C \otimes D)[1]$, and $C[1] \otimes D$ are the same, and we define $l_{C,D}$ to be the canonical isomorphism. The map $r_{C,D}$ is multiplication by $(-1)^p$ on the summand $C^p \otimes D^q$. A routine calculation verifies the following.

PROPOSITION 8A.4. *Let \mathcal{A} be an additive symmetric monoidal category. Then the chain homotopy category $\mathbf{K}^-(\mathcal{A})$ of bounded above cochain complexes is a tensor triangulated category.*

EXAMPLE 8A.5. (See [Ver96].) Let \mathcal{A} be the category of modules over a commutative ring, or more generally over a scheme. Then not only is $\mathbf{K}^-(\mathcal{A})$ a tensor triangulated category, but the total tensor product $\otimes^{\mathbb{L}}$ makes the derived category $\mathbf{D}^-(\mathcal{A})$ into a tensor triangulated category. In effect, $\mathbf{D}^-(\mathcal{A})$ is equivalent to the tensor triangulated subcategory of flat complexes in $\mathbf{K}^-(\mathcal{A})$.

EXAMPLE 8A.6. The smash product of based topological spaces leads to another example. If $A \rightarrow X \rightarrow X/A \rightarrow SA$ is a cofibration sequence, there is a natural homeomorphism $(X/A) \wedge Y \cong (X \wedge Y)/(A \wedge Y)$; see [Whi78, III.2.3]. The suspension $SX = S^1 \wedge X$ has homeomorphisms

$$X \wedge (SY) \xrightarrow[r]{\cong} S(X \wedge Y) \xleftarrow[l]{\cong} (SX) \wedge Y$$

satisfying (TTC1) and (TTC2) up to homotopy. It follows easily that the stable homotopy category, which is triangulated by [Wei94, 10.9.18] and a symmetric monoidal category by [Ada74, III.4], is a tensor triangulated category.

If W is a saturated multiplicative system of morphisms in a triangulated category \mathbf{D} , closed under \oplus , translations, and cones, Verdier proved in [Ver96] that the localization $\mathbf{D}[W^{-1}]$ is also a triangulated category.

PROPOSITION 8A.7. *Let \mathbf{D} be a tensor triangulated category. Suppose that if $C \rightarrow C'$ is in W then $C \otimes D \rightarrow C' \otimes D$ is in W for every D in \mathbf{D} . Then the localization $\mathbf{D}[W^{-1}]$ is also a tensor triangulated category.*

PROOF. Because each $\otimes D : \mathbf{D} \rightarrow \mathbf{D}$ preserves W , \otimes induces a symmetric monoidal pairing $\mathbf{D}[W^{-1}] \times \mathbf{D}[W^{-1}] \rightarrow \mathbf{D}[W^{-1}]$ by the universal property of localization (applied to $W \times W$). Similarly, the natural isomorphisms r and l descend to $\mathbf{D}[W^{-1}]$. Axiom (TTC2) is automatic, and axiom (TTC1) may be routinely verified for Verdier's description of distinguished triangles in $\mathbf{D}[W^{-1}]$. \square

Inverting twists $X \mapsto X \otimes T$ is another construction which often preserves the tensor triangulated structure. For example, it is used to construct the tensor triangulated category $\mathbf{DM}_{\text{ét}}^-(k, \mathbb{Z}/m)$ from $\mathbf{DM}_{\text{ét}}^{\text{eff}, -}(k, \mathbb{Z}/m)$; see 9.7.

Let T be an object in a symmetric monoidal category $(\mathcal{C}, \otimes, \mathbf{1})$. Let $\mathcal{C}[T^{-1}]$ denote the category whose objects are pairs (X, m) with X in \mathcal{C} and $m \in \mathbb{Z}$; morphisms $(X, m) \rightarrow (Y, n)$ in $\mathcal{C}[T^{-1}]$ are just elements of the direct limit $\lim_{i \rightarrow \infty} \text{Hom}(X \otimes T^{\otimes m+i}, Y \otimes T^{\otimes n+i})$, where the bonding maps are given by the functor $\otimes T : \mathcal{C} \rightarrow \mathcal{C}$. Composition is defined in the obvious way, and it's easy to check that $\mathcal{C}[T^{-1}]$ is a category. There is a universal functor $\mathcal{C} \rightarrow \mathcal{C}[T^{-1}]$ sending X to $(X, 0)$. Note that $(X, m) \cong X \otimes T^{\otimes m}$ in $\mathcal{C}[T^{-1}]$ for $m \geq 0$.

EXERCISE 8A.8. Let T be an object in a tensor triangulated category \mathcal{C} . Show that $\mathcal{C}[T^{-1}]$ is a triangulated category, and that $\mathcal{C} \rightarrow \mathcal{C}[T^{-1}]$ is triangulated.

In order for the formula $(X, m) \otimes (Y, n) = (X \otimes Y, m+n)$ to extend to a bifunctor on $\mathcal{C}[T^{-1}]$, we need to define the tensor $f \otimes g$ of two $\mathcal{C}[T^{-1}]$ -morphisms in a natural way. In general, $\mathcal{C}[T^{-1}]$ need not be symmetric monoidal, as exercise 8A.9 shows.

EXERCISE 8A.9. Let T be an invertible object in a symmetric monoidal category \mathcal{C} , i.e., an object such that $T \otimes U \cong \mathbf{1}$ for some U . It is well known that endomorphisms of $\mathbf{1}$ commute; show that the same must be true for endomorphisms of T . Then show that the cyclic permutation of $T \otimes (T \otimes T)$ must equal the identity morphism.

PROPOSITION 8A.10. *Let T be an object in a symmetric monoidal category $(\mathcal{C}, \otimes, \mathbf{1})$ such that the cyclic permutation on $T^{\otimes 3}$ is the identity in $\mathcal{C}[T^{-1}]$. Then $(\mathcal{C}[T^{-1}], \otimes, \mathbf{1})$ is also a symmetric monoidal category.*

PROOF. The hypothesis implies that permutations on $T^{\otimes n}$ commute with each other for $n \geq 3$. The many ways to define $f \otimes g$ on $X \otimes T^{m+i} \otimes Y \otimes T^{n+j}$ are indexed by the (i, j) -shuffles, and differ only by a permutation, so $f \otimes g$ is independent of this choice. Therefore the tensor product is a bifunctor on $\mathcal{C}[T^{-1}]$. The symmetric monoidal axioms may now be routinely verified as in [Ada74, III.4]. The hexagonal axiom, that the two isomorphisms from $X \otimes (Y \otimes Z)$ to $(Z \otimes X) \otimes Y$ agree, follows because the cyclic permutation on $T^{\otimes 3}$ is the identity. \square

COROLLARY 8A.11. *Let T be an object in a tensor triangulated category \mathcal{C} such that the cyclic permutation on $T^{\otimes 3}$ is the identity in $\mathcal{C}[T^{-1}]$. Then $\mathcal{C}[T^{-1}]$ is a tensor triangulated category.*

PROOF. By 8A.8 and 8A.10, $\mathcal{C}[T^{-1}]$ is both triangulated and symmetric monoidal. The verification of the remaining axioms is straightforward. \square

EXERCISE 8A.12. Let T be an object in a tensor triangulated category \mathbf{D} such that $\mathrm{Hom}(X, Y) \rightarrow \mathrm{Hom}(X \otimes T, Y \otimes T)$ is an isomorphism for every X and Y in \mathbf{D} . Show that $\mathbf{D}[T^{-1}]$ is a tensor triangulated category.

LECTURE 9

\mathbb{A}^1 -weak equivalence

In this section we define the notion of \mathbb{A}^1 -weak equivalence between bounded above cochain complexes of étale sheaves with transfers, and \mathbb{A}^1 -local complexes. The category $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}$ is obtained by inverting \mathbb{A}^1 -weak equivalences. The main result in this lecture (9.35) is that when we restrict to sheaves of \mathbb{Z}/m -modules the category $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}$ is equivalent to the derived category of discrete Galois modules for the group $\text{Gal}(k_{\text{sep}}/k)$. We will use these ideas in the next lecture to identify étale motivic cohomology with ordinary étale cohomology.

Since quasi-isomorphic complexes will be \mathbb{A}^1 -weak equivalent, it is appropriate to define the notion in the derived category $\mathbf{D}^- = \mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$ of étale sheaves of R -modules with transfers. In \mathbf{D}^- , we have the usual shift, and

$$A \xrightarrow{f} B \longrightarrow \text{cone}(f) \longrightarrow A[1]$$

is a distinguished triangle for each map f . We refer the reader to [GM03] or [Wei94] for basic facts about derived categories. We will also need the notion of a thick subcategory, which was introduced by Verdier in [Ver96]. We will use Rickard's definition (see [Ric89]); this is slightly different from, but equivalent to, Verdier's definition.

DEFINITION 9.1. A full additive subcategory \mathcal{E} of \mathbf{D}^- is **thick** if:

- (1) Let $A \rightarrow B \rightarrow C \rightarrow A[1]$ be a distinguished triangle. Then if two out of A, B, C are in \mathcal{E} then so is the third.
- (2) if $A \oplus B$ is in \mathcal{E} then both A and B are in \mathcal{E} .

If \mathcal{E} is a thick subcategory of \mathbf{D}^- , we can form a quotient triangulated category \mathbf{D}^-/\mathcal{E} as follows (see [Ver96]). Let $W_{\mathcal{E}}$ be the set of maps whose cone is in \mathcal{E} ; $W_{\mathcal{E}}$ is a saturated multiplicative system of morphisms. Then \mathbf{D}^-/\mathcal{E} is the localization $\mathbf{D}^-[W_{\mathcal{E}}^{-1}]$, which may be constructed using calculus of fractions; see [Wei94, 10.3.7]. In particular, a morphism $f : C \rightarrow C'$ becomes an isomorphism in $\mathbf{D}^-[W_{\mathcal{E}}^{-1}]$ if and only if f is in $W_{\mathcal{E}}$.

DEFINITION 9.2. A morphism f in \mathbf{D}^- is called an **\mathbb{A}^1 -weak equivalence** if f is in $W_{\mathcal{E}_{\mathbb{A}}}$, where $\mathcal{E}_{\mathbb{A}}$ is the smallest thick subcategory so that:

- (1) the cone of $R_{tr}(X \times \mathbb{A}^1) \rightarrow R_{tr}(X)$ is in $\mathcal{E}_{\mathbb{A}}$ for every smooth scheme X ;
- (2) $\mathcal{E}_{\mathbb{A}}$ is closed under any direct sum that exists in \mathbf{D}^- .

We set $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R) = \mathbf{D}^-[W_{\mathcal{E}_{\mathbb{A}}}^{-1}]$.

REMARK 9.3. Alternatively, we can describe $\mathcal{E}_{\mathbb{A}}$ as the thick subcategory of all complexes E such that $C_*(E)$ is acyclic (i.e., quasi-isomorphic to zero). Indeed, it follows from 2.24 that $C_*(E)$ is acyclic for every E in $\mathcal{E}_{\mathbb{A}}$. Conversely, if $C_*(E)$ is acyclic then $E \rightarrow 0$ is in $W_{\mathbb{A}}$ by 9.15 below, and hence E is in $\mathcal{E}_{\mathbb{A}}$.

It is clear that the notion of \mathbb{A}^1 -weak equivalence in $\mathbf{D}^- = \mathbf{D}^-(Sh(Cor_k, R))$ makes sense for other topologies. This includes the alternative description in 9.3. For the Nisnevich topology, we will see in 14.11 that the localization $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ of \mathbf{D}^- is the triangulated category of motivic complexes introduced and studied in [TriCa].

LEMMA 9.4. *The smallest class in \mathbf{D}^- which contains all the $R_{tr}(X)$ and is closed under quasi-isomorphisms, direct sums, shifts, and cones is all of \mathbf{D}^- .*

PROOF. First we show that for any complex D_* , if all D_n are in the class, then so is D_* . If $\beta_n D$ is the brutal truncation $0 \rightarrow D_n \rightarrow D_{n-1} \rightarrow \cdots$ of D_* , then D_* is the union of the $\beta_n D$. Each $\beta_n D$ is a finite complex, belonging to the class, as an inductive argument shows. Since there is an exact sequence

$$0 \longrightarrow \bigoplus \beta_n D \longrightarrow \bigoplus \beta_n D \longrightarrow D_* \longrightarrow 0,$$

it follows that D_* is in the class.

Thus it suffices to show that each sheaf F is in the class. Now there is a resolution $L_* \rightarrow F$ by sums of the representable sheaves $R_{tr}(X)$, given by lemma 8.1. Since each L_n is in this class, so is L_* and hence F . \square

LEMMA 9.5. *If $f : C \rightarrow C'$ is an \mathbb{A}^1 -weak equivalence, then for every D the map $f \otimes Id : C \otimes_{L, \acute{e}t}^{tr} D \rightarrow C' \otimes_{L, \acute{e}t}^{tr} D$ is an \mathbb{A}^1 -weak equivalence.*

PROOF. Since $\otimes_{L, \acute{e}t}^{tr}$ commutes with cones and f is an \mathbb{A}^1 -weak equivalence if and only if its cone is in $\mathcal{E}_{\mathbb{A}}$, it suffices to show that if C is in $\mathcal{E}_{\mathbb{A}}$, then $C \otimes_{L, \acute{e}t}^{tr} D$ is in $\mathcal{E}_{\mathbb{A}}$ for any D .

If $D = R_{tr}(X)$, consider the subcategory \mathcal{E} of all C in \mathbf{D}^- such that $C \otimes_{L, \acute{e}t}^{tr} D$ is in $\mathcal{E}_{\mathbb{A}}$. \mathcal{E} is closed under direct sums and it is thick. Moreover, if Y is a smooth scheme, then \mathcal{E} contains the cone of $R_{tr}(Y \times \mathbb{A}^1) \rightarrow R_{tr}(Y)$. Therefore $\mathcal{E}_{\mathbb{A}} \subseteq \mathcal{E}$.

Now fix C in $\mathcal{E}_{\mathbb{A}}$ and consider the full subcategory \mathcal{D} of all D in \mathbf{D}^- such that $C \otimes_{L, \acute{e}t}^{tr} D$ is in $\mathcal{E}_{\mathbb{A}}$. \mathcal{D} is closed under direct sums, it is thick and we have seen that it contains $R_{tr}(X)$ for all X . By 9.4, we conclude that $\mathcal{D} = \mathbf{D}^-$. \square

COROLLARY 9.6. *The product $\otimes_{L, \acute{e}t}^{tr}$ endows $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$ with the structure of a tensor triangulated category.*

PROOF. Given 8.17, this follows from 9.5 and proposition 8A.7. \square

REMARK 9.7. The category $\mathbf{DM}_{\acute{e}t}^-(k, R)$ is obtained from $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$ by inverting the Tate twist operation $M \mapsto M(1) = M \otimes_{L, \acute{e}t}^{tr} R(1)$. If $R = \mathbb{Z}/m$, then the Tate twist is already invertible by 8.19, so we have

$$\mathbf{DM}_{\acute{e}t}^-(k, \mathbb{Z}/m) = \mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, \mathbb{Z}/m).$$

For any coefficients R , it will follow from 8A.11 and 15.8 below that $\mathbf{DM}_{\acute{e}t}^-(k, R)$ is always a tensor triangulated category.

DEFINITION 9.8. Two morphisms $F \xrightarrow[f]{g} G$ of sheaves of R -modules with transfers are called \mathbb{A}^1 -**homotopic** if there is a map $h : F \otimes^{tr} R_{tr}(\mathbb{A}^1) \rightarrow G$ so that the restrictions of h along $R \xrightarrow[0]{1} R_{tr}(\mathbb{A}^1)$ coincide with f and g .

If G is an étale sheaf, h factors through (and is determined by) a map $F_{\acute{e}t} \otimes_{L, \acute{e}t}^{tr} R_{tr}(\mathbb{A}^1) \rightarrow G$.

EXAMPLE 9.9. Suppose we are given two maps $f, g : X \rightarrow Y$ such that the induced maps $\mathbb{Z}_{tr}(X) \rightarrow \mathbb{Z}_{tr}(Y)$ are \mathbb{A}^1 -homotopic in the sense of 9.8. By the Yoneda lemma, this is equivalent to saying that f and g are restrictions of some $h \in \text{Cor}(X \times \mathbb{A}^1, Y)$, i.e., that f and g are \mathbb{A}^1 -homotopic maps in the sense of 2.25.

LEMMA 9.10. *Let $f, g : F \rightarrow G$ be two maps between étale sheaves with transfers. If f and g are \mathbb{A}^1 -homotopic, then $f = g$ in $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$.*

PROOF. Any two sections of $\mathbb{A}^1 \rightarrow \text{Spec } k$ yield the same map $R \rightarrow R_{tr}(\mathbb{A}^1)$ in the localized category $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$, namely the inverse of the \mathbb{A}^1 -weak equivalence $R_{tr}(\mathbb{A}^1) \rightarrow R$. Therefore the maps:

$$F \xrightarrow[F \times 1]{F \times 0} R_{tr}(\mathbb{A}^1) \otimes_{L, \acute{e}t}^{tr} F \xrightarrow{h} G$$

are the same in the localized category. □

There is a mistake in the proof of the corresponding lemma 3.2.5 in [TriCa] as the proof there assumes that $\mathbb{Z}_{tr}(\mathbb{A}^1)$ is flat in Cor_k . If we replace \otimes by \otimes_L in *loc. cit.*, the proof goes through as written.

COROLLARY 9.11. *Every \mathbb{A}^1 -homotopy equivalence is an \mathbb{A}^1 -weak equivalence.*

Our next goal is to show that, $F \rightarrow C_*F$ is always an \mathbb{A}^1 -weak equivalence (see 9.15 below). Hence $F \cong C_*F$ in $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$.

By the (direct sum) total complex $\text{Tot}(B)$ of a double complex B , we mean the cochain complex with n^{th} term $\bigoplus_{p+q=n} B^{p,q}$; see [Wei94, 1.2.6].

LEMMA 9.12. *Let $f : B \rightarrow B'$ be a map of double complexes which are vertically bounded above in the sense that there is a Q so that $B^{*,q} = (B')^{*,q} = 0$ for all $q \geq Q$. Suppose that the restriction of f to each row is an \mathbb{A}^1 -weak equivalence, and that $\text{Tot}(B)$ and $\text{Tot}(B')$ are bounded above.*

Then $\text{Tot}(B) \rightarrow \text{Tot}(B')$ is an \mathbb{A}^1 -weak equivalence.

PROOF. Let $S(n)$ be the double subcomplex of B consisting of the $B^{p,q}$ for $q \geq n$. Then $\text{Tot}S(n+1)$ is a subcomplex of $\text{Tot}S(n)$ whose cokernel is a shift of the n^{th} row of B . If $S'(n)$ is defined similarly, then each $\text{Tot}S(n) \rightarrow \text{Tot}S'(n)$

is an \mathbb{A}^1 -weak equivalence by induction on n . Now $Sh_{\acute{e}t}(Cor_k, R)$ satisfies (AB4), meaning that \oplus , and hence Tot , is exact. Hence there is a short exact sequence of complexes

$$0 \longrightarrow \bigoplus_{n=1}^{\infty} \text{Tot} S(n) \xrightarrow{id\text{-shift}} \bigoplus_{n=1}^{\infty} \text{Tot} S(n) \longrightarrow \text{Tot} B \longrightarrow 0$$

and similarly for B' . Since $\bigoplus \text{Tot} S(n) \rightarrow \bigoplus \text{Tot} S'(n)$ is an \mathbb{A}^1 -weak equivalence, so is $\text{Tot} B \rightarrow \text{Tot} B'$. \square

COROLLARY 9.13. *If $f : C \rightarrow C'$ is a morphism of bounded above complexes, and $f_n : C_n \rightarrow C'_n$ is in $W_{\mathbb{A}}$ for every n , then f is in $W_{\mathbb{A}}$.*

PROOF. This is a special case of 9.12. \square

LEMMA 9.14. *For every F and every n , the map $F \xrightarrow{s} \underline{\text{Hom}}(R_{tr}(\Delta^n), F) = C_n(F)$ is an \mathbb{A}^1 -homotopy equivalence. A fortiori, it is an \mathbb{A}^1 -weak equivalence.*

PROOF. Since Δ^n is isomorphic to \mathbb{A}^n as a scheme, we have $C_n(F) \cong C_1 C_{n-1}(F)$. Thus we may suppose that $n = 1$. We define a map $m : C_1 F \rightarrow C_2 F$ as follows. For each X , the map

$$m_X : C_1(F)(X) = F(X \times \mathbb{A}^1) \rightarrow F(X \times \mathbb{A}^2) = C_2(F)$$

is induced by the multiplication map $\mathbb{A}^2 \rightarrow \mathbb{A}^1$ by crossing it with X and applying F . Since $C_2 F = \underline{\text{Hom}}(R_{tr}(\mathbb{A}^1), C_1 F)$, the adjunction of 8.2 associates to m a map $h : C_1 F \otimes^{tr} R_{tr}(\mathbb{A}^1) \rightarrow C_1 F$. Similarly the inclusions $\mathbb{A}^1 \times \{i\} \subset \mathbb{A}^2$ induce maps $\eta_i : C_2 F \rightarrow C_1 F$, and the compositions $\eta_i m : C_1 F \rightarrow C_1 F$ are adjoint to the restriction of h along $i : R \rightarrow R_{tr}(\mathbb{A}^1)$. Hence h induces an \mathbb{A}^1 -homotopy between the identity ($\eta_1 m$) and the composite

$$C_1 F \xrightarrow{\partial_0} F \xrightarrow{s} C_1 F,$$

corresponding to $\eta_0 m$. Since $\partial_0 s$ is the identity on F , s and ∂_0 are inverse \mathbb{A}^1 -homotopy equivalences. They are \mathbb{A}^1 -weak equivalences by 9.11. \square

LEMMA 9.15. *For every bounded above complex F of sheaves of R -modules with transfers, the morphism $F \rightarrow C_*(F)$ is an \mathbb{A}^1 -weak equivalence. Hence $F \cong C_*(F)$ in $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$.*

PROOF. By 9.12, we may assume that F is a sheaf. Consider the diagram whose rows are chain complexes

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & F \\ & & \downarrow & & \downarrow & & \downarrow = \\ \cdots & \xrightarrow{0} & F & \xrightarrow{1} & F & \xrightarrow{0} & F \\ & & \downarrow \simeq_{\mathbb{A}^1} & & \downarrow \simeq_{\mathbb{A}^1} & & \downarrow \simeq_{\mathbb{A}^1} \\ \cdots & \longrightarrow & C_2 F & \longrightarrow & C_1 F & \longrightarrow & F. \end{array}$$

The first two rows are quasi-isomorphic. Now $F \simeq_{\mathbb{A}^1} C_n(F)$ by 9.14. Using 9.13, we see that the second and third rows are \mathbb{A}^1 -weak equivalent. \square

EXAMPLE 9.16. The identity map on \mathcal{O} is \mathbb{A}^1 -homotopic to zero by 2.23 and 9.15. Hence \mathcal{O} is isomorphic to zero in $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k)$. When $\text{char } k = \ell > 0$ the Artin-Schrier sequence of étale sheaves [Mil80, II 2.18(c)]

$$0 \longrightarrow \mathbb{Z}/\ell \longrightarrow \mathcal{O} \xrightarrow{1-\phi} \mathcal{O} \longrightarrow 0$$

shows that $\mathbb{Z}/\ell \cong 0$ in $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k)$. Here R may be either \mathbb{Z} or \mathbb{Z}/ℓ .

Étale \mathbb{A}^1 -local complexes

In this section we will show that $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, \mathbb{Z}/m)$ can be identified with the full subcategory \mathcal{L} of \mathbb{A}^1 -local complexes in $\mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, \mathbb{Z}/m))$.

DEFINITION 9.17. An object L in \mathbf{D}^- is called \mathbb{A}^1 -local if for all \mathbb{A}^1 -weak equivalences $K' \rightarrow K$ the induced map $\text{Hom}(K, L) \rightarrow \text{Hom}(K', L)$ is bijective. We write \mathcal{L} for the full subcategory of \mathbb{A}^1 -local objects in \mathbf{D}^- .

It is easy to see that \mathcal{L} forms a thick triangulated subcategory of \mathbf{D}^- .

REMARK 9.18. We will see in 9.31 below that C_* is a functor from \mathbf{D}^- to \mathcal{L} , provided that $R = \mathbb{Z}/m$ and $cd_m(k) < \infty$. Moreover, $\text{Hom}(C_*(F), L) \cong \text{Hom}(F, L)$ for every L in \mathcal{L} and F in \mathbf{D}^- , by 9.15 and definition 9.17. Hence C_* is the left adjoint to the inclusion $\mathcal{L} \subset \mathbf{D}^-$.

LEMMA 9.19. *If L is \mathbb{A}^1 -local then for every K in \mathbf{D}^-*

$$\text{Hom}_{\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k,R)}(K, L) = \text{Hom}_{\mathbf{D}^-}(K, L).$$

Hence the natural functor $\mathcal{L} \rightarrow \mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ is full and faithful.

PROOF. By the calculus of fractions [Wei94, 10.3.7], the left side consists of equivalence classes of diagrams $K \xleftarrow{s} K' \longrightarrow L$ with s in $W_{\mathbb{A}^1}$. It suffices to show that if $K' \rightarrow K$ is an \mathbb{A}^1 -weak equivalence then $\text{Hom}(K, L) = \text{Hom}(K', L)$. But this holds since L is \mathbb{A}^1 -local. \square

LEMMA 9.20. *An object L in \mathbf{D}^- is \mathbb{A}^1 -local if and only if $\text{Hom}(R_{tr}(X)[n], L) \rightarrow \text{Hom}(R_{tr}(X \times \mathbb{A}^1)[n], L)$ is an isomorphism for all X and n .*

PROOF. Let \mathcal{K} be the full subcategory of all K for which $\text{Hom}(K[n], L) = 0$ for all n . Clearly, \mathcal{K} is a thick subcategory of \mathbf{D}^- and it is closed under direct sums and shifts. Under the given hypothesis, \mathcal{K} contains the cone of every map $R_{tr}(X \times \mathbb{A}^1) \rightarrow R_{tr}(X)$. By definition, $\mathcal{E}_{\mathbb{A}^1}$ is a subcategory of \mathcal{K} , i.e., L is \mathbb{A}^1 -local. \square

LEMMA 9.21. *If $f : L \rightarrow L'$ is an \mathbb{A}^1 -weak equivalence and L, L' are \mathbb{A}^1 -local then f is an isomorphism in \mathbf{D}^- , i.e., a quasi-isomorphism of complexes of étale sheaves with transfers.*

PROOF. By definition, f induces bijections $\mathrm{Hom}(L', L) \cong \mathrm{Hom}(L, L)$ and $\mathrm{Hom}(L', L') \cong \mathrm{Hom}(L, L')$. Hence there is a unique $g : L' \rightarrow L$ so that $fg = 1_L$, and $f(gf) = (fg)f = f$ implies that $gf = 1_{L'}$. \square

DEFINITION 9.22. An étale sheaf with transfers F is **strictly \mathbb{A}^1 -homotopy invariant** if the map $H_{\acute{e}t}^n(X, F) \rightarrow H_{\acute{e}t}^n(X \times \mathbb{A}^1, F)$ is bijective for all smooth X and every $n \in \mathbb{N}$. In particular for $n = 0$ we must have that F is homotopy invariant (2.15).

LEMMA 9.23. ([SGA4, XV 2.2]) *If R is of torsion prime to char k then any locally constant sheaf of R -modules is strictly \mathbb{A}^1 -homotopy invariant.*

LEMMA 9.24. *Let F be an étale sheaf of R -modules with transfers. Then F is \mathbb{A}^1 -local if and only if F is strictly \mathbb{A}^1 -homotopy invariant.*

PROOF. By 6.23 or 6.24, we have

$$\mathrm{Hom}_{\mathbf{D}^-}(R_{tr}(X), F[i]) = \mathrm{Ext}_{\mathrm{Sh}_{\acute{e}t}(\mathrm{Cor}_k, R)}^i(R_{tr}(X), F) = H_{\acute{e}t}^i(X, F)$$

for every smooth X . Since $R_{tr}(X \times \mathbb{A}^1)[n] \rightarrow R_{tr}(X)[n]$ is an \mathbb{A}^1 -weak equivalence for all n , 9.20 shows that F is \mathbb{A}^1 -local if and only if the induced map

$$H_{\acute{e}t}^{-n}(X, F) = \mathrm{Hom}(R_{tr}(X)[n], F) \rightarrow \mathrm{Hom}(R_{tr}(X \times \mathbb{A}^1)[n], F) = H_{\acute{e}t}^{-n}(X \times \mathbb{A}^1, F)$$

is an isomorphism, that is, if and only if F is strictly \mathbb{A}^1 -homotopy invariant. \square

Here is a special case of 9.24 which includes the sheaves $\mu_n^{\otimes q}$. It follows by combining 9.23 with 9.24.

COROLLARY 9.25. *Let M be a locally constant étale sheaf of torsion prime to char k . Then M is \mathbb{A}^1 -local.*

We now make the running assumption that R is a commutative ring and that $cd_R(k) < \infty$, i.e., k is a field having finite étale cohomological dimension for coefficients in R . If $R = \mathbb{Z}/m$ we will write $cd_m(k)$ for $cd_R(k)$. This assumption allows us to invoke a classical result from [SGA4].

LEMMA 9.26. ([SGA4], [Mil80]) *Let X be a scheme of finite type over k . If k has finite R -cohomological dimension d then $cd_R(X) \leq d + 2 \dim_k X$.*

COROLLARY 9.27. *Set $n_X = cd_R(k) + 2 \dim_k X$. Then $\mathrm{Ext}^n(R_{tr}(X), F) = 0$ when $n \geq n_X$.*

PROOF. $\mathrm{Ext}^n(R_{tr}(X), F) \cong H_{\acute{e}t}^n(X, F)$ by 6.24. \square

REMARK 9.28. If B is \mathbb{A}^1 -local then so is the complex $\underline{R}\mathrm{Hom}(R_{tr}X, B)$ of 8.21. Indeed, B is strictly \mathbb{A}^1 -homotopy invariant by 9.24, so by 6.25, we have:

$$\begin{aligned} H^* \underline{R}\mathrm{Hom}(R_{tr}X, B)(U) &\cong \mathbb{H}^*(U \times X, B) \\ &\cong \mathbb{H}^*(U \times X \times \mathbb{A}^1, B) \cong H^* \underline{R}\mathrm{Hom}(R_{tr}X, B)(U \times \mathbb{A}^1). \end{aligned}$$

If C is a cochain complex of presheaves, each cohomology $H^n(C)$ is a presheaf. We write $a_{\acute{e}t}H^n(C)$ for its associated étale sheaf.

LEMMA 9.29. *Assume that $cd_R(k) < \infty$. Then for every (bounded above) chain complex C there is a bounded, convergent spectral sequence:*

$$E_2^{p,q} = \text{Ext}^p(R_{tr}(X), a_{\acute{e}t}H^q(C)) \implies \text{Hom}_{\mathbf{D}^-}(R_{tr}(X), C[p+q]).$$

PROOF. This is well-known; see [Wei94, 5.7.9]. The spectral sequence is bounded, and hence converges, by 9.27. \square

PROPOSITION 9.30. *Let C be a bounded above cochain complex of étale sheaves of R -modules with transfers, where $cd_R(k) < \infty$. If the sheaves $a_{\acute{e}t}H^n(C)$ are all strictly \mathbb{A}^1 -homotopy invariant, then C is \mathbb{A}^1 -local.*

PROOF. Let C be a complex of étale sheaves with transfers. By 9.20, it suffices to prove that $\text{cone}(f)$ is in this class when f is the projection $R_{tr}(X \times \mathbb{A}^1) \rightarrow R_{tr}(X)$. The map f induces a morphism between the spectral sequences of 9.29 for X and $X \times \mathbb{A}^1$. Because the sheaves $L = a_{\acute{e}t}H^q C$ are strictly \mathbb{A}^1 -homotopy invariant, they are \mathbb{A}^1 -local by 9.24. Thus

$$\begin{aligned} \text{Ext}^p(R_{tr}(X), L) &= \text{Hom}_{\mathbf{D}^-}(R_{tr}(X)[-p], L) \\ &\cong \text{Hom}_{\mathbf{D}^-}(R_{tr}(X \times \mathbb{A}^1)[-p], L) = \text{Ext}^p(R_{tr}(X \times \mathbb{A}^1), L). \end{aligned}$$

Hence the morphism of spectral sequences is an isomorphism on all E_2 terms. By the Comparison Theorem [Wei94, 5.2.12], f induces an isomorphism from $\text{Hom}_{\mathbf{D}^-}(R_{tr}(X)[n], C)$ to $\text{Hom}_{\mathbf{D}^-}(R_{tr}(X \times \mathbb{A}^1)[n], C)$ for each n . Done. \square

LEMMA 9.31. *Suppose that $1/m \in k$ and $cd_m(k) < \infty$. If K is a bounded above complex of étale sheaves of \mathbb{Z}/m -modules with transfers, then $\text{Tot}C_*(K)$ is \mathbb{A}^1 -local.*

PROOF. Set $C = \text{Tot}C_*(K)$. By 2.19, each $H^i C$ is an \mathbb{A}^1 -homotopy invariant presheaf of \mathbb{Z}/m -modules with transfers. By the Rigidity Theorem 7.20, the sheaf $a_{\acute{e}t}H^i C$ is locally constant. By 9.23, $a_{\acute{e}t}H^i C$ is strictly \mathbb{A}^1 -homotopy invariant. Finally, 9.30 lets us conclude that C is \mathbb{A}^1 -local. \square

Combining 9.21 with 9.31, we obtain:

COROLLARY 9.32. *If F is \mathbb{A}^1 -local then $F \cong C_* F$ in \mathbf{D}^- .*

COROLLARY 9.33. *If $1/m \in k$ then $\mathbb{Z}/m(q)$ is \mathbb{A}^1 -local for all q .*

PROOF. Take K to be $(\mathbb{Z}/m)_{tr} \mathbb{G}_m^{\wedge q}[-q]$; $\mathbb{Z}/m(q) = C_* K$ by definition 3.1. \square

DEFINITION 9.34. If $1/m \in k$, let \mathcal{L} denote the full subcategory of \mathbf{D}^- consisting of \mathbb{A}^1 -local complexes of \mathbb{Z}/m -modules with transfers. If E and F are \mathbb{A}^1 -local, we set $E \otimes_{\mathcal{L}} F = \text{Tot}C_*(E \otimes_{L, \acute{e}t}^tr F)$. By 9.31, $E \otimes_{\mathcal{L}} F$ is \mathbb{A}^1 -local, so $\otimes_{\mathcal{L}}$ is a bifunctor from $\mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$.

Recall from 6.9 that the category of locally constant étale sheaves of \mathbb{Z}/m -modules is equivalent to the category $\mathbf{Mod}(G, \mathbb{Z}/m)$ of discrete \mathbb{Z}/m -modules over the Galois group $G = \text{Gal}(k_{\text{sep}}/k)$. Let $\mathbf{D}^-(G, \mathbb{Z}/m)$ denote the (bounded above) derived category of such modules. There is a triangulated functor π^* from $\mathbf{D}^-(G, \mathbb{Z}/m)$ to $\mathbf{D}^- = \mathbf{D}^-(\text{Sh}_{\text{ét}}(\text{Cor}_k, \mathbb{Z}/m))$.

THEOREM 9.35. *If $1/m \in k$, $(\mathcal{L}, \otimes_{\mathcal{L}})$ is a tensor triangulated category and the functors*

$$\mathbf{D}^-(G, \mathbb{Z}/m) \xrightarrow{\pi^*} \mathcal{L} \longrightarrow \mathbf{D}^-[W_{\mathbb{A}}^{-1}] = \mathbf{DM}_{\text{ét}}^{\text{eff}, -}(k, \mathbb{Z}/m)$$

are equivalences of tensor triangulated categories.

PROOF. Clearly, \mathcal{L} is a thick subcategory of \mathbf{D}^- . By 9.19, the functor $\mathcal{L} \rightarrow \mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ is fully faithful. By 9.31, every object of $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ is isomorphic to an object of \mathcal{L} . Hence \mathcal{L} is equivalent to $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ as a triangulated category.

By 9.6, $\mathbf{DM}_{\text{ét}}^{\text{eff}, -}(k, \mathbb{Z}/m)$ is a tensor triangulated category. Using the first part of this proof, we conclude that \mathcal{L} is a tensor triangulated category as well. Moreover, if E and F are \mathbb{A}^1 -local, then $E \otimes_{\mathcal{L}} F$ is isomorphic to $E \otimes_{L, \text{ét}}^{\text{tr}} F$ in $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ by 9.15, so the induced tensor operation on \mathcal{L} is isomorphic to $\otimes_{\mathcal{L}}$.

Next we consider π^* . It is easy to see from 6.9 and 6.11 that π^* induces an equivalence between $\mathbf{D}^-(G, \mathbb{Z}/m)$ and the full subcategory of complexes of locally constant sheaves in \mathbf{D}^- . By exercise 8.20, π^* sends $\otimes_{\mathbb{Z}/m}^{\mathbb{L}}$ to $\otimes_{L, \text{ét}}^{\text{tr}}$. It suffices to show that every \mathbb{A}^1 -local complex F is isomorphic to such a complex. By 9.15, 9.31, and 9.21 $F \rightarrow C_*F$ is a quasi-isomorphism. By 2.19, each $a_{\text{ét}}H^iF$ is \mathbb{A}^1 -homotopy invariant. By 7.20 the sheaves $a_{\text{ét}}H^iF$ are locally constant. Hence the canonical map $F \rightarrow \pi^*\pi_*F$ is a quasi-isomorphism of complexes of étale sheaves. But π_*F is a complex of modules in $\mathbf{Mod}(G, \mathbb{Z}/m)$. \square

Étale motivic cohomology and algebraic singular homology

There are two ways one might define an étale version of motivic cohomology. One way, which is natural from the viewpoint of these notes, is to use the morphisms in the triangulated category $\mathbf{DM}_{\acute{e}t}^-$, namely to define the integral cohomology group indexed by (p, q) as $\mathrm{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}(q)[p])$, and similarly for cohomology with coefficients in an abelian group A . The second approach, due to Lichtenbaum, is to take the étale hypercohomology of the complex $\mathbb{Z}(q)$.

DEFINITION 10.1. For any abelian group A , we define the **étale (or Lichtenbaum) motivic cohomology** of X as the hypercohomology of $A(q)$:

$$H_L^{p,q}(X, A) = \mathbb{H}_{\acute{e}t}^p(X, A(q)|_{X_{\acute{e}t}}).$$

If $q < 0$ then $H_L^{p,q}(X, A) = 0$, because $A(q) = 0$. If $q = 0$ then $H_L^{p,0}(X, A) \cong H_{\acute{e}t}^p(X, A)$, because $A(0) = A$.

The two definitions agree in some cases of interest. We will see in 10.7 below that $H_L^{p,q}(X, \mathbb{Z}/n) \cong \mathrm{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}/n(q)[p])$ when $1/n \in k$. Even further on, in 14.27, we will see that $H_L^{p,q}(X, \mathbb{Q}) \cong \mathrm{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Q}(q)[p])$. However, the two definitions do not agree for ℓ -torsion coefficients, for $\ell = \mathrm{char}(k)$. Indeed, for $q = 0$ we have $\mathrm{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}/\ell[p]) = 0$ in characteristic ℓ by 9.16, yet the groups $H_L^{p,0}(X, \mathbb{Z}/\ell) \cong H_{\acute{e}t}^p(X, \mathbb{Z}/\ell)$ can certainly be nonzero.

By corollary 6.4 we have $H_L^{p,1}(X, \mathbb{Z}/n) \cong H_{\acute{e}t}^p(X, \mu_n)$ when $1/n \in k$. Here is the generalization to all q .

THEOREM 10.2. *Let n be an integer prime to the characteristic of k . Then:*

$$H_L^{p,q}(X, \mathbb{Z}/n) = H_{\acute{e}t}^p(X, \mu_n^{\otimes q}) \quad q \geq 0, p \in \mathbb{Z}.$$

By 6.4 there is a quasi-isomorphism $\mu_n \rightarrow \mathbb{Z}/n(1)$ of complexes of étale sheaves. Because μ_n and the terms of $\mathbb{Z}/n(1)$ are flat as sheaves of \mathbb{Z}/n -modules, there is a morphism $\mu_n^{\otimes q} \rightarrow (\mathbb{Z}/n)(1)^{\otimes q}$ in the category of complexes of étale sheaves of \mathbb{Z}/n -modules. Combining with the multiplication of 3.11 gives a map

$$\mu_n^{\otimes q} \longrightarrow (\mathbb{Z}/n)(1)^{\otimes q} \longrightarrow (\mathbb{Z}/n)(q).$$

We may now reformulate theorem 10.2 as follows.

THEOREM 10.3. *The map $\mu_n^{\otimes q} \rightarrow \mathbb{Z}/n(q)$ is a quasi-isomorphism of complexes of étale sheaves.*

PROOF. The theorem is true for $q = 1$ by 6.4. By 9.25 and 9.31, both $\mu_n^{\otimes q}$ and $\mathbb{Z}/n(q)$ are \mathbb{A}^1 -local. We will show that the map $\mu_n^{\otimes q} \rightarrow \mathbb{Z}/n(q)$ is an \mathbb{A}^1 -weak equivalence in 10.6 below. By 9.21, it is also a quasi-isomorphism. \square

Let R be any commutative ring. Recall that $R(n) = R \otimes_{\mathbb{Z}} \mathbb{Z}(n)$. Clearly, the multiplication map $\mathbb{Z}(m) \otimes_{\mathbb{Z}} \mathbb{Z}(n) \rightarrow \mathbb{Z}(m+n)$ of 3.11 induces a map $R(m) \otimes_R R(n) \rightarrow R(m+n)$.

PROPOSITION 10.4. *The multiplication map $R(m) \otimes R(n) \rightarrow R(m+n)$ factors through a map $\mu : R(m) \otimes^{tr} R(n) \rightarrow R(m+n)$.*

$$\begin{array}{ccc} R(m) \otimes_R R(n) & \xrightarrow{\text{mult.}} & R(m+n) \\ \downarrow 8.9 & \nearrow \mu & \\ R(m) \otimes^{tr} R(n) & & \end{array}$$

PROOF. We first reinterpret the left vertical map in simplicial language. Recall that by definition 3.1, $R(n)[n] = C_*(R_{tr}(\mathbb{G}_m^{\wedge n}))$. Let us write A_{\bullet}^n for the underlying simplicial presheaf, viz., $A_{\bullet}^n(U) = \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})(U \times \Delta^{\bullet})$, and write the associated unnormalized chain complex as A_*^n . By 8.9, we have a natural map of bisimplicial presheaves $A_{\bullet}^m \otimes_R A_{\bullet}^n \rightarrow A_{\bullet}^m \otimes^{tr} A_{\bullet}^n$, and a map of their diagonal chain complexes, $(A^m \otimes_R A^n)_* \rightarrow (A^m \otimes^{tr} A^n)_*$. As in 3.11, the Eilenberg-Zilber theorem yields quasi-isomorphisms ∇ fitting into a commutative diagram:

$$\begin{array}{ccccc} R(m) \otimes_R R(n)[m+n] & \xrightarrow{=} & A_*^m \otimes_R A_*^n & \xrightarrow{\nabla} & (A^m \otimes_R A^n)_* \\ \downarrow 8.9 & & \downarrow 8.9 & & \downarrow 8.9 \\ R(m) \otimes^{tr} R(n)[m+n] & \xrightarrow{=} & A_*^m \otimes^{tr} A_*^n & \xrightarrow{\nabla} & (A^m \otimes^{tr} A^n)_* \end{array}$$

Comparing with 3.11, we see that it suffices to find a simplicial map for all X and Y ,

$$(10.4.1) \quad \text{diag}(C_{\bullet} R_{tr}(X) \otimes^{tr} C_{\bullet} R_{tr}(Y)) \longrightarrow C_{\bullet} R_{tr}(X \times Y)$$

compatible with the corresponding construction 3.10 for \otimes_R . The map μ will be the composite of ∇ and the map induced by 10.4.1.

Let F be any presheaf with transfers. Definitions 8.2 and 2.14 imply that $C_n(F) \cong \underline{\text{Hom}}(R_{tr}(\Delta^n), F)$ as presheaves and that $C_{\bullet}(F) \cong \underline{\text{Hom}}(R_{tr}(\Delta_{\bullet}^n), F)$ as simplicial presheaves. Using these identifications, we define the map 10.4.1 in degree n as the composition:

$$\begin{aligned} & C_n(R_{tr}(X)) \otimes^{tr} C_n(R_{tr}(Y)) \\ &= \underline{\text{Hom}}(R_{tr}(\Delta^n), R_{tr}(X)) \otimes^{tr} \underline{\text{Hom}}(R_{tr}(\Delta^n), R_{tr}(Y)) \\ &\xrightarrow{8.5} \underline{\text{Hom}}(R_{tr}(\Delta^n \times \Delta^n), R_{tr}(X \times Y)) \xrightarrow{\text{diagonal}} \underline{\text{Hom}}(R_{tr}(\Delta^n), R_{tr}(X \times Y)) \\ &= C_n(R_{tr}(X \times Y)). \end{aligned}$$

Since $\underline{Hom}(R_{tr}(\Delta^n \times \Delta^n), R_{tr}(X \times Y))(U) = R_{tr}(X \times Y)(U \times \Delta^n \times \Delta^n)$, the above composition is the right vertical composition in the following commutative diagram (see 8.5):

$$\begin{array}{ccc}
 R_{tr}(X)(U \times \Delta^n) \otimes_R R_{tr}(Y)(U \times \Delta^n) & \xrightarrow{8.9} & (C_n R_{tr}(X) \otimes^{tr} C_n R_{tr}(Y))(U) \\
 \downarrow \otimes & & \downarrow 8.5 \\
 R_{tr}(X \times Y)(U \times U \times \Delta^n \times \Delta^n) & \xrightarrow{\text{diag}(U)} & R_{tr}(X \times Y)(U \times \Delta^n \times \Delta^n) \\
 \searrow \text{diag}(U \times \Delta^n) & & \downarrow \text{diag}(\Delta^n) \\
 & & R_{tr}(X \times Y)(U \times \Delta^n).
 \end{array}$$

Since the left composite is the degree n part of construction 3.10, this shows that the triangle in 10.4 commutes. \square

PROPOSITION 10.5. *The map $\mathbb{Z}/n(1)^{\otimes_{L^1} q} \rightarrow \mathbb{Z}/n(q)$ is an \mathbb{A}^1 -weak equivalence in $\mathbf{D}^-(Sh_{\acute{e}t}(Cor_k, \mathbb{Z}/n))$.*

PROOF. The assertion follows from the diagram in figure 10.1, remembering that by definition $\mathbb{Z}/n(q)$ is $C_*(\mathbb{Z}/n)_{tr}(\mathbb{G}_m^{\wedge q})[-q]$. \square

$$\begin{array}{ccc}
 \mathbb{Z}/n(1)^{\otimes_{L^1} q} & \xrightarrow{\quad} & \mathbb{Z}/n(q) \\
 \uparrow \simeq_{\mathbb{A}^1} \quad 9.5+9.15 & & \uparrow \quad 9.15 \simeq_{\mathbb{A}^1} \\
 (\mathbb{Z}/n)_{tr}(\mathbb{G}_m)[-1]^{\otimes_{L^1} q} & & (\mathbb{Z}/n)_{tr}(\mathbb{G}_m^{\wedge q})[-q] \\
 \searrow 8.7 \simeq & & \nearrow 8.10 = \\
 & ((\mathbb{Z}/n)_{tr}(\mathbb{G}_m))^{\otimes_{tr} q}[-q] &
 \end{array}$$

FIGURE 10.1. The factorization in proposition 10.5

PROPOSITION 10.6. *The map $\mu_n^{\otimes q} \rightarrow \mathbb{Z}/n(q)$ is an \mathbb{A}^1 -weak equivalence in $\mathbf{D}^-(Sh_{\acute{e}t}(Cor_k, \mathbb{Z}/n))$.*

PROOF. Consider the following diagram, in which \otimes^{tr} and \otimes_L^{tr} are to be understood in \mathbb{Z}/n -modules.

$$\begin{array}{ccccc}
 & & \mu_n^{\otimes_L^{tr} q} & \xrightarrow{\simeq} & \mathbb{Z}/n(1)^{\otimes_L^{tr} q} \\
 & & \downarrow \simeq & & \downarrow \mu_1 \\
 & & \mu_n^{\otimes q} & \xrightarrow{\simeq} & \mu_n^{\otimes^{tr} q} \\
 \mu_n^{\otimes q} & \xrightarrow{\simeq} & \mu_n^{\otimes^{tr} q} & \longrightarrow & \mathbb{Z}/n(1)^{\otimes^{tr} q} \xrightarrow{\mu} \mathbb{Z}/n(q) \\
 & & & & \swarrow \mu_1 \circ \mu
 \end{array}$$

We already know that the top map is a quasi-isomorphism by 6.4 and 8.16. Lemma 8.13 proves that the bottom left map $\mu_n^{\otimes q} \rightarrow \mu_n^{\otimes^{tr} q}$ is a quasi-isomorphism. Lemma 8.18 proves that the left vertical map is a quasi-isomorphism. Hence the assertion follows from proposition 10.5. \square

Recall that when $1/n \in k$ we have $\mathbf{DM}_{\acute{e}t}^- = \mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, \mathbb{Z}/n)$.

PROPOSITION 10.7. *If $1/n \in k$ then*

$$H_L^{p,q}(X, \mathbb{Z}/n) \cong \text{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}/n(q)[p]).$$

PROOF. Since $A = \mathbb{Z}/n(q)$ is \mathbb{A}^1 -local by 9.33, the right side is

$$\begin{aligned}
 \text{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}/n(q)[p]) &= \text{Hom}_{\mathbf{D}^-}(\mathbb{Z}_{tr}(X), \mathbb{Z}/n(q)[p]) \\
 &= \text{Ext}^p(\mathbb{Z}_{tr}(X), \mathbb{Z}/n(q)).
 \end{aligned}$$

By 6.25, this Ext group is $\mathbb{H}_{\acute{e}t}^p(X, \mathbb{Z}/n(q))$, which is the left side. \square

As a bonus for all our hard work, we are able to give a nice interpretation of Suslin's algebraic singular homology. Recall that $R_{tr}(X) = \mathbb{Z}_{tr}(X) \otimes R$.

DEFINITION 10.8. We define the algebraic singular homology of X by:

$$H_p^{\text{sing}}(X, R) = H_p(C_* R_{tr}(X)(\text{Spec } k)).$$

By remark 7.4, $H_0^{\text{sing}}(X, \mathbb{Z})$ agrees with the group $H_0^{\text{sing}}(X/\text{Spec } k)$ of lecture 7. It is immediate from 5.2 that:

$$H^{p,q}(\text{Spec } k, R) = H_{q-p}^{\text{sing}}(\mathbb{G}_m^{\wedge q}, R).$$

Notice that $R_{tr}(\mathbb{G}_m^{\wedge q})$ is well-defined even though $\mathbb{G}_m^{\wedge q}$ is not a scheme.

The following theorem was first proven in [SV96, 7.8] under the assumption of resolution of singularities on k . The proof we give here doesn't need resolution of singularities, so it extends the result to fields of positive characteristic.

THEOREM 10.9. *Let k be a separably closed field and X a smooth scheme over k , and let l be a prime number different from $\text{char } k$. Then there exist natural isomorphisms for all i :*

$$H_p^{\text{sing}}(X, \mathbb{Z}/l)^* \cong H_{\acute{e}t}^p(X, \mathbb{Z}/l)$$

where the $*$ denotes the dual vector space over \mathbb{Z}/l .

It is amusing to note that this implies that $H_{\acute{e}t}^i(X, \mathbb{Z}/l)$ is finite, because it is a countable-dimensional dual module.

To prove 10.9, we need one more lemma. To clarify the role of the coefficient ring R , we will write \mathbf{D}_R^- for $\mathbf{D}^-(Sh_{\acute{e}t}(Cor_k, R))$, so that $\mathbf{D}_{\mathbb{Z}}^-$ is just the usual derived category of $Sh_{\acute{e}t}(Cor_k)$.

LEMMA 10.10. *Let k be a separably closed field and C a bounded above chain complex of étale sheaves of R -modules with transfers. Assume that the cohomology sheaves of C are locally constant and projective (as R -modules). Then for any $n \in \mathbb{Z}$ we have:*

$$\mathrm{Hom}_{\mathbf{D}_R^-}(C, R[n]) = \mathrm{Hom}_{R\text{-mod}}(H^n(C)(\mathrm{Spec} k), R).$$

PROOF. For simplicity, let us write Ext^* for Ext in the category $Sh_{\acute{e}t}(Cor_k, R)$. (There are enough injectives to define Ext by 6.19.)

If P is a summand of $\bigoplus_{\alpha} R$, then $\mathrm{Ext}^n(P, R)$ injects into

$$\mathrm{Ext}^n(\bigoplus_{\alpha} R, R) = \prod \mathrm{Ext}^n(R, R) = \prod \mathrm{Ext}^n(R_{tr}(\mathrm{Spec} k), R).$$

But $\mathrm{Ext}^n(R_{tr}(\mathrm{Spec} k), R) = H_{\acute{e}t}^n(\mathrm{Spec} k, R)$ by 6.24 and this vanishes if $n \neq 0$ as k is separably closed. If $n = 0$, this calculation yields $\mathrm{Ext}^0(R, R) = R$ and $\mathrm{Ext}^0(P, R) = \mathrm{Hom}_{R\text{-mod}}(P, R)$.

Now recall that $\mathrm{Ext}^n(F, R) = \mathrm{Hom}_{\mathbf{D}_R^-}(F, R[n])$ for every sheaf F ; see [Wei94, 10.7.5]. More generally, if $R \rightarrow I^*$ is an injective resolution then the total Hom cochain complex $\mathbb{R}\mathrm{Hom}(C, R)$ of $\mathrm{Hom}^*(C, I[n])$ satisfies

$$H^n \mathbb{R}\mathrm{Hom}(C, R) \cong \mathrm{Hom}_{\mathbf{D}_R^-}(C, R[n]).$$

(See [Wei94, 10.7.4].) Since $\mathrm{Hom}^*(C, I[n])$ is a bounded double complex, it gives rise to a convergent spectral sequence which, as in [Wei94, 5.7.9], may be written

$$E_2^{pq} = \mathrm{Ext}^p(H^q C, R) \implies H^{p+q} \mathbb{R}\mathrm{Hom}(C, R) = \mathrm{Hom}_{\mathbf{D}_R^-}(C, R[p+q]).$$

The assumption on $H^q C$ makes the spectral sequence collapse to yield $\mathrm{Ext}^0(H^n C, R) \cong \mathrm{Hom}_{\mathbf{D}_R^-}(C, R[n])$, whence the result. \square

PROOF OF 10.9. Taking $R = \mathbb{Z}/l$, this means that all R -modules are projective. Consider the diagram:

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbf{D}_R^-}(C_* R_{tr}(X), R[n]) & \xrightarrow[10.10]{\cong} & \mathrm{Hom}_{R\text{-mod}}(H_n^{sing}(X, R), R) \\ \downarrow 9.25 \cong & & \\ \mathrm{Hom}_{\mathbf{D}_R^-}(R_{tr}(X), R[n]) & \xrightarrow[6.24]{\cong} & H_{\acute{e}t}^n(X, R). \end{array}$$

By 2.19, each $H^n = H^n C_* R_{tr}(X)$ is a homotopy invariant presheaf of \mathbb{Z}/l -modules with transfers. Hence the sheaves $a_{\acute{e}t} H^n$ are locally constant by the Rigidity Theorem 7.20. Hence the top map is an isomorphism by 10.10. Since R is \mathbb{A}^1 -local by 9.25, the left map is an isomorphism by 9.15. The bottom map is an isomorphism by 6.24. \square

COROLLARY 10.11. *Let k be a separably closed field and X a smooth scheme over k , and let n be an integer relatively prime to $\text{char } k$. Then there exist natural isomorphisms for all i :*

$$H_p^{\text{sing}}(X, \mathbb{Z}/n)^* \cong H_{\text{ét}}^p(X, \mathbb{Z}/n)$$

where the $*$ denotes the Pontrjagin dual \mathbb{Z}/n -module.

PROOF. Using the sequences $0 \rightarrow \mathbb{Z}/l \rightarrow \mathbb{Z}/lm \rightarrow \mathbb{Z}/m \rightarrow 0$, the 5-lemma shows that we may assume that n is prime. \square

Part 3

Nisnevich Sheaves with Transfers

LECTURE 11

Standard triples

Our goal in this lecture is to prove proposition 11.1 below, which is one of the main properties of homotopy invariant presheaves with transfers. It (or rather its corollary 11.2) will be used in subsequent lectures to promote results from the Nisnevich topology to the Zariski topology. It depends primarily upon the relative Picard group introduced in lecture 7.

For all of this lecture, F will be a homotopy invariant presheaf with transfers.

Recall that a subgroup A of an abelian group B is called **pure** if $nA = nB \cap A$ for every integer n . A homomorphism $f : A \rightarrow B$ of abelian groups is called **pure injective** if it is injective and $f(A)$ is a pure subgroup of B .

Any semilocal subscheme S of a smooth X is the intersection of the open sets X_α which contain it; by abuse we call S smooth and write $F(S)$ for $\varinjlim F(X_\alpha)$, as in exercise 2.10. (If S is local, this is the stalk of F at the closed point of S .)

PROPOSITION 11.1. *For any smooth semilocal S over k , any Zariski dense open subset $V \subset S$, and any homotopy invariant presheaf with transfers F , the map $F(S) \rightarrow F(V)$ is pure injective.*

The intersection of all such V is the coproduct of the generic points $\text{Spec } E_i$ of S . Hence $F(S)$ injects (as a pure subgroup) into $\bigoplus F(\text{Spec } E_i) = \varinjlim F(V)$.

COROLLARY 11.2. *Let F be a homotopy invariant presheaf with transfers. If $F(\text{Spec } E) = 0$ for every field E over k , then $F_{\text{Zar}} = 0$.*

Here is the proof of proposition 11.1; it is a consequence of a more precise result, theorem 11.3, whose proof will take up most of this lecture.

PROOF. The semilocal scheme S is the intersection of a family X_α of smooth varieties of finite type over k and V is the intersection of dense open subschemes $V_\alpha \subset X_\alpha$. Hence $F(S) \rightarrow F(V)$ is the filtered colimit of the maps $i_\alpha : F(X_\alpha) \rightarrow F(V_\alpha)$. Since the U_α given by 11.3 contains some X_β , the kernel of i_α vanishes in $F(X_\beta)$ and the colimit is an injection. If $a \in F(X_\alpha)$ equals $nb \in F(V_\alpha)$ for some $b \in F(V_\alpha)$, then the image of a in $F(U_\alpha)$, and hence in $F(S)$, is n -divisible. \square

THEOREM 11.3. *Let X be smooth of finite type over a field k and let V be a dense open subset. Then for every finite set of points $x_1, \dots, x_n \in X$ there exists an open neighborhood U of these points such that the restriction $F(X) \rightarrow F(U)$ factors through $F(X) \rightarrow F(V)$. That is, there is a map $F(V) \rightarrow F(U)$ such that the*

following diagram commutes.

$$\begin{array}{ccc}
 F(X) & & \\
 \downarrow & \searrow & \\
 F(V) & \xrightarrow{\exists} & F(U)
 \end{array}$$

EXAMPLE 11.4. If $V \subsetneq X$ is a dense open subset, then $F = \mathbb{Z}_{tr}(X)/\mathbb{Z}_{tr}(V)$ is a presheaf with transfers, but $F(X) \rightarrow F(V)$ is not injective. (1_X is nonzero in $F(X)$ but vanishes in $F(V)$.) This shows that homotopy invariance is necessary in 11.3.

To prepare for the proof of theorem 11.3, we need a technical digression.

DEFINITION 11.5. A **standard triple** is a triple $(\bar{X} \xrightarrow{\bar{p}} S, X_\infty, Z)$ where \bar{p} is a proper morphism of relative dimension 1 and Z and X_∞ are closed subschemes of \bar{X} . The following conditions must be satisfied:

- (1) S is smooth and \bar{X} is normal,
- (2) $\bar{X} - X_\infty$ is quasi-affine and smooth over S ,
- (3) $Z \cap X_\infty = \emptyset$,
- (4) $X_\infty \cup Z$ lies in an affine open neighborhood in \bar{X} .

Given a standard triple as above, we usually write X for $\bar{X} - X_\infty$. Note that \bar{X} is a good compactification of both X and $X - Z$ (see 7.8) by parts (2) and (4).

Conversely, if \bar{X} is a good compactification of a smooth quasi-affine curve $X \rightarrow S$ (see 7.8), then $(\bar{X}, \bar{X} - X, \emptyset)$ is a standard triple.

We will see in 11.17 below that any pair of smooth quasi-projective varieties $Z \subset X$ is locally part of a standard triple, at least when k is infinite.

REMARK 11.6. (Gabber) Parts (4) and (2) imply that S is affine, and that Z and X_∞ are finite over S . Indeed, X_∞ is finite and surjective over S by part (2), and affine by part (4), so Chevalley's theorem ([Har77, III Ex.4.2]) implies that S is affine.

We will make use of the following observation. Recall from 7.10 that $\text{Pic}(\bar{X}, X_\infty)$ is the group of isomorphism classes of pairs (\mathcal{L}, s) where \mathcal{L} is a line bundle on \bar{X} and s is a trivialization on X_∞ .

Given a standard triple (\bar{X}, X_∞, Z) , any section $x : S \rightarrow X$ of p defines an element $[x]$ of $\text{Pic}(\bar{X}, X_\infty)$. Indeed, there is a homomorphism $C_0(X/S) \rightarrow \text{Pic}(\bar{X}, X_\infty)$.

REMARK 11.7. Let F be a homotopy invariant presheaf with transfers. Given a standard triple (\bar{X}, X_∞, Z) , by 7.5 there is a pairing:

$$(\ , \) : \text{Pic}(\bar{X}, X_\infty) \otimes F(X) \rightarrow F(S).$$

Let $x : S \rightarrow X$ be a section of p . If $[x]$ is the class of x in $\text{Pic}(\bar{X}, X_\infty)$, then $([x], f) = F(x)(f)$ for all $f \in F(X)$.

LEMMA 11.8. *Let (\bar{X}, X_∞, Z) be a standard triple over S and $X = \bar{X} - X_\infty$. Then there is a commutative diagram for every homotopy invariant presheaf with transfers F .*

$$\begin{array}{ccc} \text{Pic}(\bar{X}, X_\infty \amalg Z) \otimes F(X) & \longrightarrow & \text{Pic}(\bar{X}, X_\infty \amalg Z) \otimes F(X - Z) \\ \downarrow & & \downarrow \\ \text{Pic}(\bar{X}, X_\infty) \otimes F(X) & \longrightarrow & F(S) \end{array}$$

PROOF. By definition, \bar{X} is a good compactification of both X and $X - Z$. Thus the pairings exist by 7.5 (or 7.16) and are induced by the transfers pairing $\text{Cor}_k(S, X) \otimes F(X) \rightarrow F(S)$. Commutativity of the diagram is a restatement of the fact that any presheaf with transfers is a functor on Cor_k . \square

COROLLARY 11.9. *If $x : S \rightarrow X$ is a section and $[x] \in \text{Pic}(\bar{X}, X_\infty)$ lifts to $\lambda \in \text{Pic}(\bar{X}, X_\infty \amalg Z)$, there is a commutative diagram:*

$$\begin{array}{ccc} F(X) & \longrightarrow & F(X - Z) \\ \downarrow [x] & \searrow \lambda & \\ F(S) & & \end{array}$$

Moreover, if $\lambda' \in C_0(X - Z/S) \subset \text{Cor}(S, X - Z)$ is any representative of λ (see 7.16 and 1A.12), the composition of λ' with the inclusion $X - Z \subset X$ is \mathbb{A}^1 -homotopic to x in $\text{Cor}(S, X)$.

EXERCISE 11.10. Use example 7.14 with $F = \mathcal{O}^*$ to show that there can be more than one lift $\lambda : F(X - Z) \rightarrow F(S)$.

More generally, observe that any unit s of $\mathcal{O}(Z)$ gives a trivialization of $\mathcal{O}(\bar{X})$ on Z ; combining this with the trivialization 1 on X_∞ gives an element $\sigma(s) = (\mathcal{O}, 1 \amalg s)$ of $\text{Pic}(\bar{X}, X_\infty \amalg Z)$. Show that $\lambda + \sigma(s)$ is also a lift of $[x]$ to $\text{Pic}(\bar{X}, X_\infty \amalg Z)$, and that every other lift has this form for some $s \in \mathcal{O}^*(Z)$.

DEFINITION 11.11. A standard triple is **split** over an open subset $U \subset X$ if $\mathcal{L}_\Delta|_{U \times_S Z}$ is trivial, where \mathcal{L}_Δ is the line bundle on $U \times_S \bar{X}$ corresponding to the graph of the diagonal map.

EXAMPLE 11.12. For any affine S , the standard triple $(S \times \mathbb{P}^1, S \times \infty, S \times 0)$ is split over any U in $X = S \times \mathbb{A}^1$. Indeed, the line bundle \mathcal{L}_Δ is trivial on all of $X \times X$.

EXERCISE 11.13. Let \bar{X} be a smooth projective curve over k , with affine open $X = \text{Spec}(A)$ and set $X_\infty = \bar{X} - X$. Then (\bar{X}, X_∞, Z) is a standard triple for every finite Z in X . Let P_1, \dots be the prime ideals of A defining the points of Z , and suppose for simplicity that $A/P_i \cong k$ for all i . Show that the standard triple splits over $D(f)$ if and only if each P_i becomes a principal ideal in the ring $A[1/f]$.

In particular, if $\bar{X} = \mathbb{P}^1$, the triple splits over all X because in this case A is a principal ideal domain.

LEMMA 11.14. *Any finite set of points in X has an open neighborhood U such that the triple is split over U .*

PROOF. The map $f : X \times_S Z \rightarrow X$ is finite, as Z is finite over S . Given points $x_i \in X$, each $f^{-1}(x_i)$ is finite. Now the line bundle \mathcal{L}_Δ is trivial in some neighborhood V of $\cup_i f^{-1}(x_i)$, because every line bundle on a semilocal scheme is trivial. But every such V contains an open of the form $U \times_S Z$, and the triple is split over such a U . \square

PROPOSITION 11.15. *Consider a standard triple split over an affine U . Then there is an \mathbb{A}^1 -equivalence class of finite correspondences $\lambda : U \rightarrow (X - Z)$ such that the composite of λ with $(X - Z) \subset X$ is \mathbb{A}^1 -homotopic to the inclusion $U \subset X$.*

In particular, $F(X) \rightarrow F(U)$ factors through $\lambda : F(X - Z) \rightarrow F(U)$:

$$\begin{array}{ccc} F(X) & \longrightarrow & F(X - Z) \\ \downarrow & \nearrow \exists \lambda & \\ & & F(U) \end{array}$$

PROOF. Pulling back yields a standard triple $(U \times_S \bar{X}, U \times_S X_\infty, U \times_S Z)$ over the affine U . The diagonal $\Delta : U \rightarrow U \times_S X$ is a section and its class in $\text{Pic}(U \times_S \bar{X}, U \times_S X_\infty)$ is represented by the line bundle \mathcal{L}_Δ . If the triple is split over an affine U , then \mathcal{L}_Δ has a trivialization on $U \times_S Z$ as well, so $[\Delta]$ lifts to a class λ in $\text{Pic}(U \times_S \bar{X}, U \times_S (X_\infty \amalg Z))$. By 7.2 and 7.16, λ is an \mathbb{A}^1 -equivalence class of maps in $\text{Cor}(U, X - Z)$. By 11.9 we have a commutative diagram

$$\begin{array}{ccccc} U & \xrightarrow{\lambda} & U \times_S (X - Z) & \xrightarrow{[pr]} & X - Z \\ & \searrow [\Delta] & \downarrow & & \downarrow \\ & & U \times_S X & \xrightarrow{[pr]} & X \end{array}$$

and it suffices to observe that $pr \circ \Delta : U \rightarrow U \times_S X \rightarrow X$ is the inclusion. \square

A different splitting (trivialization on $U \times_S Z$) may yield a different lifting λ' . By exercise 11.10, $\lambda' = \lambda + \sigma(s)$ for some unit s of $\mathcal{O}(U \times_S Z)$.

EXERCISE 11.16. Suppose that λ is represented by an element D of $\text{Cor}(U, X - Z) = C_0(U \times (X - Z)/U)$, as in exercise 7.15. Show that the element $D - [\Delta(U)]$ of $\text{Cor}(U, X)$ is represented by a principal divisor (f) on $U \times \bar{X}$, with f equal to 1 on $U \times X_\infty$.

THEOREM 11.17. *Let W be a connected quasi-projective smooth scheme over an infinite field k , Y a proper closed subset of W and $y_1, \dots, y_n \in Y$. Then there is*

an affine open neighborhood X of these points in W and a standard triple $(\bar{X} \rightarrow S, X_\infty, Z)$ such that $(X, X \cap Y) \cong (\bar{X} - X_\infty, Z)$.

PROOF. (Mark Walker) We may assume that W is affine, a closed $(d+1)$ -dimensional subscheme of \mathbb{A}^n . Embed \mathbb{A}^n in \mathbb{A}^N by

$$(x_1, \dots, x_n) \mapsto (x_1, \dots, x_n, x_1^2, x_1x_2, \dots, x_ix_j, \dots, x_n^2).$$

Given a closed point $x \in W$, Bertini's Theorem (see [SGA4, XI.2.1]) implies that the general linear projection $p : \mathbb{A}^N \rightarrow \mathbb{A}^d$ is smooth near each point of W lying on $p^{-1}(p(x))$. It is also finite when restricted to Y , because Y has dimension $\leq d$.

Let \bar{W} denote the closure of W in \mathbb{P}^N , $H = \mathbb{P}^N - \mathbb{A}^N$, and $W_\infty = \bar{W} \cap H$. The general projection defines a rational map $p : \bar{W} \dashrightarrow \mathbb{P}^d$ whose center C is finite, because C lies in the intersection of W_∞ with a codimension d linear subspace of H . Let \bar{X}_1 be the closure of the graph of $p : (\bar{W} - C) \rightarrow \mathbb{P}^d$ in $\bar{W} \times \mathbb{P}^d$. Then W is naturally an open subscheme of \bar{X}_1 and $\bar{X}_1 - W$ has finite fibers over \mathbb{A}^d .

The singular points Σ of the projection $\bar{X}_1 \rightarrow \mathbb{P}^d$ are closed, and finite over each $p(y_i)$ because p is smooth near $W \cap p^{-1}(p(y_i))$. Therefore there is an affine open neighborhood S in \mathbb{A}^d of $\{p(y_i)\}$ over which Σ is finite and disjoint from Y . Define X to be $p^{-1}(S) \cap W - \Sigma$; by construction $p : X \rightarrow S$ is smooth. Define $\bar{X} \subset \bar{X}_1$ to be the inverse image of S , and $X_\infty = \bar{X} - X$. Then $X \cap Y \rightarrow S$ and $X_\infty \rightarrow S$ are both finite.

It remains to show that $X_\infty \amalg (X \cap Y)$ lies in an affine open neighborhood of \bar{X} . As \bar{X} is projective over S , there is a global section of some very ample line bundle \mathcal{L} whose divisor D misses all of the finitely many points of X_∞ and $X \cap Y$ over any $p(y_i)$. Because \mathcal{L} is very ample and S is affine, $\bar{X} - D$ is affine. Replacing S by a smaller affine neighborhood of the $p(y_i)$, we can assume that D misses X_∞ and $X \cap Y$, i.e., that X_∞ and $X \cap Y$ lie in $\bar{X} - D$, as desired. \square

PORISM 11.18. If k is finite, the proof shows that there is a finite extension k' and an affine open X' of the points in $W \times_k \text{Spec } k'$ so that $(X', X' \cap Y')$ comes from a standard triple over k' , where $Y' = Y \times_k \text{Spec } k'$. In fact, for each prime l we can assume that $[k' : k]$ is a power of l .

Finally, we will use 11.15, 11.14 and 11.17 to prove 11.3.

PROOF OF 11.3. We first assume that k is infinite. Since we may replace V by $V - \{x_1, \dots, x_n\}$, we may assume that the closed points x_1, \dots, x_n of X lie in $Z = X - V$. We can use 11.17 to shrink X about these points to assume that there exists a standard triple with $X = \bar{X} - X_\infty$. By 11.14 the triple splits over an open neighborhood U of the points. As X is quasi-projective, we may shrink U to make it affine. By 11.15 we get the map $F(X - Z) \rightarrow F(U)$ factoring $F(X) \rightarrow F(U)$.

If k is finite, we proceed as follows. We see by porism 11.18 that there is an open X' of $X \times_k \text{Spec}(k')$ fitting into a standard triple over k' . The argument above shows that there is an open neighborhood U of x_1, \dots, x_n (depending on k') such that if $U' = U \times_k \text{Spec}(k')$ and $V' = V \times_k \text{Spec}(k')$, then $F(X') \rightarrow F(U')$ factors through a map $\Phi' : F(V') \rightarrow F(U')$. Let $\Phi(k') : F(V) \rightarrow F(U)$ be the composite of

Φ' and the transfer $F(U') \rightarrow F(U)$. By 1.11, $[k' : k]$ times $F(X) \rightarrow F(U)$ factors through $\Phi(k')$. By 11.18, we can choose two such extensions k', k'' with $[k' : k]$ and $[k'' : k]$ relatively prime. Shrinking U , we may assume that $F(U)$ is the target of both $\Phi(k')$ and $\Phi(k'')$. But then $F(X) \rightarrow F(U)$ factors through a linear combination of $\Phi(k')$ and $\Phi(k'')$. \square

LECTURE 12

Nisnevich sheaves

We have already mentioned the Nisnevich topology several times in previous lectures, as an alternative to the étale and Zariski topologies. In this lecture we develop some of its more elementary properties.

We begin by recalling the definition of the Nisnevich topology (see [Nis89]). A family of étale morphisms $\{p_i : U_i \rightarrow X\}$ is said to be a **Nisnevich covering** of X if it has the Nisnevich lifting property:

- for all $x \in X$, there is an i and a $u \in U_i$ so that $p_i(u) = x$ and the induced map $k(x) \rightarrow k(u)$ is an isomorphism.

It is easy to check that this notion of cover satisfies the axioms for a Grothendieck topology (in the sense of [Mil80, I.1.1], or pre-topology in the sense of [SGA4]). The Nisnevich topology is the class of all Nisnevich coverings.

EXAMPLE 12.1. Here is an example to illustrate the arithmetic nature of a Nisnevich cover. When $\text{char } k \neq 2$, the two morphisms $U_0 = \mathbb{A}^1 - \{a\} \xrightarrow{c \rightarrow j} \mathbb{A}^1$ and $U_1 = \mathbb{A}^1 - \{0\} \xrightarrow{z \rightarrow z^2} \mathbb{A}^1$ form a Nisnevich covering of \mathbb{A}^1 if and only if $a \in (k^*)^2$. They form an étale covering of \mathbb{A}^1 for any nonzero $a \in k$.

EXAMPLE 12.2. Let k be a field. The small Nisnevich site on $\text{Spec } k$ consists of the étale U over $\text{Spec } k$, together with their Nisnevich coverings. Every étale U over $\text{Spec } k$ is a finite disjoint union $\coprod \text{Spec } l_i$ with the l_i finite and separable over k ; to be a Nisnevich cover, one of the l_i must equal k . Thus a Nisnevich sheaf F on $\text{Spec } k$ merely consists of a family of sets $F(l)$, natural in the finite separable extension fields l of k . In fact, each such l determines a “point” of $(\text{Spec } k)_{\text{Nis}}$ in the sense of [SGA4, IV 6.1].

From this description it follows that $\text{Spec } k$ has Nisnevich cohomological dimension zero. This implies that the Nisnevich cohomological dimension of any Noetherian scheme X is at most $\dim X$; see [KS86].

LEMMA 12.3. *If $\{U_i \rightarrow X\}$ is a Nisnevich covering then there is a nonempty open $V \subset X$ and an index i such that $U_i|_V \rightarrow V$ has a section.*

PROOF. For each generic point x of X , there is a generic point $u \in U_i$ so that $k(x) \cong k(u)$. Hence $U_i \rightarrow X$ induces a rational isomorphism between the corresponding components of U_i and X , i.e., $U_i \rightarrow X$ has a section over an open subscheme V of X containing x . □

EXAMPLE 12.4. A **Hensel local ring or scheme** (R, \mathfrak{m}) is a local ring such that any finite R -algebra S is a product of local rings. It is well-known (see [Mil80, I.4.2]) that if S is finite and étale over R , and if $R/\mathfrak{m} \cong S/\mathfrak{m}_i$ for some maximal ideal \mathfrak{m}_i of S , then $R \rightarrow S$ splits; one of the factors of S is isomorphic to R . If $\{U_i \rightarrow \text{Spec } R\}$ is a Nisnevich covering then some U_i is finite étale, so $U_i \rightarrow \text{Spec } R$ splits. Thus every Nisnevich covering of $\text{Spec } R$ has the trivial covering as a refinement. Consequently, the Hensel local schemes $\text{Spec } R$ determine “points” for the Nisnevich topology.

As with any Grothendieck topology, the category $Sh_{Nis}(Sm/k)$ of Nisnevich sheaves of abelian groups is abelian, and sheafification $F \mapsto F_{Nis}$ is an exact functor. We know that exactness in $Sh_{Nis}(Sm/k)$ may be tested at the Hensel local rings $\mathcal{O}_{X,x}^h$ of all smooth X at all points x (see [Nis89, 1.17]). That is, for every presheaf F :

- $F_{Nis} = 0$ if and only if $F(\text{Spec } \mathcal{O}_{X,x}^h) = 0$ for all (X, x) ;
- $F_{Nis}(\text{Spec } \mathcal{O}_{X,x}^h) = F(\text{Spec } \mathcal{O}_{X,x}^h)$.

By abuse of notation, we shall write $F(\mathcal{O}_{X,x}^h)$ for $F(\text{Spec } \mathcal{O}_{X,x}^h)$, and refer to it as the *stalk* of F_{Nis} at x .

DEFINITION 12.5. A commutative square $Q = Q(X, Y, A)$ of the form

$$\begin{array}{ccc} B & \xrightarrow{i} & Y \\ \downarrow f & & \downarrow f \\ A & \xrightarrow{i} & X \end{array}$$

is called **upper distinguished** if $B = A \times_X Y$, f is étale, $i: A \rightarrow X$ is an open embedding and $(Y - B) \rightarrow (X - A)$ is an isomorphism. Clearly, any upper distinguished square determines a Nisnevich covering of X : $\{Y \rightarrow X, A \rightarrow X\}$.

EXERCISE 12.6. If $\dim X \leq 1$ show that any Nisnevich cover of X admits a refinement $\{U, V\}$ such that $Q(X, U, V)$ is upper distinguished. Show that this fails if $\dim X \geq 2$. *Hint:* Cover \mathbb{P}^n by copies of \mathbb{A}^n .

By definition, $F(Q)$ is a pullback square if and only if $F(X)$ is the pullback $F(Y) \times_{F(B)} F(A)$, i.e., the kernel of $f - i: F(Y) \times F(A) \rightarrow F(B)$.

LEMMA 12.7. A presheaf F is a Nisnevich sheaf if and only if $F(Q)$ is a pullback square for every upper distinguished square Q .

PROOF. For the “if” part, suppose that each $F(Q)$ is a pullback square. To prove that F is a Nisnevich sheaf, fix a Nisnevich covering $\{U_i \rightarrow X\}$. Let us say that an open subset $V \subset X$ is *good* (for the covering) if

$$F(V) \longrightarrow \prod F(U_i \times_X V) \rightrightarrows \prod F(U_i \times_X U_j \times_X V)$$

is an equalizer diagram. We need to show that X itself is *good*.

By Noetherian induction, we may assume that there is a largest *good* $V \subset X$. Suppose that $V \neq X$ and let $Z = X - V$. By lemma 12.3, there is a nonempty open $W \subset Z$ and an index i such that $U_i|_W \rightarrow W$ splits. Let $X' \subset X$ be the complement of the closed set $Z - W$. Then V and $U'_i = U_i|_{X'}$ form an upper distinguished square Q over X' . Pulling back along each $U'_j = U_j|_{X'}$ also yields an upper distinguished square. Thus we have pullback squares

$$\begin{array}{ccc}
 F(X') & \longrightarrow & F(U'_i) & & F(U'_j) & \longrightarrow & F(U'_i \times_X U'_j) \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 F(V) & \longrightarrow & F(U'_i|_V) & & F(U'_j|_V) & \longrightarrow & F(U'_i \times_X U'_j|_V).
 \end{array}$$

A diagram chase shows that X' is also *good*, contradicting the assumption that $V \neq X$. Hence X is *good* for each cover, i.e., F is a Nisnevich sheaf.

For “only if”, we assume that F is a Nisnevich sheaf and Q is upper distinguished and we need to prove that the map $F(X) \rightarrow F(Y) \times_{F(B)} F(A)$ is an isomorphism. We already know the map is monic because $\{A, Y\}$ is a Nisnevich cover of X . For the surjectivity, note that the sheaf axiom for this covering yields the equalizer sequence

$$F(X) \rightarrow F(Y) \times F(A) \rightrightarrows F(B) \times F(A) \times F(Y \times_X Y).$$

Since $\{\Delta(Y), B \times_A B\}$ is a cover of $Y \times_X Y$, we have an injection $F(Y \times_X Y) \rightarrow F(Y) \times F(B \times_A B)$. Now $(a, y) \in F(A) \times F(Y)$ lies in $F(A) \times_{F(B)} F(Y)$ if the two restrictions to $F(B)$ are the same. The two maps to $F(A)$ and $F(Y)$ are the same, so it suffices to consider the maps from $F(Y)$ to $F(B \times_A B)$. These both factor through $F(B)$, so the images of y are the same as the images of a . By construction the two maps $F(A) \rightrightarrows F(B \times_A B)$ are the same. \square

PORISM 12.8. Suppose more generally that F is a sheaf for some Grothendieck topology, and that $Q = Q(X, Y, A)$ is a pullback square whose horizontal maps are monomorphisms. If $\{A, Y\}$ is a cover of X and $\{B \times_A B, Y\}$ is a cover of $Y \times_X Y$, the proof of lemma 12.7 shows that $F(Q)$ is a pullback square.

EXERCISE 12.9. Write $\mathcal{O}^*/\mathcal{O}^{*l}$ for the presheaf $U \mapsto \mathcal{O}^*(U)/\mathcal{O}^{*l}(U)$, and \mathcal{O}^*/l for the Zariski sheaf associated to $\mathcal{O}^*/\mathcal{O}^{*l}$. Show that there is an exact sequence

$$0 \rightarrow \mathcal{O}^*(U)/\mathcal{O}^{*l}(U) \rightarrow \mathcal{O}^*/l(U) \rightarrow \text{Pic}(U) \xrightarrow{l} \text{Pic}(U)$$

for all smooth U . Then show that \mathcal{O}^*/l is a Nisnevich sheaf on Sm/k . If $1/l \in k$, this is an example of a Nisnevich sheaf which is not an étale sheaf. In fact, $(\mathcal{O}^*/l)_{\acute{e}t} = 0$.

EXERCISE 12.10. If F is a Nisnevich sheaf, consider the presheaf $E^0(F)$ defined by:

$$E^0(F)(X) = \prod_{\substack{\text{closed} \\ x \in X}} F(\mathcal{O}_{X,x}^h).$$

Show that $E^0(F)$ is a Nisnevich sheaf, and that the canonical map $F \rightarrow E^0(F)$ is an injection. Using 12.2, show that $E^0(F)$ is a flasque sheaf, i.e., that it has no higher cohomology (see [SGA4, V.4.1]). Iteration of this construction yields the canonical flasque resolution $0 \rightarrow F \rightarrow E^0(F) \rightarrow \cdots$ of a Nisnevich sheaf, which may be used to compute the cohomology groups $H_{Nis}^*(X, F)$.

DEFINITION 12.11. Consider the presheaf h_X sending U to $\mathbb{Z}[\text{Hom}_{Sm/k}(U, X)]$. We write $\mathbb{Z}(X)$ for its sheafification $(h_X)_{Nis}$ with respect to the Nisnevich topology. It is easily checked that $\mathbb{Z}(X)(U) = \mathbb{Z}[\text{Hom}(-, X)](U)$ for every connected open U . This is false for non-connected U , since $\mathbb{Z}(X)(U_1 \amalg U_2) = \mathbb{Z}(X)(U_1) \oplus \mathbb{Z}(X)(U_2)$ but $h_X(U_1 \amalg U_2) = h_X(U_1) \otimes h_X(U_2)$.

By the Yoneda lemma, $\text{Hom}(\mathbb{Z}(X), G) = G(X)$ for every sheaf G . Since $\mathbb{Z}_{tr}(X)$ is a Nisnevich sheaf by 6.2, we see that $\mathbb{Z}(X)$ is a subsheaf of $\mathbb{Z}_{tr}(X)$.

Let \mathbf{D}_{Nis}^- denote the derived category of cohomologically bounded above complexes in $Sh_{Nis}(Sm/k)$. If F and G are Nisnevich sheaves, it is well known that $\text{Ext}_{Nis}^n(F, G) = \text{Hom}_{\mathbf{D}_{Nis}^-}(F, G[n])$ (see [Wei94, 10.7.5]).

LEMMA 12.12. *Let G be a complex of Nisnevich sheaves. Then for all X :*

$$\text{Ext}_{Nis}^n(\mathbb{Z}(X), G) = \mathbb{H}_{Nis}^n(X, G).$$

PROOF. First suppose that G is a sheaf. If $G \rightarrow I^*$ is a resolution by injective Nisnevich sheaves, then the n^{th} cohomology of G is H^n of $I^*(X)$. But by [Wei94, 10.7.4] we know that the left side is H^n of $\text{Hom}_{Sh_{Nis}(Sm/k)}(\mathbb{Z}(X), I^*) = I^*(X)$. A similar argument applies when G is a complex. \square

LEMMA 12.13. *The smallest class in \mathbf{D}_{Nis}^- which contains all the $\mathbb{Z}(X)$ and is closed under quasi-isomorphisms, direct sums, shifts, and cones is all of \mathbf{D}_{Nis}^- .*

PROOF. The proof of 9.4 goes through using $\mathbb{Z}(X)$ in place of $R_{tr}(X)$. \square

For the rest of this lecture, we shall write \otimes for the presheaf tensor product, $(F \otimes G)(U) = F(U) \otimes_{\mathbb{Z}} G(U)$, and \otimes_{Nis} for the tensor product of Nisnevich sheaves, i.e., the sheafification of \otimes . Note that if a sheaf F is flat as a presheaf then F is also flat as a sheaf. This is true for example of the sheaves $\mathbb{Z}(X)$.

LEMMA 12.14. $\mathbb{Z}(X \times Y) = \mathbb{Z}(X) \otimes_{Nis} \mathbb{Z}(Y)$.

PROOF. Since $\text{Hom}(U, X \times Y) = \text{Hom}(U, X) \times \text{Hom}(U, Y)$, we see that $\mathbb{Z}[\text{Hom}(U, X \times Y)] = \mathbb{Z}[\text{Hom}(U, X)] \otimes \mathbb{Z}[\text{Hom}(U, Y)]$. Thus $\mathbb{Z}[\text{Hom}(-, X \times Y)] \cong \mathbb{Z}[\text{Hom}(-, X)] \otimes \mathbb{Z}[\text{Hom}(-, Y)]$ as presheaves. Now sheafify. \square

LEMMA 12.15. *Let G be a Nisnevich sheaf on Sm/k such that $H_{Nis}^n(-, G)$ is homotopy invariant for all n . Then for all n and all bounded above C :*

$$\mathrm{Hom}_{\mathbf{D}_{Nis}^-}(C, G[n]) \cong \mathrm{Hom}_{\mathbf{D}_{Nis}^-}(C \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1), G[n]).$$

PROOF. By 12.12, our assumption yields $\mathrm{Ext}^n(\mathbb{Z}(X), G) \cong \mathrm{Ext}^n(\mathbb{Z}(X \times \mathbb{A}^1), G)$ for all X . Since $\mathbb{Z}(X \times \mathbb{A}^1) = \mathbb{Z}(X) \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1)$ by 12.14, the conclusion holds for $C = \mathbb{Z}(X)$. If C and C' are quasi-isomorphic, then so are $C \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1)$ and $C' \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1)$, because $\mathbb{Z}(\mathbb{A}^1)$ is a flat sheaf. But the class of all complexes C for which the conclusion holds is closed under direct sums and cones, so by 12.13 the conclusion holds for all C . \square

We borrow yet another topological definition: deformation retract. For each F , note that the presheaf $F \otimes \mathbb{Z}[\mathrm{Hom}(-, \mathrm{Spec} k)]$ is just F .

DEFINITION 12.16. An injection of presheaves $i : F \rightarrow G$ is called a **(strong) deformation retract** if there is a map $r : G \rightarrow F$ such that $r \circ i = id_F$ and a homotopy $h : G \otimes \mathbb{Z}[\mathrm{Hom}(-, \mathbb{A}^1)] \rightarrow G$ so that the restriction $h|_F$ is the projection $F \otimes \mathbb{Z}[\mathrm{Hom}(-, \mathbb{A}^1)] \rightarrow F$, $h(G \otimes 0) = i \circ r$ and $h(G \otimes 1) = id$.

If F and G are sheaves, the condition in the definition is equivalent to the condition that there is a sheaf map $h : G \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1) \rightarrow G$ so that the restriction $h|_F$ is the projection $F \otimes_{Nis} \mathbb{Z}(\mathbb{A}^1) \rightarrow F$, $h(G \otimes 0) = i \circ r$ and $h(G \otimes 1) = id$.

For example, the zero-section $\mathrm{Spec} k \xrightarrow{0} \mathbb{A}^1$ induces a deformation retract $\mathbb{Z} \rightarrow \mathbb{Z}(\mathbb{A}^1)$; the homotopy map h is induced by the multiplication $\mathbb{A}^1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^1$ using 12.14. If I^1 is the quotient presheaf $\mathbb{Z}(\mathbb{A}^1)/\mathbb{Z}$, so that $\mathbb{Z}(\mathbb{A}^1) \cong \mathbb{Z} \oplus I^1$, then $0 \subset I^1$ is also a deformation retract.

LEMMA 12.17. *If $F \rightarrow G$ is a deformation retract, then the quotient presheaf G/F is a direct summand of $G/F \otimes I^1$.*

PROOF. The inclusion $0 \subset G/F$ is a deformation retract, whose homotopy is induced from h . Therefore we may assume that $F = 0$.

Let K denote the kernel of h . Since the evaluation “ $t = 1$ ” : $G = G \otimes \mathbb{Z} \rightarrow G \otimes \mathbb{Z}(\mathbb{A}^1)$ is a section of both h and the projection $G \otimes \mathbb{Z}(\mathbb{A}^1) \rightarrow G$, we see that K is isomorphic to $G \otimes I^1$. But “ $t = 0$ ” : $G \rightarrow G \otimes \mathbb{Z}(\mathbb{A}^1)$ embeds G as a summand of K . \square

For every presheaf F we define $\tilde{C}_m(F)$ to be the quotient presheaf $C_m(F)/F$. That is, $\tilde{C}_m(F)(U)$ is $F(U \times \mathbb{A}^m)/F(U)$. Thus we have split exact sequences $0 \rightarrow F \rightarrow C_m(F) \rightarrow \tilde{C}_m(F) \rightarrow 0$.

COROLLARY 12.18. *$\tilde{C}_m(F)$ is a direct summand of $\tilde{C}_m(F) \otimes I^1$ for all $m \geq 0$.*

PROOF. It is easy to see that $F \rightarrow C_m F$ is a deformation retract, so 12.18 is a special case of 12.17. \square

PROPOSITION 12.19. *Let G be a Nisnevich sheaf on Sm/k such that $H_{Nis}^n(-, G)$ is homotopy invariant for all n . Then for all n and for all presheaves F , there is an isomorphism*

$$\mathrm{Hom}_{\mathbf{D}_{Nis}^-}((C_*F)_{Nis}, G[n]) \xrightarrow{\cong} \mathrm{Hom}_{\mathbf{D}_{Nis}^-}(F_{Nis}, G[n]).$$

PROOF. Write $\mathrm{Ext}^n(C, G)$ for $\mathrm{Hom}_{\mathbf{D}_{Nis}^-}(C, G[n])$. For each complex C , lemma 12.15 implies that $\mathrm{Ext}^q(C \otimes_{Nis} I^1, G) = 0$ for all q . For $C = (\tilde{C}_p F)_{Nis}$, 12.18 yields $\mathrm{Ext}^q((\tilde{C}_p F)_{Nis}, G) = 0$.

Note that $\mathrm{Ext}^q(C, G) = H^q \mathbb{R} \mathrm{Hom}(C, G)$ for any C ; see [Wei94, 10.7.4]. As in the proof of 10.10, a resolution $G \rightarrow I^*$ yields a first quadrant Hom double complex $\mathrm{Hom}((\tilde{C}_* F)_{Nis}, I^*)$ and hence a first quadrant spectral sequence

$$E_1^{p,q} = \mathrm{Ext}^q((\tilde{C}_p F)_{Nis}, G) \Rightarrow \mathrm{Ext}^{p+q}((\tilde{C}_* F)_{Nis}, G)$$

(see [Wei94, 5.6.1]). Since every $E_1^{p,q}$ vanishes, this implies that $\mathrm{Ext}^n((\tilde{C}_* F)_{Nis}, G) = 0$ for all n . In turn, this implies the conclusion of 12.19, viz., $\mathrm{Ext}^n((C_* F)_{Nis}, G) \cong \mathrm{Ext}^n(F, G)$ for all n . \square

EXERCISE 12.20. If U is open in X , $\mathbb{Z}(U)$ is a subsheaf of $\mathbb{Z}(X)$; write $\mathbb{Z}(X, U)$ for the quotient Nisnevich sheaf. If $f : Y \rightarrow X$ is an étale morphism of smooth schemes over k , and $Z \subset X$ is a closed subscheme isomorphic to $f^{-1}(Z)$, show that $\mathbb{Z}(Y, Y - f^{-1}(Z)) \xrightarrow{\cong} \mathbb{Z}(X, X - Z)$.

The cdh topology

In order to extend the main results of the following chapters to (possibly) singular schemes, we need to introduce another topology: the cdh topology on the category Sch/k of schemes of finite type over k . A crucial part will be played by the following notion.

DEFINITION 12.21. Let X be a scheme of finite type over a field k and let $i : Z \rightarrow X$ be a closed immersion. Then an **abstract blow-up** of X with center Z is a proper map $p : X' \rightarrow X$ which induces an isomorphism $(X' - Z')_{red} \cong (X - Z)_{red}$, where $Z' = X' \times_X Z$. We will often refer to the cartesian square

$$\begin{array}{ccc} Z' & \longrightarrow & X' \\ \downarrow & & \downarrow p \\ Z & \xrightarrow{i} & X. \end{array}$$

We will say that $p : X' \rightarrow X$ is an abstract blow-up if there exists a $Z \subseteq X$ which satisfies the conditions above.

EXERCISE 12.22. Let $X' \rightarrow X$ be an abstract blow-up with center Z . Show that both $0 \rightarrow \mathbb{Z}(Z') \rightarrow \mathbb{Z}(X') \oplus \mathbb{Z}(Z) \rightarrow \mathbb{Z}(X)$ and $0 \rightarrow \mathbb{Z}_{tr}(Z') \rightarrow \mathbb{Z}_{tr}(X') \oplus \mathbb{Z}_{tr}(Z) \rightarrow \mathbb{Z}_{tr}(X)$ are exact sequences of Nisnevich sheaves on Sm/k .

DEFINITION 12.23. The cdh topology on Sch/k is the minimal Grothendieck topology generated by Nisnevich covers and covers $X' \amalg Z \rightarrow X$ corresponding to abstract blow-ups. A **proper cdh cover** is a proper map which is also a cdh cover. A proper cdh cover of a reduced scheme is called a **proper birational cover** if it is an isomorphism over a dense open subscheme.

If F is any presheaf on Sch/k we will write F_{cdh} for its sheafification with respect to this topology.

The name “cdh” stands for “completely decomposed h-topology”; “completely decomposed” is the original term for the Nisnevich topology (see [Nis89]), and the h-topology was introduced in Voevodsky’s thesis.

EXAMPLE 12.24. It is easy to see from the definition that $X_{red} \rightarrow X$ is a proper cdh cover, and that every cdh cover has the Nisnevich lifting property (see p. 89). In particular, it follows as in 12.2 that every cover of $Spec k$ has a section, and that any 0-dimensional scheme has cdh cohomological dimension zero. In fact, the cdh cohomological dimension of any Noetherian scheme is at most $\dim X$; see [SV00, 5.13].

If $X' \rightarrow X$ is an abstract blow-up with center Z , and Z contains no generic point of X , then $X' \amalg Z \rightarrow X$ is a proper birational cdh cover.

EXAMPLE 12.25. If X is reduced, every proper cdh cover $\tilde{X} \rightarrow X$ has a refinement which is a proper birational cdh cover. To see this, note that the Nisnevich lifting property applied to the generic point of X yields a closed subscheme X' of \tilde{X} such that $X' \rightarrow X$ is a birational isomorphism, i.e., an isomorphism over a dense open of the form $X - Z$. But then $X' \amalg Z \rightarrow X$ is a cdh cover, and $X' \amalg (\tilde{X} \times_X Z) \rightarrow X$ is a proper birational cdh cover.

To better understand the structure of the cdh topology, we need to study some properties of its coverings.

LEMMA 12.26. (See [SV00, 5.8].) *A proper map is a proper cdh cover if and only if it satisfies the Nisnevich lifting property.*

PROOF. Let $\tilde{X} \rightarrow X$ be a proper map satisfying the Nisnevich lifting property; we must show that it is a cdh cover. By 12.24, this is true if $\dim X = 0$, and we may assume that X is reduced and irreducible. We will proceed by induction on $\dim X$.

Consider the proper birational cdh cover $X' \amalg Z \rightarrow X$ constructed in 12.25. The pullback of $\tilde{X} \rightarrow X$ along this cover consists of $\tilde{X} \times_X X' \rightarrow X'$ (which is a cover because it has a section) and $\tilde{X} \times_X Z \rightarrow Z$ (which is a cdh cover by induction on $\dim X$ because the Nisnevich lifting property is satisfied). Since $\tilde{X} \rightarrow X$ is a cdh cover locally in the cdh topology, it is a cdh cover. \square

For example, if X is smooth then any blow-up $X' \rightarrow X$ along a smooth center Z is a proper birational cdh cover. Indeed, the inverse image of Z is a projective bundle over Z , and such a bundle always satisfies the Nisnevich lifting property.

PROPOSITION 12.27. Any cdh cover of the form $T \xrightarrow{p} U \xrightarrow{q} X$, where X is integral, p is a proper cdh cover, and q is a Nisnevich cover, has a refinement of the form

$$\begin{array}{ccc} V & \xrightarrow{f} & S \\ \downarrow & & \downarrow g \\ T & \xrightarrow{p} U \xrightarrow{q} & X \end{array}$$

where f is a Nisnevich cover and g is a proper cdh cover.

PROOF. Let $U = \coprod_i U_i$ be the decomposition of U into its irreducible components and let $T_i = T \times_U U_i$. Refining $T \rightarrow U$ as in 12.25, we may assume that each $T_i \rightarrow U_i$ is a proper birational cdh cover. By platification (see [RG71] or 1A.1) applied to $T \rightarrow X$, there is a blow-up $X' \rightarrow X$ along a $Z \subseteq X$ such that the proper transform T'_i of each T_i is flat over X' . We set $U'_i = U_i \times_X X'$. The situation is described by the following diagram in which all squares are cartesian:

$$\begin{array}{ccccccc} T'_i & \longrightarrow & T_i \times_X X' & \longrightarrow & U'_i & \xrightarrow{h} & X' \\ & & \downarrow & & \downarrow & & \downarrow \\ & & T_i & \longrightarrow & U_i & \longrightarrow & X. \end{array}$$

Now $T'_i \rightarrow X'$ is flat, and h is étale by base change, so $g : T'_i \rightarrow U'_i$ is flat. But g is also proper and birational, and therefore g is an isomorphism since both T'_i and U'_i are irreducible. Hence the pullback of $T \rightarrow X$ to X' admits the refinement $U' = \coprod U'_i \cong \coprod T'_i \rightarrow X'$.

By induction on $\dim X$, the induced cover $T \times_X Z \rightarrow U \times_X Z \rightarrow Z$ admits a refinement $V' \rightarrow S' \rightarrow Z$ with $V' \rightarrow S'$ a Nisnevich cover and $S' \rightarrow Z$ a proper cdh cover. But then the required refinement of $T \rightarrow X$ is the composition

$$V = V' \amalg U' \xrightarrow{f} S = S' \amalg X' \rightarrow Z \amalg X' \rightarrow X. \quad \square$$

PROPOSITION 12.28. Every cdh cover of X in Sch/k has a refinement of the form $U \xrightarrow{q} X' \xrightarrow{p} X$, where p is a proper cdh cover and q is a Nisnevich cover.

PROOF. Since $X_{\text{red}} \rightarrow X$ is a proper cdh cover, we may assume X itself is reduced. It will suffice to prove the statement for the irreducible components, and therefore we may assume that X itself is an integral scheme.

By definition, each cdh cover of X can be refined to a cover of the form $X_n \rightarrow X_{n-1} \rightarrow X_{n-2} \rightarrow \cdots \rightarrow X_1 \rightarrow X$, where each map is either a Nisnevich cover or a proper cdh cover. Using 12.27 we can move all Nisnevich covers to the left and all proper cdh covers to the right, which is the statement. \square

PROPOSITION 12.29. Let F be a Nisnevich sheaf on Sch/k . Then $F_{\text{cdh}} = 0$ if and only if for any scheme X and any $a \in F(X)$, there is a proper cdh cover $p : X' \rightarrow X$ such that $p^*(a) = 0 \in F(X')$.

PROOF. If $F_{cdh} = 0$ and $a \in F(X)$, there is a cdh cover $U \rightarrow X$ such that $a|_U = 0$. But by 12.28 we may assume that the cover is of the form $U \xrightarrow{q} X' \xrightarrow{p} X$, where q is a Nisnevich cover and p is a proper cdh cover. We know that $0 = (p \circ q)^*(a) = q^*(p^*(a))$. Since q is a Nisnevich cover and F is a Nisnevich sheaf, q^* is injective, and therefore $p^*(a) = 0$ in $F(X')$.

Now let us assume that the condition holds and consider $a \in F_{cdh}(X)$. Replacing X by a cover, we may assume that $a \in F(X)$. By assumption there is a proper cdh cover $p : X' \rightarrow X$ such that $p^*(a) = 0$. But then $a = 0$ since $F_{cdh}(X)$ injects into $F_{cdh}(X')$. \square

Consider the composite morphism of sites $r : (Sch/k)_{cdh} \rightarrow (Sch/k)_{Nis} \rightarrow (Sm/k)_{Nis}$. If F is a Nisnevich sheaf on Sm/k , the inverse image sheaf $r^*(F)$ is a cdh sheaf on Sch/k . By abuse of notation, we will write F_{cdh} for $r^*(F)$.

If we have resolution of singularities, then not only does every X in Sch/k have an abstract blow-up $X' \rightarrow X$ with X' smooth, but every proper birational cdh cover $X' \rightarrow X$ has a refinement $X'' \rightarrow X$ with X'' smooth, obtained as a composite of blow-ups along smooth centers. Thus every cdh sheaf of Sch/k is determined by its restriction to Sm/k . In fact, assuming resolution of singularities, the functor $F \mapsto F_{cdh}$ from $Sh_{Nis}(Sm/k)$ to $Sh_{cdh}(Sch/k)$ is an exact functor by [SV00, 5.11].

LEMMA 12.30. *Assume that k admits resolution of singularities. Let F be a Nisnevich sheaf on Sm/k . Then $F_{cdh} = 0$ if and only if for any smooth scheme X and any $a \in F(X)$, there is a composition of blow-ups along smooth centers $p : X_r \rightarrow X_{r-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X$ such that $p^*(a) = 0 \in F(X')$.*

PROOF. Assume $F_{cdh} = 0$ and let $a \in F(X)$ for a smooth X . By 12.29 and 12.25, there is a proper birational cdh cover $p : X' \rightarrow X$ such that $p^*(a) = 0$. Refining the cover, we may assume p is a composition of blow-ups along smooth centers.

Conversely, assume that the condition holds. Let $X \in Sch/k$ and let $a \in F_{cdh}(X)$. Passing to a covering, we may assume that $a \in F(X)$ and that X is smooth. By assumption, there is a proper cdh cover $p : X_r \rightarrow X_{r-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X$ such that $p^*(a) = 0 \in F(X')$. \square

EXERCISE 12.31. If C is a nodal curve, show that $H_{cdh}^1(C, \mathbb{Z}) = \mathbb{Z}$.

EXERCISE 12.32. This exercise shows that $H_{Nis}^2(X, \mathbb{Z}) \neq H_{cdh}^2(X, \mathbb{Z})$ for some normal surfaces.

- (1) If X is normal, use [SGA1, I.10.1] to show that $H_{Nis}^n(X, \mathbb{Z}) = 0$ for $n > 0$.
- (2) Let X be a normal surface with a point singularity, whose exceptional fiber is a node. Show that $H_{cdh}^2(X, \mathbb{Z}) \neq 0$.

Nisnevich sheaves with transfers

We now consider the category $Sh_{Nis}(Cor_k)$ of Nisnevich sheaves with transfers. As with étale sheaves, we say that a presheaf with transfers F is a **Nisnevich sheaf with transfers** if its underlying presheaf is a Nisnevich sheaf on Sm/k . Clearly, every étale sheaf with transfers is a Nisnevich sheaf with transfers.

THEOREM 13.1. *Let F be a presheaf with transfers, and write F_{Nis} for the sheafification of the underlying presheaf. Then F_{Nis} has a unique structure of presheaf with transfers such that $F \rightarrow F_{Nis}$ is a morphism of presheaves with transfers.*

Consequently, $Sh_{Nis}(Cor_k)$ is an abelian category, and the forgetful functor $Sh_{Nis}(Cor_k) \rightarrow \mathbf{PST}(k)$ has a left adjoint ($F \mapsto F_{Nis}$) which is exact and commutes with the forgetful functor to (pre)sheaves on Sm/k .

Finally, $Sh_{Nis}(Cor_k)$ has enough injectives.

PROOF. The Nisnevich analogue of 6.16, is valid; just replace ‘étale cover’ by ‘Nisnevich cover’ in the proof. As explained after 6.12, the Čech complex $\mathbb{Z}_{tr}(\check{U})$ is a Nisnevich resolution of $\mathbb{Z}_{tr}(X)$. With these two observations, the proofs of 6.17, 6.18, and 6.19 go through for the Nisnevich topology. \square

EXAMPLE 13.2. By theorem 4.1, $\mathbb{Z}(1) \simeq \mathcal{O}^*[-1]$ as complexes of Nisnevich sheaves with transfers. By 12.9, $\mathcal{O}^*/l = \mathcal{O}^* \otimes_{Nis} \mathbb{Z}/l$. Since $\mathbb{Z}/l(1) = \mathbb{Z}(1) \otimes_{Nis}^L \mathbb{Z}/l$, it follows that there is a distinguished triangle of Nisnevich sheaves with transfers for each l :

$$\mu_l \rightarrow \mathbb{Z}/l(1) \rightarrow \mathcal{O}^*/l[-1] \rightarrow \mu_l[1].$$

Since $(\mathcal{O}^*/l)_{\acute{e}t} = 0$, this recovers 4.8: $\mu_l \simeq \mathbb{Z}/l(1)_{\acute{e}t}$.

EXERCISE 13.3. If F is a Nisnevich sheaf with transfers, modify example 6.20 to show that the sheaf $E^0(F)$ defined in 12.10 is a Nisnevich sheaf with transfers, and that the canonical flasque resolution $F \rightarrow E^*(F)$ is a complex of Nisnevich sheaves with transfers.

LEMMA 13.4. *Let F be a Nisnevich sheaf with transfers. Then:*

- (1) *its cohomology presheaves $H_{Nis}^n(-, F)$ are presheaves with transfers;*
- (2) *for any smooth X , we have $F(X) \cong \mathrm{Hom}_{Sh_{Nis}(Cor_k)}(\mathbb{Z}_{tr}(X), F)$;*
- (3) *for any smooth X and any $n \in \mathbb{Z}$,*

$$H_{Nis}^n(X, F) \cong \mathrm{Ext}_{Sh_{Nis}(Cor_k)}^n(\mathbb{Z}_{tr}(X), F).$$

PROOF. (Cf. 6.3, 6.21, and 6.23.) Assertion (2) is immediate from 13.1 and the Yoneda isomorphism $F(X) \cong \text{Hom}_{\mathbf{PST}}(\mathbb{Z}_{tr}(X), F)$. Now consider the canonical flasque resolution $F \rightarrow E^*(F)$ in $Sh_{Nis}(Sm/k)$. By 13.3, this is a resolution of sheaves with transfers. Since $H_{Nis}^n(-, F)$ is the cohomology of $E^*(F)$ as a presheaf, and hence as a presheaf with transfers, we get part (1).

For part (3), it suffices by part (2) to show that if F is an injective sheaf with transfers and $n > 0$, then $H_{Nis}^n(-, F) = 0$. Since $F \rightarrow E^0(F)$ must split in $Sh_{Nis}(Cor_k)$, $H_{Nis}^n(X, F)$ is a summand of $H_{Nis}^n(X, E^0(F)) = 0$, and must vanish. \square

EXERCISE 13.5. (Cf. 6.25.) Let K be any complex of Nisnevich sheaves of R -modules with transfers and let X be a smooth scheme. Use the fact that $cd_{Nis}(X) \leq \dim(X)$ (by 12.2) to generalize 13.4, by showing that hyperext and hypercohomology agree in the sense that for $n \in \mathbb{Z}$:

$$\text{Ext}^n(R_{tr}(X), K) \cong \mathbb{H}_{Nis}^n(X, K).$$

Since Nisnevich hypercohomology commutes with infinite direct sums, this shows that $\text{Ext}^n(R_{tr}(X), \bigoplus K_\alpha) \cong \bigoplus_\alpha \text{Ext}^n(R_{tr}(X), K_\alpha)$.

EXERCISE 13.6. Let F be a homotopy invariant Nisnevich sheaf of R -modules with transfers. Show that $F(X) \cong \text{Hom}_{\mathbf{D}^-}(C_*\mathbb{Z}_{tr}(X), F)$, where $\mathbf{D}^- = \mathbf{D}^- Sh_{Nis}(Cor_k, R)$.

The following result allows us to bootstrap quasi-isomorphism results from the field level to the sheaf level.

PROPOSITION 13.7. *Let $A \rightarrow B$ be a morphism of complexes of presheaves with transfers. Assume that their cohomology presheaves H^*A and H^*B are homotopy invariant, and that $A(\text{Spec } E) \rightarrow B(\text{Spec } E)$ is a quasi-isomorphism for every field E over k . Then $A_{Zar} \rightarrow B_{Zar}$ is a quasi-isomorphism in the Zariski topology.*

PROOF. Let C be the mapping cone. By the 5-lemma, each $H^n C$ is a homotopy invariant presheaf with transfers, which vanishes on $\text{Spec } E$ for every field E over k . Corollary 11.2 states that $(H^n C)_{Zar} = 0$. This implies that C_{Zar} is acyclic as a complex of Zariski sheaves, i.e., that A_{Zar} and B_{Zar} are quasi-isomorphic in the Zariski topology. \square

The main result of this lecture, 13.12, as well as the next few lectures, depends upon the following result, whose proof will not be completed until 24.1. Theorem 13.8 allows us to bypass the notion of strictly \mathbb{A}^1 -homotopy invariance (see 9.22) used in lecture 9. The case $n = 0$ of 13.8, that F_{Nis} is homotopy invariant, will be completed in 22.3.

THEOREM 13.8. *Let k be a perfect field and F a homotopy invariant presheaf with transfers. Then each presheaf $H_{Nis}^n(-, F_{Nis})$ is homotopy invariant.*

The proofs of the following results are all based upon a combination of theorem 13.8, lemma 13.4, and proposition 13.7.

PROPOSITION 13.9. *Let k be a perfect field. If F is a homotopy invariant Nisnevich sheaf with transfers, then for all n and all smooth X :*

$$H_{Zar}^n(X, F) \cong H_{Nis}^n(X, F).$$

We will prove in 22.15 that F_{Zar} is a presheaf with transfers. This would simplify the proof of 13.9.

PROOF. For $n = 0$ we have $H_{Nis}^0(X, F) = H_{Zar}^0(X, F) = F(X)$ for every sheaf. By the Leray spectral sequence, it now suffices to prove that $H_{Nis}^n(S, F) = 0$ for all $n > 0$ when S is a local scheme. By 13.4 and 13.8, each $H_{Nis}^n(-, F)$ is a homotopy invariant presheaf with transfers. By 11.2, it suffices to show that $H_{Nis}^n(\text{Spec } E, F) = 0$ for every field E over k . But fields are Hensel local rings, and as such have no higher cohomology, i.e., $H_{Nis}^n(\text{Spec } E, -) = 0$ for $n > 0$. \square

PROPOSITION 13.10. *Let C be a bounded above complex of Nisnevich sheaves with transfers, whose cohomology sheaves are homotopy invariant. Then its Zariski and Nisnevich hypercohomology agree:*

$$\mathbb{H}_{Zar}^n(X, C) \cong \mathbb{H}_{Nis}^n(X, C) \quad \text{for all smooth } X \text{ and for all } n.$$

PROOF. We will proceed by descending induction on $n - p$, where $C^i = 0$ for $i > p$. If $\dim X = d$, then $\mathbb{H}_{Zar}^n(X, C) = \mathbb{H}_{Nis}^n(X, C) = 0$ for all $n > p + d$, because $\text{cd}_{Zar}(X)$ and $\text{cd}_{Nis}(X)$ are at most d . By 13.1, both the good Nisnevich truncation τC and the p^{th} -cohomology sheaf $H^p = (C/\tau C)_{Nis}$ are Nisnevich sheaves with transfers. Setting $m = n - p$, we have a diagram

$$\begin{array}{ccccccccc} H_{Zar}^{m-1}(X, H^p) & \longrightarrow & \mathbb{H}_{Zar}^n(X, \tau C) & \longrightarrow & \mathbb{H}_{Zar}^n(X, C) & \longrightarrow & H_{Zar}^m(X, H^p) & \longrightarrow & \mathbb{H}_{Zar}^{n+1}(X, \tau C) \\ \downarrow \cong & & \downarrow \cong & & \downarrow & & \downarrow \cong & & \downarrow \cong \\ H_{Nis}^{m-1}(X, H^p) & \longrightarrow & \mathbb{H}_{Nis}^n(X, \tau C) & \longrightarrow & \mathbb{H}_{Nis}^n(X, C) & \longrightarrow & H_{Nis}^m(X, H^p) & \longrightarrow & \mathbb{H}_{Nis}^{n+1}(X, \tau C). \end{array}$$

The four outer verticals are isomorphisms, by induction and 13.9. The statement now follows from the 5-lemma. \square

EXAMPLE 13.11. The motivic complex $R(i)$ is bounded above, and has homotopy invariant cohomology by 2.19. If A is an R -module, the same is true for $A(i) = A \otimes_R R(i)$. By 13.10, the motivic cohomology of a smooth X could be computed using Nisnevich hypercohomology:

$$H^{n,i}(X, A) = \mathbb{H}_{Zar}^n(X, A(i)) = \mathbb{H}_{Nis}^n(X, A(i)).$$

This is the definition of motivic cohomology used in [VSF00]. Note that the motivic cohomology groups $H^{n,i}(X, A)$ are presheaves with transfers by 13.5.

By 12.12 and 13.5,

$$H^{n,i}(X, A) \cong \text{Hom}_{\mathbf{D}_{Nis}^-}(\mathbf{R}(X), A(i)[n]) \cong \text{Ext}_{Sh_{Nis}(Cor_k)}^n(\mathbf{R}_{tr}(X), A(i)).$$

So motivic cohomology is representable in \mathbf{D}_{Nis}^- and $\mathbf{D}^-(Sh_{Nis}(Cor_k))$.

THEOREM 13.12. *Let k be a perfect field and F a presheaf with transfers such that $F_{Nis} = 0$. Then $(C_*F)_{Nis} \simeq 0$ in the Nisnevich topology, and $(C_*F)_{Zar} \simeq 0$ in the Zariski topology.*

PROOF. Let F be a presheaf with transfers such that $F_{Nis} = 0$. We will first prove that $(C_*F)_{Nis} \simeq 0$ or, equivalently, that the homology presheaves $H_i = H_i C_*F$ satisfy $(H_i)_{Nis} = 0$ for all i . For $i < 0$ this is trivial; $C_i F = 0$ implies that $H_i C_*F = 0$. Since $(H_0)_{Nis}$ is a quotient of $F_{Nis} = 0$, it is also true for $i = 0$.

We shall proceed by induction on i , so we assume that $(H_j)_{Nis} = 0$ for all $j < i$. That is, we assume that $\tau(C_*F)_{Nis} \simeq (C_*F)_{Nis}$, where $\tau(C_*F)_{Nis}$ denotes the subcomplex of $(C_*F)_{Nis}$ obtained by good truncation at level i :

$$\tau(C_*F)_{Nis} \text{ is } \cdots \rightarrow (C_{i+1}F)_{Nis} \rightarrow (C_i F)_{Nis} \rightarrow d(C_i F)_{Nis} \rightarrow 0.$$

There is a canonical morphism $\tau(C_*F)_{Nis} \rightarrow (H_i)_{Nis}[i]$ and hence a morphism $f : (C_*F)_{Nis} \rightarrow (H_i)_{Nis}[i]$ in the derived category \mathbf{D}_{Nis}^- . Since f induces an isomorphism on the i th homology sheaves, it suffices to prove that $f = 0$.

The presheaf with transfers H_i is homotopy invariant by 2.19, so by 13.1 and 13.8 the sheaf $G = (H_i)_{Nis}$ satisfies the hypothesis of 12.19. Since $F_{Nis} = 0$, 12.19 yields

$$\mathrm{Hom}_{\mathbf{D}_{Nis}^-}((C_*F)_{Nis}, (H_i)_{Nis}[i]) \cong \mathrm{Hom}_{\mathbf{D}_{Nis}^-}(F_{Nis}, (H_i)_{Nis}[i]) = 0.$$

Hence $f = 0$ in \mathbf{D}_{Nis}^- , and this implies that $(H_i)_{Nis} = 0$.

We can now prove that $C_*F_{Zar} \simeq 0$. Each cohomology presheaf $H^i = H^i C_*F$ is a homotopy invariant presheaf with transfers by 2.19. Since $(C_*F)_{Nis} \simeq 0$, we have $C_*(F)(\mathrm{Spec} E) \simeq 0$ for every finitely generated field extension E of k (and hence for every field over k). Indeed, E is $\mathcal{O}_{X,x}^h$ for the generic point of some smooth X . Now apply 13.7 to $C_*F \rightarrow 0$. \square

Here is a stalkwise restatement of theorem 13.12.

COROLLARY 13.13. *Let k be a perfect field and F a presheaf with transfers so that $F(\mathrm{Spec} \mathcal{O}_{X,x}^h) = 0$ for all smooth X and all $x \in X$. Then $(C_*F)(\mathrm{Spec} \mathcal{O}_{X,x}) \simeq 0$ for all X and all $x \in X$.*

COROLLARY 13.14. *Let $f : C_1 \rightarrow C_2$ be a map of bounded above cochain complexes of presheaves with transfers. If f induces a quasi-isomorphism over all Hensel local rings $\mathrm{Spec} \mathcal{O}_{X,x}^h$, then $\mathrm{Tot}(C_*C_1) \rightarrow \mathrm{Tot}(C_*C_2)$ induces a quasi-isomorphism over all local rings.*

PROOF. Let $K = \mathrm{cone}(f)$ denote the mapping cone of f . By assumption, each $H^p K$ is a presheaf with transfers which vanishes on all Hensel local schemes, i.e., $K_{Nis} \simeq 0$. By 13.12, $C_*H^p K \simeq 0$ in the Zariski topology.

Since K is a bounded above cochain complex, the double complex $C_*(K)$ is bounded. Hence the usual spectral sequence of a double complex (see [Wei94, 5.6.2]) converges to $H_* \mathrm{Tot} C_*(K)$. Since $C_q K(X) = K(X \times \Delta^q)$ we have $H^p C_q K =$

$C_q H^p K$ for all p and q , and we have seen that each $H^q C_* H^p K$ vanishes on every local scheme X . The resulting collapse in the spectral sequence shows that $H_* \text{Tot} C_*(K)$ vanishes on every local scheme, which yields the result. \square

If $\mathcal{U} = \{U_1, \dots, U_n\}$ is a Zariski covering of X , we saw in 6.12 that the Čech complex

$$\mathbb{Z}_{\text{tr}}(\check{\mathcal{U}}) : 0 \rightarrow \mathbb{Z}_{\text{tr}}(U_1 \cap \dots \cap U_n) \rightarrow \dots \rightarrow \bigoplus_i \mathbb{Z}_{\text{tr}}(U_i) \rightarrow 0$$

is a resolution of $\mathbb{Z}_{\text{tr}}(X)$ in the étale topology (and the Nisnevich topology). Surprisingly, this gets even better when we apply C_* .

PROPOSITION 13.15. *If \mathcal{U} is a Zariski covering of X then the Čech resolution $\text{Tot} C_* \mathbb{Z}_{\text{tr}}(\check{\mathcal{U}}) \rightarrow C_* \mathbb{Z}_{\text{tr}}(X)$ is a quasi-isomorphism in the Zariski topology.*

PROOF. Apply 13.14 to 6.14. \square

COROLLARY 13.16. *If E is a vector bundle over X , $C_* \mathbb{Z}_{\text{tr}}(E) \rightarrow C_* \mathbb{Z}_{\text{tr}}(X)$ is a quasi-isomorphism.*

PROOF. Choose a Zariski cover \mathcal{U} of X on which E is a trivial bundle. By 2.24 and 13.15, the left vertical and the horizontal maps are quasi-isomorphisms in the diagram:

$$\begin{array}{ccc} C_* \mathbb{Z}_{\text{tr}}(\check{\mathcal{U}} \times_X E) & \xrightarrow{\simeq} & C_* \mathbb{Z}_{\text{tr}}(E) \\ \simeq \downarrow & & \downarrow \\ C_* \mathbb{Z}_{\text{tr}}(\check{\mathcal{U}}) & \xrightarrow{\simeq} & C_* \mathbb{Z}_{\text{tr}}(X). \end{array}$$

Hence the right vertical map is a quasi-isomorphism. \square

EXAMPLE 13.17. Applying 13.15 to the usual cover of \mathbb{P}^1 (by $\mathbb{P}^1 - \{0\}$ and $\mathbb{P}^1 - \{\infty\}$) allows us to deduce that $C_*(\mathbb{Z}_{\text{tr}}(\mathbb{P}^1)/\mathbb{Z}) \simeq C_* \mathbb{Z}_{\text{tr}}(\mathbb{G}_m)[1] = \mathbb{Z}(1)[2]$ for the Zariski topology, because $C_* \mathbb{Z}_{\text{tr}}(\mathbb{A}^1)/\mathbb{Z} \simeq 0$ by 2.24. This was already observed in example 6.15 for the étale topology. This example will be generalized in theorem 15.2 below.

PROPOSITION 13.18. *Let k be a perfect field and F a homotopy invariant Nisnevich sheaf with transfers. Then the Zariski sheaf H_{Zar} associated to $H(U) = H^q(U \times X, F)$ vanishes for every $q > \dim(X)$.*

PROOF. By 13.4 and 13.8, $H(U)$ is a homotopy invariant presheaf with transfers. If E is a field over k then the Nisnevich cohomological dimension of X_E is at most $\dim(X_E) = \dim(X)$, so $H(\text{Spec } E) = H^q(X_E, F) = 0$. By 11.2, $H_{\text{Zar}} = 0$. \square

We now consider the behavior of cohomology with respect to blow-ups along smooth centers. We assume that k is perfect in order to invoke 13.8.

PROPOSITION 13.19. *Let $p : X' \rightarrow X$ be the blow-up of a smooth X with a smooth center Z . Let C (respectively, Q) denote the cokernel of $\mathbb{Z}(X') \rightarrow \mathbb{Z}(X)$ (respectively, $\mathbb{Z}(Z) \oplus \mathbb{Z}(X') \rightarrow \mathbb{Z}(X)$). Then for any homotopy invariant Nisnevich sheaf with transfers F , $\text{Ext}^n(C, F) = \text{Ext}^n(Q, F) = 0$ for all n .*

If L is a complex of Nisnevich sheaves with transfers whose cohomology sheaves are homotopy invariant, $\text{Ext}^n(C, L) = \text{Ext}^n(Q, L) = 0$ for all n .

PROOF. Set $U = X - Z$; by assumption $\mathbb{Z}(U)$ is a subsheaf of both $\mathbb{Z}(X')$ and $\mathbb{Z}(X)$. By the 5-lemma, C is the cokernel of $\mathbb{Z}(X')/\mathbb{Z}(U) \rightarrow \mathbb{Z}(X)/\mathbb{Z}(U)$.

Since Z is smooth, and exercise 12.20 allows us to pass to an étale neighborhood of Z without changing C or Q , we may assume that $X = T \times \mathbb{A}^d$ and that Z is identified with $T \times 0$. But then the projection $X' \rightarrow T \times \mathbb{P}^{d-1}$ defining the blow-up is a vector bundle with fiber \mathbb{A}^1 , with section $Z' = Z \times_X X' \cong T \times \mathbb{P}^{d-1}$. Since F is a homotopy invariant presheaf with transfers, theorem 13.8 implies that $\text{Ext}^n(\mathbb{Z}(X'), F) \cong H^n(X', F) = H^n(T \times \mathbb{P}^{d-1}, F)$ and $H^n(T \times \mathbb{A}^d, F) \cong H^n(Z, F)$.

The result for F is now a straightforward calculation. Since $Z \rightarrow X$ factors through X' in this special case, we have $C = Q$. If K denotes the kernel of $\mathbb{Z}(X) \rightarrow C$, then $\text{Ext}^*(\mathbb{Z}(Z'), F) \cong \text{Ext}^*(\mathbb{Z}(X'), F)$ implies that $\text{Ext}^*(K, F) \cong \text{Ext}^*(\mathbb{Z}(Z), F)$. This in turn implies that $\text{Ext}^*(\mathbb{Z}(X), F) \rightarrow \text{Ext}^*(K, F)$ is an isomorphism, whence $\text{Ext}^*(C, F) = 0$.

The result for L follows from the hyperext spectral sequence $E_2^{p,q} = \text{Ext}^q(-, H^p L) \Rightarrow \text{Ext}^{p+q}(-, L)$. \square

COROLLARY 13.20. *Let F be a homotopy invariant Nisnevich sheaf with transfers, and let $p : X' \rightarrow X$ be the blow-up of a smooth X along a smooth center Z . Then there is a long exact sequence in Nisnevich cohomology (and, by 13.9, Zariski cohomology)*

$$\cdots \rightarrow H^{i-1}(Z', F) \rightarrow H^i(X, F) \rightarrow H^i(X', F) \oplus H^i(Z, F) \rightarrow H^i(Z', F) \rightarrow \cdots$$

There is an analogous long exact sequence of hypercohomology groups $\mathbb{H}^(-, L)$ (either Nisnevich or Zariski by 13.10) if L is a complex of Nisnevich sheaves with transfers whose cohomology sheaves are homotopy invariant.*

PROOF. Since $H_{\text{Nis}}^i(X, F) \cong \text{Ext}^i(\mathbb{Z}(X), F)$, and $\text{Ext}^*(Q, F) = 0$ by 13.19, this follows from the Ext sequences associated to exercise 12.22. \square

COROLLARY 13.21. *Let $X_r \rightarrow X_{r-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X$ be a sequence of blow-ups along smooth centers and let $C = \mathbb{Z}(X_r, X)$ be the sheaf cokernel of $\mathbb{Z}(X_r) \rightarrow \mathbb{Z}(X)$. Then $\text{Ext}^*(C, F) = 0$ if F is a homotopy invariant Nisnevich sheaf with transfers.*

PROOF. We proceed by induction on r , the case $r = 1$ being 13.19. If $X_1 \rightarrow X$ has center Z , and we set $Z_1 = Z \times_X X_1$, $Z_r = Z \times_X X_r$, then $Z_r \rightarrow Z_1$ is a composition of $r - 1$ blow-ups along smooth centers. Consider the diagram

$$\begin{array}{ccccccc} \mathbb{Z}(X_r) & \longrightarrow & \mathbb{Z}(X_1) & \longrightarrow & \mathbb{Z}(X_r, X_1) & \longrightarrow & 0 \\ \downarrow = & & \downarrow & & \downarrow & & \\ \mathbb{Z}(X_r) & \longrightarrow & \mathbb{Z}(X) & \longrightarrow & \mathbb{Z}(X_r, X) & \longrightarrow & 0. \end{array}$$

It follows from 12.22 and a diagram chase that there is an exact sequence

$$0 \rightarrow \mathbb{Z}(Z_r, Z_1) \rightarrow \mathbb{Z}(X_r, X_1) \rightarrow C \rightarrow \mathbb{Z}(X_1, X) \rightarrow 0.$$

By induction, $\text{Ext}^*(\mathbb{Z}(Z_r, Z_1), F)$ and $\text{Ext}^*(\mathbb{Z}(X_r, X_1), F)$ vanish. Since $\text{Ext}^*(\mathbb{Z}(X_1, X), F)$ also vanishes, it follows from the Ext sequences that $\text{Ext}^*(C, F) = 0$. \square

Cdh sheaves with transfers

The main result of this section will be theorem 13.25.

LEMMA 13.22. *Let C be the sheaf cokernel of $\mathbb{Z}(X') \rightarrow \mathbb{Z}(X)$, where $X' \rightarrow X$ is a cdh cover. Then $C_{cdh} = 0$.*

PROOF. To see this, pick a generator $f \in \text{Hom}(U, X)$ of $C(U)$ and consider the pullback cdh cover $U' = U \times_X X' \rightarrow U$ of $X' \rightarrow X$ along f . The image of f in $\text{Hom}(U', X)$ comes from $f' \in \text{Hom}(U', X')$ and so vanishes in $C(U')$. Since $U' \rightarrow U$ is a cdh cover, f vanishes in $C_{cdh}(U)$. \square

PROPOSITION 13.23. *Suppose that k admits resolution of singularities. If F is a Nisnevich sheaf on Sm/k such that $F_{cdh} = 0$, and H is a homotopy invariant Nisnevich sheaf with transfers, then $\text{Ext}_{\text{Nis}}^n(F, H) = 0$ for all n .*

PROOF. We proceed by induction, the case $n < 0$ being a definition. Consider the canonical surjection of sheaves

$$\bigoplus_{\alpha \in F(X)} \mathbb{Z}(X) \xrightarrow{\pi} F.$$

Since $F_{cdh} = 0$, 12.30 implies that π factors through $\bigoplus_{\alpha} C_{\alpha} \xrightarrow{p} F$, where each C_{α} is the sheaf cokernel of a sequence of blow-ups along smooth centers. Write K for the kernel of the surjection p . The sheaf K_{cdh} vanishes because it is a subsheaf of $\bigoplus (C_{\alpha})_{cdh}$, which is zero by 13.22. By 13.21, each $\text{Ext}^*(C_{\alpha}, H) = 0$, and therefore $\text{Ext}^n(F, H) \cong \text{Ext}^{n-1}(K, H)$, which is zero by induction on n . \square

COROLLARY 13.24. *Let H be a homotopy invariant Nisnevich sheaf with transfers. Then $H(X)$ injects into $H(X')$ for any cdh cover $X' \rightarrow X$.*

In particular, $H(X) \rightarrow H_{cdh}(X)$ is an injection for all smooth X .

PROOF. Let C be the sheaf cokernel of $\mathbb{Z}(X') \rightarrow \mathbb{Z}(X)$. Since $\text{Hom}(-, H)$ is left exact, $0 \rightarrow \text{Hom}(C, H) \rightarrow \text{Hom}(\mathbb{Z}(X), H) \rightarrow \text{Hom}(\mathbb{Z}(X'), H)$ is exact. But $\text{Hom}(C, H) = 0$ by 13.22 and 13.23, and $\text{Hom}(\mathbb{Z}(X), H) = H(X)$ by Yoneda. \square

THEOREM 13.25. *Assume that resolution of singularities holds over a perfect field k . Let F be a Nisnevich sheaf with transfers such that $F_{cdh} = 0$. Then the complex $C_*(F)$ is acyclic.*

PROOF. If $C_*(F)$ is not acyclic, there is a smallest $n \geq 0$ such that the sheaf $H = H_n(C_*(F))$ is non-zero. Using the good truncation of $C_*(F)$, we define a non-zero map $C_*(F) \rightarrow H[n]$ in $\mathbf{D}_{\text{Nis}}^-$. By 2.19 and 13.8, the sheaf with transfers H is strictly homotopy invariant. By 12.19, $\text{Hom}_{\mathbf{D}_{\text{Nis}}^-}(C_*F, H[n]) \cong \text{Hom}_{\mathbf{D}_{\text{Nis}}^-}(F, H[n]) = \text{Ext}_{\text{Nis}}^n(F, H)$. But this Ext group is zero by 13.23. \square

THEOREM 13.26. *Assume that resolution of singularities holds over k . Let $X' \rightarrow X$ be an abstract blow-up with center Z , and set $Z' = Z \times_X X'$. Then there is a distinguished triangle in $\mathbf{D}^-(Sh_{Nis}(Cor_k))$:*

$$C_*\mathbb{Z}_{tr}(Z') \rightarrow C_*\mathbb{Z}_{tr}(Z) \oplus C_*\mathbb{Z}_{tr}(X') \rightarrow C_*\mathbb{Z}_{tr}(X) \rightarrow C_*\mathbb{Z}_{tr}(Z')[1].$$

PROOF. Let Φ be the sequence $\mathbb{Z}_{tr}(Z') \rightarrow \mathbb{Z}_{tr}(Z) \oplus \mathbb{Z}_{tr}(X') \rightarrow \mathbb{Z}_{tr}(X)$. Let Q denote the cokernel of $\mathbb{Z}_{tr}(X') \oplus \mathbb{Z}_{tr}(Z) \rightarrow \mathbb{Z}_{tr}(X)$; by exercise 12.22, $\Phi \cong Q$. We have to show that $C_*(Q)$ is acyclic; by 13.25, it suffices to show that $Q_{cdh} = 0$.

Pick a finite set of elementary correspondences $W_i \subset U \times X$ representing generators a_i of $\mathbb{Z}_{tr}(X)(U)$ and hence $Q(U)$. We may assume that no W_i lies in $U \times Z$. Let W'_i be the proper transform of W_i in $U \times X'$. By platication [RG71], there is a blow-up $U' \rightarrow U$ such that the proper transforms W''_i of W'_i in $U' \times X'$ are flat over U' . Since each $W''_i \rightarrow U'$ is generically finite, and flat, it is finite. Using resolution of singularities, we can find $U'' \rightarrow U'$ with U'' smooth such that $U'' \rightarrow U$ is a (proper birational) cdh cover; we may replace U' by U'' so that the W''_i represent elements $b_i \in \mathbb{Z}_{tr}(X')(U'')$. But the map $\mathbb{Z}_{tr}(X)(U) \rightarrow \mathbb{Z}_{tr}(X)(U'')$ sends each a_i to the image of b_i , and hence the injection $Q_{cdh}(U) \rightarrow Q_{cdh}(U'')$ sends each a_i to zero. By 12.30, this proves that $Q_{cdh} = 0$. \square

PROPOSITION 13.27. *Assume that resolution of singularities holds over k , and let F be a homotopy invariant Nisnevich sheaf with transfers. Then for all smooth X :*

- $F_{cdh}(X) \cong F(X)$;
- $H^n_{cdh}(X, F_{cdh}) \cong H^n_{Nis}(X, F)$ for all n .

PROOF. Let us first show that $F_{cdh}(X) = F(X)$. By 13.24, $F(X) \rightarrow F_{cdh}(X)$ is an injection for all smooth X . Letting G be the sheaf cokernel, we have an exact sequence of Nisnevich sheaves:

$$0 \rightarrow F \rightarrow F_{cdh} \rightarrow G \rightarrow 0.$$

By 13.23, $\text{Ext}^1(G, F) = 0$, and therefore the sequence splits. Hence F is a direct summand of F_{cdh} , and hence the restriction of a cdh sheaf to Sm/k . So $G = 0$.

Let $F \rightarrow I^*$ be a cdh injective resolution of F . The restriction $I^*|_{Nis}$ of I^* to the Nisnevich topology is a complex of injective Nisnevich sheaves. It suffices to show that this is also a resolution for the Nisnevich topology. Let B^i, Z^i and $H^i = Z^i/B^i$ be the i^{th} boundaries, cycles and cohomology sheaves of the complex $I^*|_{Nis}$, respectively. Since $H^0 = F$ by left exactness, we only need to show that $H^i = 0$ for $i > 0$. If not, there is a minimal $i > 0$ such that $H^i \neq 0$. By hypothesis $(H^i)_{cdh} = 0$, so by 13.23 and dimension shifting [Wei94, Exercise 2.4.3], $0 = \text{Ext}^{i+1}(H^i, F) \cong \text{Ext}^1(H^i, B^i)$. This implies that the sequence $0 \rightarrow B^i \rightarrow Z^i \rightarrow H^i \rightarrow 0$ splits, and therefore that H^i is a summand of Z^i . Since Z^i is a cdh sheaf, so is H^i . But then $H^i = (H^i)_{cdh} = 0$. \square

Part 4

The Triangulated Category of Motives

The category of motives

In this lecture, we define the triangulated category of (effective) motives over k , and the motive of a scheme in this category. The construction of $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R)$ is parallel to the construction of $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ in 9.2, but more central. We list the main properties of this category in 14.5 below. If k admits resolution of singularities, this category allows us to extend motivic cohomology to all schemes of finite type, as a cdh hypercohomology group. If $\mathbb{Q} \subseteq R$, we will show that $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R)$ and $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ are equivalent.

Write \mathbf{D}^- for $\mathbf{D}^- \text{Sh}_{Nis}(\text{Cor}_k, R)$, and let $\mathcal{E}_{\mathbb{A}^1}$ denote the smallest thick subcategory of \mathbf{D}^- containing every $R_{tr}(X \times \mathbb{A}^1) \rightarrow R_{tr}(X)$ and closed under direct sums. (See 9.1 and 9.2.) The quotient $\mathbf{D}^- / \mathcal{E}_{\mathbb{A}^1}$ is the localization $\mathbf{D}^- [W_{\mathbb{A}^1}^{-1}]$, where $W_{\mathbb{A}^1} = W_{\mathcal{E}_{\mathbb{A}^1}}$ is the class of maps in \mathbf{D}^- whose cone is in $\mathcal{E}_{\mathbb{A}^1}$. A map in $W_{\mathbb{A}^1}$ is called an \mathbb{A}^1 -**weak equivalence**.

As pointed out in 9.3, it follows from 2.24 and 14.4 below that $\mathcal{E}_{\mathbb{A}^1}$ is the thick subcategory of all complexes E in \mathbf{D}^- such that $C_*(E)$ is acyclic.

DEFINITION 14.1. The triangulated category of motives over k is defined to be the localization $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R) = \mathbf{D}^- [W_{\mathbb{A}^1}^{-1}]$ of $\mathbf{D}^- = \mathbf{D}^- \text{Sh}_{Nis}(\text{Cor}_k, R)$. (Cf. 9.2.) If X is a smooth scheme over k , we write $M(X)$ for the class of $\mathbb{Z}_{tr}(X)$ in $\mathbf{DM}_{Nis}^{\text{eff},-}(k, \mathbb{Z})$ and call it the **motive** of X .

We define $\mathbf{DM}_{gm}^{\text{eff}}(k, R)$ to be the thick subcategory of $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R)$ generated by the motives $M(X)$, where X is smooth over k . Objects in $\mathbf{DM}_{gm}^{\text{eff}}(k, R)$ will be called **effective geometric motives**. If k admits resolution of singularities, it follows from (14.5.3) and (14.5.5) that $\mathbf{DM}_{gm}^{\text{eff}}$ contains $M(Y)$ for every Y in Sch/k , and is generated by $M(X)$, where X is smooth and projective.

In 8.17, we showed that the derived category $\mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$ is a tensor triangulated category. The same argument works in the Nisnevich topology for $\mathbf{D}^- \text{Sh}_{Nis}(\text{Cor}_k, R)$. Here are the details.

DEFINITION 14.2. If C and D are bounded above complexes of presheaves with transfers, we write $C \otimes_{L, Nis}^{\text{tr}} D$ for $(C \otimes_L^{\text{tr}} D)_{Nis}$. Because 6.12 holds for the Nisnevich topology, the Nisnevich analogues of 8.14, 8.15, 8.16, 8.17, and 8.18 hold. In particular, the derived category \mathbf{D}^- of bounded above complexes of Nisnevich sheaves with transfers is a tensor triangulated category under $\otimes_{L, Nis}^{\text{tr}}$. By 8.10, $M(X) \otimes_{L, Nis}^{\text{tr}} M(Y) \cong M(X \times Y)$.

Given 14.2, the proofs of 9.5 and 9.6 go through to show that the tensor $\otimes_{L, Nis}^{tr}$ on \mathbf{D}^- also endows the localization $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ of $\mathbf{D}^- Sh_{Nis}(Cor_k, R)$ with the structure of a tensor triangulated category.

The category $\mathbf{DM}^-(k, R)$ is obtained from $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ by inverting the Tate twist operation $M \mapsto M(1) = M \otimes_{L, Nis}^{tr} R(1)$. Thus every object in $\mathbf{DM}^-(k, R)$ is isomorphic to $M(-n)$ for some $n \geq 0$ and some M in $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$. For any coefficients R , it will follow from 8A.11 and 15.8 below that $\mathbf{DM}^-(k, R)$ is always a tensor triangulated category. The localization $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R) \rightarrow \mathbf{DM}^-(k, R)$ is fully faithful, by Voevodsky's Cancellation Theorem 16.25 below.

The category $\mathbf{DM}_{gm}(k, R)$ of **geometric motives** is obtained from $\mathbf{DM}_{gm}^{\text{eff}}(k, R)$ by inverting the Tate twist operation $M \mapsto M(1) = M \otimes_{L, Nis}^{tr} R(1)$. From the previous paragraph, it is clear that $\mathbf{DM}_{gm}(k, R)$ is a full tensor triangulated subcategory of $\mathbf{DM}^-(k, R)$ and that the localization $\mathbf{DM}_{gm}^{\text{eff}}(k, R) \rightarrow \mathbf{DM}_{gm}(k, R)$ is fully faithful.

REMARK 14.3. Because sheafification is exact, it induces a triangulated functor from $\mathbf{D}_{Nis}^- = \mathbf{D}^-(Sh_{Nis}(Cor_k, R))$ to $\mathbf{D}_{\acute{e}t}^- = \mathbf{D}^-(Sh_{\acute{e}t}(Cor_k, R))$. By lemma 9.15 and definition 14.2, we have $(K \otimes_{L, Nis}^{tr} L)_{\acute{e}t} = K \otimes_{L, \acute{e}t}^{tr} L$. Comparing definitions, we see that $\mathbf{D}_{Nis}^- \rightarrow \mathbf{D}_{\acute{e}t}^-$ sends Nisnevich \mathbb{A}^1 -weak equivalences to étale \mathbb{A}^1 -weak equivalences, so it induces a tensor triangulated functor σ from $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ to $\mathbf{DM}_{\acute{e}t}^{\text{eff}, -}(k, R)$. We will show in 14.30 below that σ is an equivalence when $R = \mathbb{Q}$.

Our definitions of $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ and $M(X)$ are equivalent to the definitions in [TriCa, p. 205]. This follows by comparing the definition in *loc. cit.* to theorem 14.11 below, using the following lemma.

LEMMA 14.4. *For every bounded above complex K of sheaves of R -modules with transfers, the morphism $K \rightarrow \text{Tot}C_*(K)$ is an \mathbb{A}^1 -weak equivalence. Hence $K \cong \text{Tot}C_*(K)$ in $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$.*

In particular, there is a natural isomorphism $M(X) \cong C_\mathbb{Z}_{tr}(X)$.*

PROOF. The proof of lemmas 9.12 and 9.15 go through in this setting. \square

PROPERTIES 14.5. We now summarize the main properties of the category $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$ for the convenience of the reader.

- By 14.4, we have $M(X) = \mathbb{Z}_{tr}(X) \cong C_*\mathbb{Z}_{tr}(X)$. By 8.10, we see that $M(X) \otimes M(Y) \cong M(X \times Y)$, and 2.24 yields $M(X) \cong M(X \times \mathbb{A}^1)$.
- For every smooth X and every Y , it follows from 14.16 that $\mathbb{H}^n(X, C_*R_{tr}(Y)) \cong \text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff}, -}}(M(X), M(Y)[n])$. In particular,

$$H^{n,i}(X, R) \cong \text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff}, -}}(M(X), R(i)[n]).$$

For non-smooth X , $H^{n,i}(X, R)$ is defined via this formula; see 14.17.

- (Mayer-Vietoris) For each open cover $\{U, V\}$ of a smooth scheme X , proposition 13.15 yields the Mayer-Vietoris triangle in $\mathbf{DM}_{Nis}^{\text{eff}, -}(k, R)$:

$$(14.5.1) \quad M(U \cap V) \rightarrow M(V) \oplus M(U) \rightarrow M(X) \rightarrow M(U \cap V)[1].$$

- (Vector bundle) If $E \rightarrow X$ is a vector bundle, by 13.16 we have an isomorphism $M(E) \xrightarrow{\cong} M(X)$.
- (Projective bundle) We will prove in 15.12 that if $\mathbb{P}(\mathcal{E}) \rightarrow X$ is a projective bundle of rank $n + 1$, then the canonical map induces an isomorphism:

$$(14.5.2) \quad \bigoplus_{i=0}^n M(X)(i)[2i] \xrightarrow{\cong} M(\mathbb{P}(\mathcal{E})).$$

- (Blow-up triangle) Assume that resolution of singularities holds over k . Let $X' \rightarrow X$ be an abstract blow-up with center Z , and set $Z' = Z \times_X X'$. By 13.26, there is a distinguished triangle:

$$(14.5.3) \quad M(Z') \rightarrow M(X') \oplus M(Z) \rightarrow M(X) \rightarrow M(Z')[1].$$

If moreover X and Z are smooth, and Z has codimension c , we show in 15.13 that (14.5.2) and (14.5.3) easily yield an isomorphism:

$$(14.5.4) \quad M(X') \cong M(X) \oplus \left(\bigoplus_{i=1}^{c-1} M(Z)(i)[2i] \right).$$

- (Gysin triangle) Let X be a smooth scheme over k and Z a smooth closed subscheme of X of codimension c . We will show in 15.15 that there is a distinguished triangle:

$$(14.5.5) \quad M(X - Z) \rightarrow M(X) \rightarrow M(Z)(c)[2c] \rightarrow M(X - Z)[1].$$

- (Cancellation) Assume that k admits resolution of singularities. Let M and N be in $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R)$. Then we will see in 16.25 that there is an isomorphism $\text{Hom}(M, N) \rightarrow \text{Hom}(M(1), N(1))$.
- (Chow motives) We will show in 20.1 that Grothendieck's category of effective Chow motives embeds contravariantly into $\mathbf{DM}_{gm}^{\text{eff}}(k, \mathbb{Z})$, and hence into $\mathbf{DM}_{Nis}^{\text{eff},-}(k, \mathbb{Z})$, in the sense that if X and Y are two smooth projective schemes, then:

$$(14.5.6) \quad \text{Hom}(M(X), M(Y)) \cong CH^{\dim X}(X \times Y) = \text{Hom}_{\text{Chow}}(Y, X).$$

We will define the notion of a motive with compact support in lecture 16. We will investigate its properties there and in lecture 20.

Nisnevich \mathbb{A}^1 -local complexes

In this section we will show that $\mathbf{DM}_{Nis}^{\text{eff},-}(k, R)$ can be identified with the full subcategory \mathcal{L} of \mathbb{A}^1 -local complexes in $\mathbf{D}^- = \mathbf{D}^-(\text{Sh}_{Nis}(\text{Cor}_k, R))$.

DEFINITION 14.6. As is 9.17, we say that an object L of \mathbf{D}^- is called \mathbb{A}^1 -**local** (for the Nisnevich topology) if $\text{Hom}_{\mathbf{D}^-}(-, L)$ sends \mathbb{A}^1 -weak equivalences to isomorphisms. We write \mathcal{L} for the full subcategory of \mathbb{A}^1 -local objects in \mathbf{D}^- . The proof of 9.19 goes through in the Nisnevich setting to show that if L is \mathbb{A}^1 -local then for every K :

$$(14.6.1) \quad \text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff},-}}(K, L) \cong \text{Hom}_{\mathbf{D}^-}(K, L).$$

REMARK 14.7. We will see in 14.9 below that C_* is a functor from \mathbf{D}^- to \mathcal{L} . Moreover, $\mathrm{Hom}_{\mathcal{L}}(C_*(K), L) \cong \mathrm{Hom}_{\mathbf{D}^-}(K, L)$ for every L in \mathcal{L} and K in \mathbf{D}^- by 14.4 and definition 14.6. Hence C_* is the left adjoint to the inclusion $\mathcal{L} \subset \mathbf{D}^-$.

Let F be a Nisnevich sheaf with transfers. Then F is \mathbb{A}^1 -local if and only if F is homotopy invariant, because the proof of 9.24 goes through using 13.4 and 13.8. This is the easy case of the following proposition.

PROPOSITION 14.8. *Let k be a perfect field and K a bounded above cochain complex of Nisnevich sheaves of R -modules with transfers. Then K is \mathbb{A}^1 -local if and only if the sheaves $a_{\mathrm{Nis}}(H^n K)$ are all homotopy invariant.*

Hence \mathcal{L} is the category $\mathbf{D}_{hi}^-(\mathrm{Sh}_{\mathrm{Nis}}(\mathrm{Cor}_k, R))$ of complexes with homotopy invariant cohomology sheaves.

PROOF. Suppose first that the cohomology sheaves of K are homotopy invariant. By 13.8 applied to $F = a_{\mathrm{Nis}}(H^q K)$, the presheaves $H_{\mathrm{Nis}}^n(-, F)$ are homotopy invariant. As in the proof of 9.24, this implies that each $a_{\mathrm{Nis}}(H^q K)$ is \mathbb{A}^1 -local. Because $cd_{\mathrm{Nis}}(X) < \infty$, the hyperext spectral sequence (see [Wei94, 5.7.9])

$$E_2^{pq}(X) = \mathrm{Ext}^p(R_{tr}(X), a_{\mathrm{Nis}}H^q K) \implies \mathrm{Hom}_{\mathbf{D}^-}(R_{tr}(X), K[p+q])$$

is bounded and converges. The map f induces a morphism from it to the corresponding spectral sequence for $X \times \mathbb{A}^1$. By the Comparison Theorem ([Wei94, 5.2.12]), f induces an isomorphism from $\mathrm{Hom}_{\mathbf{D}^-}(R_{tr}(X)[n], K)$ to $\mathrm{Hom}_{\mathbf{D}^-}(R_{tr}(X \times \mathbb{A}^1)[n], K)$ for each n . By 9.20, K is \mathbb{A}^1 -local.

Now suppose that K is \mathbb{A}^1 -local. The cohomology presheaves of $K' = \mathrm{Tot}C_*(K)$ are homotopy invariant by 2.19. Theorem 13.8 applied to the cohomology presheaves $H^q K'$ shows that the sheaves $a_{\mathrm{Nis}}(H^q K')$ are homotopy invariant. The first part of this proof shows that K' is \mathbb{A}^1 -local. By lemma 14.4, the canonical map $K \rightarrow K'$ is an \mathbb{A}^1 -weak equivalence. By 9.21, which goes through for the Nisnevich topology, $K \rightarrow K'$ is an isomorphism in \mathbf{D}^- . Hence the sheaves $a_{\mathrm{Nis}}(H^n K) \cong a_{\mathrm{Nis}}(H^n K')$ are homotopy invariant. \square

COROLLARY 14.9. *Let k be a perfect field and K a bounded above cochain complex of Nisnevich sheaves of R -modules with transfers. If the presheaves $H^n(K)$ are all homotopy invariant, then K is \mathbb{A}^1 -local.*

In particular, $C_(K)$ is \mathbb{A}^1 -local, and if K is \mathbb{A}^1 -local then $K \cong C_*(K)$ in \mathbf{D}^- .*

PROOF. Combine 13.8 and 14.8. The hypothesis applies to $C_*(K)$ by 2.19, and the final assertion follows immediately from 14.4, as in the étale case 9.32. \square

EXAMPLE 14.10. Here is an example to show that the converse does not hold in 14.9. Consider the complex K of example 6.15:

$$0 \rightarrow \mathbb{Z}_{tr}(\mathbb{G}_m) \rightarrow 2\mathbb{Z}_{tr}(\mathbb{A}^1, 1) \rightarrow \mathbb{Z}_{tr}(\mathbb{P}^1, 1) \rightarrow 0.$$

Evaluating at $\mathrm{Spec}(k)$ and at \mathbb{A}^1 , it is easy to see that the cohomology presheaf $H^2 K$ is not homotopy invariant (consider an embedding of \mathbb{A}^1 in \mathbb{P}^1 whose image contains both 0 and 1). On the other hand K is \mathbb{A}^1 -local, because its cohomology sheaves $a_{\mathrm{Nis}}H^*(K)$ vanish by 6.14.

If E and F are in \mathcal{L} , then we define $E \otimes_{\mathcal{L}} F = \mathrm{Tot}C_*(E \otimes_{L, Nis}^{tr} F)$.

THEOREM 14.11. *The category $(\mathcal{L}_{Nis}, \otimes_{\mathcal{L}})$ is a tensor triangulated category, and the canonical functor*

$$\mathcal{L}_{Nis} \rightarrow \mathbf{D}^-[W_{\mathbb{A}}^{-1}] = \mathbf{DM}_{Nis}^{\mathrm{eff}, -}(k, R)$$

is an equivalence of tensor triangulated categories.

PROOF. The category $\mathcal{L} = \mathcal{L}_{Nis}$ is a thick subcategory of \mathbf{D}^- . By 14.6.1 and 14.8, the functor $\mathcal{L} \rightarrow \mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ is fully faithful. By 14.4, every object K of $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ is isomorphic to $\mathrm{Tot}C_*(K)$, which is in \mathcal{L} by 14.9. Hence \mathcal{L} is equivalent to $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$.

It follows that \mathcal{L} is a tensor triangulated category, because $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$ is. If E and F are \mathbb{A}^1 -local, we have seen that the tensor product $E \otimes_{L, Nis}^{tr} F$ is naturally isomorphic to $E \otimes_{\mathcal{L}} F$ in $\mathbf{D}^-[W_{\mathbb{A}}^{-1}]$. That is, $\otimes_{\mathcal{L}}$ is isomorphic to the induced tensor operation on \mathcal{L} . \square

In [TriCa, p. 210], the tensor structure on \mathcal{L}_{Nis} was defined using $\otimes_{\mathcal{L}}$.

REMARK 14.12. If X is smooth and F is a Nisnevich sheaf with transfers, we define $\underline{RHom}(R_{tr}X, F)$ to be the complex $\mathrm{RHom}(- \times X, F)$ of sheaves with transfers, as in 8.21. If k is perfect, this complex is bounded above by 13.18. The $\underline{RHom}(R_{tr}X, F)$ construction extends in an evident way to the more general situation when F is replaced by a bounded above complex L , and $R_{tr}X$ is replaced by a complex representing an effective geometric motive M . Moreover, if L is an \mathbb{A}^1 -local complex, then $\underline{RHom}(M, L)$ is also \mathbb{A}^1 -local, by 14.9.

If K is another bounded above complex, then a short calculation shows that in either the derived category $\mathbf{D}^-(\mathrm{Sh}(\mathrm{Cor}_k, R))$ or in $\mathbf{DM}_{Nis}^{\mathrm{eff}, -}(k, R)$ we have the adjunction (where \otimes is either $\otimes_{L, Nis}^{tr}$ or $\otimes_{\mathcal{L}}$):

$$\mathrm{Hom}(K \otimes_{R_{tr}}(X), L) \cong \mathrm{Hom}(K, \underline{RHom}(R_{tr}X, L)).$$

EXERCISE 14.13. (a) Show that $\underline{RHom}(R_{tr}X, L[n]) \cong \underline{RHom}(R_{tr}X, L)[n]$.

(b) Use 14.16 and 15.2 to show that $\underline{RHom}(X, R) \cong R$ and $\underline{RHom}(X(r), R) = 0$ for all smooth X and $r > 0$.

Next, recall from 9.8 that two parallel morphisms f and g of sheaves are said to be \mathbb{A}^1 -**homotopic** if there is a map $F \otimes_L^{tr} \mathbb{Z}_{tr}(\mathbb{A}^1) \rightarrow G$ whose restrictions along 0 and 1 coincide with f and g , respectively. The proof of 9.10 shows that \mathbb{A}^1 -homotopic morphisms between Nisnevich sheaves become equal in $\mathbf{DM}_{Nis}^{\mathrm{eff}, -}(k, R)$.

PROPOSITION 14.14. *Let C and D be bounded above complexes of Nisnevich sheaves with transfers, whose cohomology sheaves are homotopy invariant. If C and D are \mathbb{A}^1 -local, then \mathbb{A}^1 -homotopic maps $f, g : C \rightarrow D$ induce the same maps on hypercohomology:*

$$f = g : \mathbb{H}_{Zar}^*(X, C) \rightarrow \mathbb{H}_{Zar}^*(X, D).$$

PROOF. To prove the proposition, we write \mathbf{DM} for $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}(k)$. Combining 13.10 with 13.5, we see that

$$\mathbb{H}_{\text{Zar}}^n(X, C) \cong \text{Hom}_{\mathbf{D}^-}(\mathbb{Z}_{tr}(X), C[n]).$$

If C is \mathbb{A}^1 -local, this equals $\text{Hom}_{\mathbf{DM}}(\mathbb{Z}_{tr}(X), C[n])$ by 14.6.1. Since f and g agree in \mathbf{DM} , they induce the same map from $\mathbb{H}^n(X, C) \cong \text{Hom}_{\mathbf{DM}}(\mathbb{Z}_{tr}(X), C[n])$ to $\mathbb{H}^n(X, D) \cong \text{Hom}_{\mathbf{DM}}(\mathbb{Z}_{tr}(X), D[n])$, as asserted. \square

We will need the following elementary result for $R = \mathbb{Q}$ in 14.30 below. It is proven by replacing $\mathbb{Z}(X)$ by $R_{tr}(X)$ in the proof of 12.13.

LEMMA 14.15. *The smallest class in \mathbf{D}^- which contains all the $R_{tr}(X)$ and is closed under quasi-isomorphisms, direct sums, shifts, and cones is all of \mathbf{D}^- .*

One of the features of motivic cohomology is that it is representable by the \mathbb{A}^1 -local complexes $R(i)[n]$.

PROPOSITION 14.16. *Let L be \mathbb{A}^1 -local. Then for any $X \in \text{Sm}/k$*

$$\mathbb{H}_{\text{Zar}}^n(X, L) \cong \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(R_{tr}(X), L[n]).$$

In particular, the motivic cohomology functors $X \mapsto H^{n,i}(X, R)$ are representable on Sm/k by $R(i)[n]$ in $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}(k, R)$:

$$H^{n,i}(X, R) \cong \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(R_{tr}(X), R(i)[n]).$$

PROOF. By 14.8 and 13.10, the left hand side is $\mathbb{H}_{\text{Nis}}^n(X, L)$. By 14.6.1, the right side equals $\text{Hom}_{\mathbf{D}^-}(R_{tr}(X), L[n])$. These are isomorphic by 13.5. The final representability formula follows from this and 13.11, because $R(i)[n]$ is \mathbb{A}^1 -local by 3.1 and 14.9. \square

DEFINITION 14.17. Let X be any scheme of finite type over k and $i \geq 0$. We define the motivic cohomology of X with coefficients in R to be:

$$H^{n,i}(X, R) = \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(R_{tr}(X), R(i)[n]),$$

where $R_{tr}(X)$ was defined in 2.11. This agrees with our original definition 3.4 for smooth X by 14.16.

Dually, we define the motivic homology $H_{n,i}(X, R)$ to be

$$H_{n,i}(X, R) = \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(R(i)[n], R_{tr}(X)).$$

Suslin's algebraic singular homology, defined in 10.8, is the special case $i = 0$ of motivic homology.

PROPOSITION 14.18. *If X is any scheme of finite type over k , then*

$$H_n^{\text{sing}}(X, R) \cong H_{n,0}(X, R).$$

PROOF. Because $H_{Nis}^0(\mathrm{Spec} k, -)$ is an exact functor, we have $\mathbb{H}_{Nis}^*(\mathrm{Spec} k, K) = H^*(K(\mathrm{Spec} k))$ for every complex K . Setting $R = R_{tr}(\mathrm{Spec} k)$, we compute:

$$\begin{aligned}
 H_n^{sing}(X, R) &= H_n(C_*R_{tr}(X)(\mathrm{Spec} k)) && \text{by definition 10.8,} \\
 &= \mathbb{H}_{Nis}^{-n}(\mathrm{Spec} k, C_*R_{tr}(X)) && \text{by remark above,} \\
 &= \mathrm{Hom}_{\mathbf{D}^-}(R[n], C_*R_{tr}(X)) && \text{by 13.5,} \\
 &= \mathrm{Hom}_{\mathbf{DM}_{Nis}^{\mathrm{eff}, -}(k, R)}(R[n], C_*R_{tr}(X)) && \text{by 14.9 and (14.6.1),} \\
 &= H_{n,0}(X, R) && \text{by definition. } \square
 \end{aligned}$$

If we assume that k admits resolution of singularities, we may use the cdh topology to extend some of the previous results from smooth schemes to all schemes of finite type. For example, applying $\mathrm{Hom}(R(i)[n], -)$ to the triangle (14.5.3) yields:

PROPOSITION 14.19. *Assume that resolution of singularities holds over k . Let $X' \rightarrow X$ be an abstract blow-up with center Z , and set $Z' = Z \times_X X'$. Then there is a long exact sequence in motivic homology:*

$$H_{n,i}(Z', R) \rightarrow H_{n,i}(Z, R) \oplus H_{n,i}(X', R) \rightarrow H_{n,i}(X, R) \rightarrow H_{n-1,i}(Z', R).$$

Now if L is \mathbb{A}^1 -local, we can identify $\mathrm{Hom}_{\mathbf{D}^-}(K, L)$ with $\mathrm{Hom}_{\mathbf{DM}_{Nis}^{\mathrm{eff}, -}}(K, L)$ by (14.6.1). When $K = \mathbb{Z}_{tr}(X)$, we can identify these Hom groups with cdh hypercohomology. Note that cdh hypercohomology makes sense by 12.24, because the cdh cohomological dimension of any Noetherian scheme is finite.

THEOREM 14.20. *Assume that resolution of singularities holds over k . Let L be an \mathbb{A}^1 -local complex. Then for any X in Sch/k and all $n \geq 0$ we have:*

$$\mathrm{Hom}_{\mathbf{D}^-}(\mathbb{Z}_{tr}(X), L[n]) \cong \mathbb{H}_{cdh}^n(X, L_{cdh}).$$

In particular, for any scheme X , and all positive n and i :

$$H^{n,i}(X, R) \cong \mathbb{H}_{cdh}^n(X, R(i)_{cdh}).$$

PROOF. For any X , the inclusion of Nisnevich sheaves $\iota : \mathbb{Z}(X) \rightarrow \mathbb{Z}_{tr}(X)$ induces a sequence of maps:

$$\begin{aligned}
 \mathrm{Hom}_{\mathbf{D}^-}(\mathbb{Z}_{tr}(X), L[n]) &= \mathrm{Ext}_{\mathbf{D}^-}^n(\mathbb{Z}_{tr}(X), L) && \text{by definition,} \\
 &\xrightarrow{u} \mathrm{Ext}_{Nis}^n(\mathbb{Z}_{tr}(X), L) && \text{by forgetting transfers,} \\
 &\xrightarrow{\iota^*} \mathrm{Ext}_{Nis}^n(\mathbb{Z}(X), L) && \text{by } \iota, \\
 &\xrightarrow{r} \mathrm{Ext}_{cdh}^n((\mathbb{Z}(X))_{cdh}, L_{cdh}) && \text{by change of sites,} \\
 &= \mathbb{H}_{cdh}^n(X, L_{cdh}) && \text{by definition.}
 \end{aligned}$$

If X is smooth, the composite $\iota^* \circ u$ is an isomorphism by 13.5 and 12.12. By mimicking the proof of 13.10 (using 13.27 in place of 13.9), we see that the map r is an isomorphism too. This finishes the proof when X is smooth.

When X is any scheme, we proceed by induction on $\dim X$. We may assume that X is reduced. By resolution of singularities, there is a smooth X' and a proper birational morphism $p : X' \rightarrow X$. Let Z be a proper subscheme (of lower dimension) such that p is an isomorphism over $X - Z$. Note that the cdh sheafification of the sequence 12.22 is exact for the cdh cover $X' \amalg Z \rightarrow X$. By 13.22, we have a long exact sequence of cdh hypercohomology groups. There is also a corresponding long exact sequence of hyperext groups obtained by applying $\mathrm{Hom}(-, L)$ to the triangle (14.5.3). Now consider the following morphism of long exact sequences:

$$\begin{array}{ccccc} \mathrm{Ext}^{n-1}(\mathbb{Z}_{tr}(Z'), L) & \longrightarrow & \mathrm{Ext}^n(\mathbb{Z}_{tr}(X), L) & \longrightarrow & \mathrm{Ext}^n(\mathbb{Z}_{tr}(X'), L) \oplus \mathrm{Ext}^n(\mathbb{Z}_{tr}(Z), L) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{H}_{cdh}^{n-1}(Z', L_{cdh}) & \longrightarrow & \mathbb{H}_{cdh}^n(X, L_{cdh}) & \longrightarrow & \mathbb{H}_{cdh}^n(X', L_{cdh}) \oplus \mathbb{H}_{cdh}^n(Z, L_{cdh}). \end{array}$$

The outside verticals are isomorphisms by induction and the smooth case. We conclude the proof of the general result using the 5-lemma.

Now the complex $R(i)$ is \mathbb{A}^1 -local by 3.1 and 14.9, so by 14.17 and (14.6.1) $H^{n,i}(X, R) \cong \mathrm{Hom}_{\mathbf{D}^-}(R_{tr}(X), R(i)[n])$. This yields the final assertion. \square

Motives with \mathbb{Q} -coefficients

We now consider the case when the coefficient ring R contains \mathbb{Q} . Our first goal is to identify étale and Nisnevich motivic cohomology (14.24). We will then describe $\mathbf{DM}_{\acute{e}t}^{\mathrm{eff},-}(k, R)$ (in 14.28), and finally show that $\mathbf{DM}_{Nis}^{\mathrm{eff},-}(k, R) \cong \mathbf{DM}_{\acute{e}t}^{\mathrm{eff},-}(k, R)$ (in 14.30).

LEMMA 14.21. *Let F be a Zariski sheaf of \mathbb{Q} -modules with transfers. Then F is also an étale sheaf with transfers.*

PROOF. It suffices to show that the presheaf kernel and cokernel of $F \rightarrow F_{\acute{e}t}$ vanish. By 6.17, these are presheaves with transfers. Thus we may suppose that $F_{\acute{e}t} = 0$. If $F \neq 0$ then there is a point $x \in X$ and a nonzero element $c \in F(S)$, $S = \mathrm{Spec} \mathcal{O}_{X,x}$. Since $F_{\acute{e}t} = 0$, there is a finite étale map $S' \rightarrow S$ with $c|_{S'} = 0$. As in 1.11, the composition of the transfers and inclusion

$$F(S) \rightarrow F(S') \rightarrow F(S)$$

is multiplication by d , the degree of $S' \rightarrow S$. Hence this composition is an isomorphism. Since it sends c to zero, we have $c = 0$. This contradiction shows that $F = 0$, as desired. \square

COROLLARY 14.22. *If F is a presheaf of \mathbb{Q} -modules with transfers, then $F_{Nis} = F_{\acute{e}t}$.*

PROOF. By 13.1, F_{Nis} is a sheaf with transfers, so 14.21 applies. \square

PROPOSITION 14.23. *If F is an étale sheaf of \mathbb{Q} -modules, then*

$$H_{\acute{e}t}^n(-, F) = H_{Nis}^n(-, F).$$

PROOF. We need to prove that $H_{\acute{e}t}^n(S, F) = 0$ for $n > 0$ when S is Hensel local. Given this, the result will follow from the Leray spectral sequence. Since F is uniquely divisible, so is its cohomology. But if s is the closed point of S then $H_{\acute{e}t}^n(S, F)$ equals $H_{\acute{e}t}^n(s, F)$, which is a torsion group. Hence $H_{\acute{e}t}^n(S, F) = 0$. \square

Recall from 10.1 that the étale (or Lichtenbaum) motivic cohomology $H_L^{p,q}(X, \mathbb{Q})$ is defined to be the étale hypercohomology of the complex $\mathbb{Q}(q)$.

THEOREM 14.24. *Let k be a perfect field. If K is a bounded above complex of presheaves of \mathbb{Q} -modules with transfers, then $K_{Nis} = K_{\acute{e}t}$ and*

$$\mathbb{H}_{\acute{e}t}^*(X, K_{\acute{e}t}) = \mathbb{H}_{Nis}^*(X, K_{Nis})$$

for every X in Sm/k . In particular, $H_L^{p,q}(X, \mathbb{Q}) = H^{p,q}(X, \mathbb{Q})$.

PROOF. Consider $F = H^q C$. By 14.22, $K_{Nis} = K_{\acute{e}t}$ and $F_{Nis} = F_{\acute{e}t}$. By 14.23, we have isomorphisms $H_{\acute{e}t}^p(X, F_{\acute{e}t}) \rightarrow H_{Nis}^p(X, F_{Nis})$, and these groups vanish for $p > \dim(X)$ by 14.23. Comparing the hypercohomology spectral sequences for the Nisnevich and the étale topology yields the result.

In particular, the result applies to the complex $C = \mathbb{Q}(q)$. \square

For clarity, let us say that a complex K is **étale \mathbb{A}^1 -local** if it is \mathbb{A}^1 -local for the étale topology (as in 9.17), and **Nisnevich \mathbb{A}^1 -local** if it is \mathbb{A}^1 -local for the Nisnevich topology.

These notions coincide for any étale sheaf of R -modules with transfers F , where $\mathbb{Q} \subseteq R$. To see this, note that by 14.8, F is Nisnevich \mathbb{A}^1 -local if and only if it is homotopy invariant: $F(X) \cong F(X \times \mathbb{A}^1, F)$ for all smooth X . On the other hand, we see from 9.24 that F is étale \mathbb{A}^1 -local if and only if it is strictly \mathbb{A}^1 -homotopy invariant in the sense of 9.22: $H_{\acute{e}t}^n(X, F) \cong H_{\acute{e}t}^n(X \times \mathbb{A}^1, F)$ for all smooth X and all n . Hence if F is étale \mathbb{A}^1 -local then it is Nisnevich \mathbb{A}^1 -local. Conversely, if F is Nisnevich \mathbb{A}^1 -local then $H_{Nis}^n(X, F) \cong H_{Nis}^n(X \times \mathbb{A}^1, F)$ for all smooth X by 13.8. By 14.23, F is strictly \mathbb{A}^1 -homotopy invariant, i.e., étale \mathbb{A}^1 -local. This proves the following lemma:

LEMMA 14.25. *Let k be a perfect field, and $\mathbb{Q} \subseteq R$. The following are equivalent for every étale sheaf of R -modules with transfers F : F is homotopy invariant; F is Nisnevich \mathbb{A}^1 -local; and F is étale \mathbb{A}^1 -local.*

PROPOSITION 14.26. *Let k be a perfect field and suppose that $\mathbb{Q} \subseteq R$. If K is a bounded above cochain complex of étale sheaves of R -modules with transfers, the following are equivalent: K is étale \mathbb{A}^1 -local; K is Nisnevich \mathbb{A}^1 -local; and the sheaves $a_{\acute{e}t}(H^n K)$ are homotopy invariant.*

In particular, each $R(j)$ is an étale \mathbb{A}^1 -local complex.

PROOF. By 14.22, $a_{Nis}(H^n K) = a_{\acute{e}t}(H^n K)$ so, by 14.8, K is Nisnevich \mathbb{A}^1 -local if and only if the sheaves $a_{\acute{e}t}(H^n K)$ are homotopy invariant.

Suppose that the sheaves $a_{\acute{e}t} H^n(K)$ are homotopy invariant. By 14.25, they are étale \mathbb{A}^1 -local. Since $\mathbb{Q} \subseteq R$, we have $\text{cd}_R(k) = 0$, so K is étale \mathbb{A}^1 -local by 9.30.

Conversely, suppose that K is étale \mathbb{A}^1 -local and set $K' = \text{Tot}C_*(K)$. By 2.19, each $H^n(K')$ is \mathbb{A}^1 -homotopy invariant. By theorem 13.8 and 14.21, each étale sheaf $a_{\acute{e}t}(H^n K')$ is homotopy invariant. The first part of this proof shows that K' is étale \mathbb{A}^1 -local. By 9.15, $K \rightarrow K'$ is an (étale) \mathbb{A}^1 -weak equivalence, and an isomorphism in $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$. By 9.21, $K \rightarrow K'$ is an isomorphism in $\mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$. Hence each sheaf $a_{\acute{e}t}(H^n K)$ is isomorphic to $a_{\acute{e}t}(H^n K')$, and is therefore also homotopy invariant. \square

COROLLARY 14.27. *If k is perfect and $\mathbb{Q} \subseteq R$, the étale motivic cohomology functors $X \mapsto H_L^{n,i}(X, R)$ are representable by $R(i)[n]$ in $\mathbf{DM}_{\acute{e}t}^- = \mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$:*

$$H_L^{n,i}(X, R) \cong \text{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(R_{tr}(X), R(i)[n]).$$

PROOF. Write $\mathbf{D}_{\acute{e}t}^-$ for $\mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$. Since $R(i)$ is étale \mathbb{A}^1 -local by 14.26, we know from 9.19 that

$$\text{Hom}_{\mathbf{DM}_{\acute{e}t}^-}(R_{tr}(X), R(i)[n]) = \text{Hom}_{\mathbf{D}_{\acute{e}t}^-}(R_{tr}(X), R(i)[n]) = \text{Ext}^n(R_{tr}(X), R(i)).$$

By 6.25, this Ext group is $H_L^{n,i}(X, R) = \mathbb{H}_{\acute{e}t}^n(X, R(i))$. \square

Let $\mathcal{L}_{\acute{e}t}$ denote the full subcategory of $\mathbf{D}_{\acute{e}t}^- = \mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$ consisting of complexes with homotopy invariant cohomology sheaves. By 14.26, $\mathcal{L}_{\acute{e}t}$ is also the subcategory of étale \mathbb{A}^1 -local complexes.

THEOREM 14.28. *The natural functor $\mathcal{L}_{\acute{e}t} \rightarrow \mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ is an equivalence of triangulated categories if $\mathbb{Q} \subseteq R$.*

PROOF. The functor is full and faithful by 9.19 and 14.26. Since every K in $\mathbf{D}_{\acute{e}t}^-$ becomes isomorphic to $\text{Tot}C_*(K)$ in $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}$ by 9.15, and $\text{Tot}C_*(K)$ is in $\mathcal{L}_{\acute{e}t}$ by 2.19 and 14.26, the functor is an equivalence. \square

REMARK 14.29. Theorem 14.28 implies that $\mathcal{L}_{\acute{e}t}$ is a tensor triangulated category. As in the proof of 9.35 and 14.11, 14.4 and 14.9 show that the tensor operation of $\mathcal{L}_{\acute{e}t}$ is isomorphic to the operation $\otimes_{\mathcal{L}}$ defined in 9.34.

THEOREM 14.30. *If $\mathbb{Q} \subseteq R$, then $\sigma : \mathbf{DM}_{Nis}^{\text{eff},-}(k, R) \rightarrow \mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ is an equivalence of tensor triangulated categories.*

PROOF. Recall from 14.3 that there are tensor triangulated functors $\mathbf{D}^- \rightarrow \mathbf{D}_{\acute{e}t}^-$ and $\mathbf{DM}_{Nis}^{\text{eff},-} \rightarrow \mathbf{DM}_{\acute{e}t}^{\text{eff},-}$. Since they are onto on objects, it suffices to show that the functor σ is full and faithful, i.e., that we have $\text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff},-}}(K, L) \cong \text{Hom}_{\mathbf{DM}_{\acute{e}t}^{\text{eff},-}}(K_{\acute{e}t}, L_{\acute{e}t})$. By theorem 14.11, we may assume that L is in \mathcal{L}_{Nis} . The class of objects K so that $\text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff},-}}(K, L[n]) \cong \text{Hom}_{\mathbf{DM}_{\acute{e}t}^{\text{eff},-}}(K_{\acute{e}t}, L_{\acute{e}t}[n])$ for all n is closed under quasi-isomorphisms, direct sums, shifts, and cones. By 14.15, it suffices to show that each $R_{tr}(X)$ is in this class. But then by 14.6.1, 13.5, 9.19, and 6.25, we have $\text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff},-}}(R_{tr}(X), L[n]) \cong \text{Hom}_{\mathbf{D}^-}(R_{tr}(X), L[n]) \cong \mathbb{H}_{Nis}^n(X, L)$ and $\text{Hom}_{\mathbf{DM}_{\acute{e}t}^{\text{eff},-}}(R_{tr}(X), L_{\acute{e}t}[n]) \cong \text{Hom}_{\mathbf{D}_{\acute{e}t}^-}(R_{tr}(X), L_{\acute{e}t}[n]) \cong \mathbb{H}_{\acute{e}t}^n(X, L_{\acute{e}t})$. By theorem 14.24, these groups are isomorphic as required. \square

LECTURE 15

The complex $\mathbb{Z}(n)$ and \mathbb{P}^n

The goal of this lecture is to interpret the motivic complex $\mathbb{Z}(n)$ in terms of $\mathbb{Z}_{tr}(\mathbb{P}^n)$ and use this to show that the product on motivic cohomology is graded-commutative. We also apply this to give a projective bundle theorem and a Gysin map associated to a smooth blow-up. We begin by observing that $M(\mathbb{P}^n - 0) \cong M(\mathbb{P}^{n-1})$.

LEMMA 15.1. *There is a chain homotopy equivalence:*

$$C_*\mathbb{Z}_{tr}(\mathbb{P}^n - 0) \simeq C_*\mathbb{Z}_{tr}(\mathbb{P}^{n-1}).$$

PROOF. Consider the projection $(\mathbb{P}^n - 0) \rightarrow \mathbb{P}^{n-1}$ which sends $(x_0 : \cdots : x_n)$ to $(x_1 : \cdots : x_n)$, where 0 is $(1 : 0 : \cdots : 0)$. This map has affine fibers. The self homotopy $\lambda(x_0 : \cdots : x_n) \rightarrow (\lambda x_0 : x_1 : \cdots : x_n)$ is well defined on $\mathbb{P}^n - 0 \times \mathbb{A}^1$, even for $\lambda = 0$, because one of x_1, \dots, x_n is always non zero. Hence the projection and the section $(x_1 : \cdots : x_n) \mapsto (0 : x_1 : \cdots : x_n)$ are inverse \mathbb{A}^1 -homotopy equivalences. The lemma now follows from 2.26. \square

THEOREM 15.2. *If k is a perfect field, there is a quasi-isomorphism of Zariski sheaves for each n :*

$$C_* (\mathbb{Z}_{tr}(\mathbb{P}^n) / \mathbb{Z}_{tr}(\mathbb{P}^{n-1})) \simeq C_*\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})[n] = \mathbb{Z}(n)[2n].$$

In particular, $C_ (\mathbb{Z}_{tr}(\mathbb{P}^n) / \mathbb{Z}_{tr}(\mathbb{P}^{n-1})) (X) \simeq \mathbb{Z}(n)[2n](X)$ for any smooth local X .*

Our proof will use theorem 13.12, whose proof depended upon theorem 13.8, a result whose proof we have postponed until lecture 24. The requirement that k be perfect is only needed for 13.8 (and hence 13.12).

PROOF. Let \mathcal{U} be the usual cover of \mathbb{P}^n by $(n+1)$ copies of \mathbb{A}^n and note that n of these form a cover \mathcal{V} of $\mathbb{P}^n - 0$. The intersection of $i+1$ of these \mathbb{A}^n is $\mathbb{A}^{n-i} \times (\mathbb{A}^1 - 0)^i$. By 6.14, we have quasi-isomorphisms $\mathbb{Z}_{tr}(\mathcal{U}) \rightarrow \mathbb{Z}_{tr}(\mathbb{P}^n)$ and $\mathbb{Z}_{tr}(\mathcal{V}) \rightarrow \mathbb{Z}_{tr}(\mathbb{P}^n - 0)$ of complexes of Nisnevich sheaves with transfers. Hence the quotient complex $Q_* = \mathbb{Z}_{tr}(\mathcal{U}) / \mathbb{Z}_{tr}(\mathcal{V})$ is a resolution of $\mathbb{Z}_{tr}(\mathbb{P}^n) / \mathbb{Z}_{tr}(\mathbb{P}^n - 0)$ as a Nisnevich sheaf. By 13.14 and 15.1, or by 13.15, $\text{Tot} C_* Q_*$ is quasi-isomorphic to $C_* (\mathbb{Z}_{tr}(\mathbb{P}^n) / \mathbb{Z}_{tr}(\mathbb{P}^n - 0))$ and hence to $C_* (\mathbb{Z}_{tr}(\mathbb{P}^n) / \mathbb{Z}_{tr}(\mathbb{P}^{n-1}))$ for the Zariski topology.

On the other hand, we know from 2.13 that for $T = \mathbb{A}^1 - 0$ the sequence

$$0 \rightarrow \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n}) \rightarrow \mathbb{Z}_{tr}(T^n) \rightarrow \oplus_i \mathbb{Z}_{tr}(T^{n-1}) \rightarrow \oplus_{i,j} \mathbb{Z}_{tr}(T^{n-2}) \rightarrow \dots \rightarrow \oplus_{i,j} \mathbb{Z}_{tr}(T^2) \rightarrow \oplus_i \mathbb{Z}_{tr}(T) \rightarrow \mathbb{Z} \rightarrow 0$$

is split exact. Rewriting this as $0 \rightarrow \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n}) \rightarrow R_n \rightarrow R_{n-1} \rightarrow \dots \rightarrow R_0 \rightarrow 0$, with $R_n = \mathbb{Z}_{tr}(T^n)$, $R_{n-1} = \oplus_i \mathbb{Z}_{tr}(T^{n-1})$, and $R_0 = \mathbb{Z}$, we may regard it as a chain homotopy equivalence $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})[n] \rightarrow R_*$. With this indexing there is a natural map $Q_* \rightarrow R_*$ whose typical term is a direct sum of the projections

$$\mathbb{Z}_{tr}(\mathbb{A}^{n-i} \times T^i) \rightarrow \mathbb{Z}_{tr}(T^i).$$

These are \mathbb{A}^1 -homotopy equivalences (see 2.25). Applying C_* turns them into quasi-isomorphisms by 2.26. Hence we have quasi-isomorphisms of total complexes of presheaves with transfers

$$\text{Tot } C_* Q_* \xrightarrow{\simeq} \text{Tot } C_* R_* \xleftarrow{\simeq} C_* \mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})[n].$$

Combining with $\text{Tot } C_* Q_* \simeq C_*(\mathbb{Z}_{tr}(\mathbb{P}^n)/\mathbb{Z}_{tr}(\mathbb{P}^{n-1}))$ yields the result in the Zariski topology. \square

If $n = 1$, it is easy to see that the isomorphisms of 13.17 and 15.2 agree. Figure 15.1 illustrates the proof of theorem 15.2 when $n = 2$. We have written ‘X’ for $C_* \mathbb{Z}_{tr}(X)$ in order to save space, and ‘ \mathbb{A}^1 -h.e.’ for \mathbb{A}^1 -homotopy equivalence.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & 0 & \longrightarrow & \mathbb{A}^1 \times (\mathbb{A}^1 - 0) & \longrightarrow & 2\mathbb{A}^2 & \longrightarrow & \mathbb{P}^2 - 0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & (\mathbb{A}^1 - 0)^2 & \longrightarrow & 3(\mathbb{A}^1 \times (\mathbb{A}^1 - 0)) & \longrightarrow & 3\mathbb{A}^2 & \longrightarrow & \mathbb{P}^2 \\
 \downarrow & & \downarrow = & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & (\mathbb{A}^1 - 0)^2 & \longrightarrow & 2(\mathbb{A}^1 \times (\mathbb{A}^1 - 0)) & \longrightarrow & \mathbb{A}^2 & \longrightarrow & \mathbb{P}^2/(\mathbb{P}^2 - 0) \\
 \downarrow & & \downarrow = & & \downarrow \mathbb{A}^1\text{-h.e.} & & \downarrow \mathbb{A}^1\text{-h.e.} & & \downarrow \\
 \mathbb{G}_m^{\wedge 2} & \longrightarrow & (\mathbb{A}^1 - 0)^2 & \longrightarrow & 2(\mathbb{A}^1 - 0) & \longrightarrow & \text{pt} & \longrightarrow & 0
 \end{array}$$

FIGURE 15.1. The case $n = 2$ of theorem 15.2

COROLLARY 15.3. *For each n there is a quasi-isomorphism for the Zariski topology*

$$C_*(\mathbb{Z}_{tr}(\mathbb{A}^n - 0)/\mathbb{Z}) \simeq \mathbb{Z}(n)[2n - 1].$$

PROOF. Applying 13.15 and 15.1 to the cover of \mathbb{P}^n by \mathbb{A}^n and $\mathbb{P}^n - 0$, we see that the sequence

$$0 \rightarrow C_* \mathbb{Z}_{tr}(\mathbb{A}^n - 0) \rightarrow C_* \mathbb{Z}_{tr}(\mathbb{A}^n) \oplus C_* \mathbb{Z}_{tr}(\mathbb{P}^{n-1}) \rightarrow C_* \mathbb{Z}_{tr}(\mathbb{P}^n) \rightarrow 0$$

becomes exact for the Zariski topology. The result now follows from theorem 15.2, since $C_*\mathbb{Z}_{tr}(\mathbb{A}^n) \simeq C_*\mathbb{Z}_{tr}(\text{Spec } k) \simeq \mathbb{Z}$ by 2.24 and 2.14. \square

EXERCISE 15.4. Show that the map $C_*\mathbb{Z}_{tr}(\mathbb{P}^i) \rightarrow \mathbb{Z}(i)[2i]$ of theorem 15.2 factors through the natural inclusion $C_*\mathbb{Z}_{tr}(\mathbb{P}^i) \rightarrow C_*\mathbb{Z}_{tr}(\mathbb{P}^n)$ for all $n > i$. *Hint:* Fix U_0 in \mathcal{U} and construct $\mathbb{Z}_{tr}(\check{\mathcal{U}}) \rightarrow \mathbb{Z}(1)[2]$ vanishing on $\mathbb{Z}_{tr}(U_0)$. Then form

$$\mathbb{Z}_{tr}(\check{\mathcal{U}}) \xrightarrow{\Delta} \mathbb{Z}_{tr}(\check{\mathcal{U}}) \otimes \cdots \otimes \mathbb{Z}_{tr}(\check{\mathcal{U}}) \rightarrow \mathbb{Z}(1)[2]^{\otimes i} \rightarrow \mathbb{Z}(i)[2i].$$

COROLLARY 15.5. *There is a quasi-isomorphism*

$$M(\mathbb{P}^n) = C_*\mathbb{Z}_{tr}(\mathbb{P}^n) \xrightarrow{\simeq} \mathbb{Z} \oplus \mathbb{Z}(1)[2] \oplus \cdots \oplus \mathbb{Z}(n)[2n].$$

PROOF. We proceed by induction, the case $n = 1$ being 13.17. By exercise 15.4, the map $\mathbb{Z}_{tr}(\mathbb{P}^{n-1}) \rightarrow \mathbb{Z}_{tr}(\mathbb{P}^n)$ is split injective in **DM**, because the quasi-isomorphism $\mathbb{Z}_{tr}(\mathbb{P}^{n-1}) \rightarrow \bigoplus \mathbb{Z}(i)[i]$ factors through it. Hence the distinguished triangle of theorem 15.2 splits:

$$C_*\mathbb{Z}_{tr}(\mathbb{P}^{n-1}) \longrightarrow C_*\mathbb{Z}_{tr}(\mathbb{P}^n) \longrightarrow \mathbb{Z}(n)[n] \longrightarrow C_*\mathbb{Z}_{tr}(\mathbb{P}^{n-1})[1]. \quad \square$$

Our re-interpretation of the motivic complexes allows us to show that the product in motivic cohomology is skew-commutative. This will be a consequence of the following construction, and some linear algebra.

EXAMPLE 15.6. Consider the reflection automorphism τ of \mathbb{P}^n , $n \geq 1$, sending $(x_0 : x_1 : \cdots : x_n)$ to $(-x_0 : x_1 : \cdots : x_n)$. We claim that the induced automorphism of $C_*\mathbb{Z}_{tr}(\mathbb{P}^n)$ is \mathbb{A}^1 -homotopic to the identity map, so that it is the identity map in $\mathbf{DM}_{Nis}^{\text{eff}, -}$ (see 14.1 and 9.10).

To see this, consider the elementary correspondence from $\mathbb{P}^n \times \mathbb{A}^1$ (parametrized by x_0, \dots, x_n and t) to \mathbb{P}^n (parametrized by y_0, \dots, y_n) given by the subvariety Z of $\mathbb{P}^n \times \mathbb{A}^1 \times \mathbb{P}^n$ defined by the homogeneous equation(s)

$$y_i(x_0y_i + tx_iy_0) = (t^2 - 1)x_iy_0^2, \quad i = 1, \dots, n$$

together with $x_iy_j = x_jy_i$ for $1 \leq i, j \leq n$ if $n \geq 2$. (Exercise: check that this is an elementary correspondence!) The restrictions along $t = \pm 1$ yield two finite correspondences from \mathbb{P}^n to itself, whose difference is $id_{\mathbb{P}^n} - \tau$.

Restricted to $\mathbb{P}^{n-1} \times \mathbb{A}^1$, this correspondence is the projection onto \mathbb{P}^{n-1} . Thus it induces an \mathbb{A}^1 -homotopy between τ and the identity of $\mathbb{Z}_{tr}(\mathbb{P}^n)/\mathbb{Z}_{tr}(\mathbb{P}^{n-1})$. Applying C_* , we see from theorem 15.2 that it induces an \mathbb{A}^1 -homotopy between the reflection automorphism τ of $\mathbb{Z}(n)[2n]$ and the identity, so that τ is the identity map in $\mathbf{DM}_{Nis}^{\text{eff}, -}$.

The symmetric group Σ_n acts canonically on \mathbb{A}^n by permuting coordinates. By inspection, this induces a Σ_n -action on the sheaf with transfers $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge n})$ and hence on the motivic complexes $\mathbb{Z}(n)$.

PROPOSITION 15.7. *The action of the symmetric group Σ_n on $C_*\mathbb{Z}_{tr}(\mathbb{A}^n - 0)$ is \mathbb{A}^1 -homotopic to the trivial action.*

PROOF. Because the action is induced from an embedding $\Sigma_n \hookrightarrow GL_n(k)$, and every transposition acts as the reflection τ times an element of $SL_n(k)$, we see from example 15.6 that it suffices to show that the action of $SL_n(k)$ on $C_*\mathbb{Z}_{tr}(\mathbb{A}^n - 0)$ is chain homotopic to the trivial action.

Since every matrix in $SL_n(k)$ is a product of elementary matrices, it suffices to consider one elementary matrix $e_{ij}(a)$, $a \in k$. But multiplication by this matrix is \mathbb{A}^1 -homotopic to the identity of $\mathbb{A}^n - 0$, by the homotopy $(x, t) \mapsto e_{ij}(at)x$ (see 9.9). In particular it is an \mathbb{A}^1 -homotopy equivalence (see 2.25). By 2.26, the resulting endomorphism of $C_*\mathbb{Z}_{tr}(\mathbb{A}^n - 0)$ is chain homotopic to the identity. \square

COROLLARY 15.8. *The action of the symmetric group Σ_n on $\mathbb{Z}(n)$ is \mathbb{A}^1 -homotopic to the trivial action. Hence it is trivial in $\mathbf{DM}_{Nis}^{\text{eff}, -}$, and on the motivic cohomology groups $\mathbb{H}^p(X, \mathbb{Z}(n))$.*

Tensoring with a coefficient ring R does not affect the action, so it follows that Σ_n also acts trivially on $R(n)[2n]$, and on $\mathbb{H}^p(X, R(n))$.

PROOF. The action of Σ_n on \mathbb{A}^n extends to an action on \mathbb{P}^n fixing \mathbb{P}^{n-1} . In fact, all the constructions in the proof of theorem 15.2 and corollary 15.3 are equivariant. By 15.3, it suffices to show that the action of Σ_n on $C_*\mathbb{Z}_{tr}(\mathbb{A}^n - 0)$ is \mathbb{A}^1 -homotopic to the trivial action. This follows from 15.7 and 14.14. \square

Recall from 3.11 that there is a pairing of presheaves $\mathbb{Z}(i) \otimes \mathbb{Z}(j) \rightarrow \mathbb{Z}(i+j)$. By inspection of 3.10, this pairing is compatible with the action of the subgroup $\Sigma_i \times \Sigma_j$ of Σ_{i+j} , as well as with the permutation τ interchanging the first i and last j coordinates of $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge i+j})$.

THEOREM 15.9. *The pairing defined in 3.12 is skew-commutative:*

$$\mathbb{H}_{Zar}^p(X, \mathbb{Z}(i)) \otimes \mathbb{H}_{Zar}^q(X, \mathbb{Z}(j)) \rightarrow \mathbb{H}_{Zar}^{p+q}(X, \mathbb{Z}(i+j)).$$

PROOF. As in 8A.2, the permutation τ fits into the commutative diagram

$$\begin{array}{ccccc} \mathbb{H}^p(X, \mathbb{Z}(i)) \otimes \mathbb{H}^q(X, \mathbb{Z}(j)) & \longrightarrow & \mathbb{H}^{p+q}(X, \mathbb{Z}(i) \otimes \mathbb{Z}(j)) & \longrightarrow & \mathbb{H}^{p+q}(X, \mathbb{Z}(i+j)) \\ \downarrow (-1)^{pq} & & \downarrow \tau & & \downarrow \tau \\ \mathbb{H}^q(X, \mathbb{Z}(j)) \otimes \mathbb{H}^p(X, \mathbb{Z}(i)) & \longrightarrow & \mathbb{H}^{p+q}(X, \mathbb{Z}(j) \otimes \mathbb{Z}(i)) & \longrightarrow & \mathbb{H}^{p+q}(X, \mathbb{Z}(j+i)) \end{array}$$

and the right vertical map is the identity by corollary 15.8. \square

We conclude this lecture with a generalization of the decomposition 15.5 of $M(\mathbb{P}^n)$ to a projective bundle theorem.

CONSTRUCTION 15.10. Let $\mathbb{P} = \mathbb{P}(\mathcal{E}) \rightarrow X$ be a projective bundle associated to a vector bundle \mathcal{E} of rank $n+1$. From 4.2, 13.11, and 13.5 we have an isomorphism

$$\text{Pic}(\mathbb{P}) \cong H_{Nis}^2(\mathbb{P}, \mathbb{Z}(1)) \cong \text{Hom}_{\mathbf{D}^-}(\mathbb{Z}_{tr}(\mathbb{P}), \mathbb{Z}(1)[2]).$$

Therefore the canonical line bundle yields a canonical map $\tau : \mathbb{Z}_{tr}(\mathbb{P}) \rightarrow \mathbb{Z}(1)[2]$ in \mathbf{D}^- . Recall from 10.4 that there are multiplication maps for all $i \geq 1$, from $\mathbb{Z}(1)^{\otimes tr i} = \mathbb{Z}(1) \otimes^{tr} \cdots \otimes^{tr} \mathbb{Z}(1)$ to $\mathbb{Z}(i)$. For $i > 1$, we let τ^i denote the composite

$$\mathbb{Z}_{tr}(\mathbb{P}) \xrightarrow{\Delta} \mathbb{Z}_{tr}(\mathbb{P} \times \cdots \times \mathbb{P}) \xrightarrow{\cong} \mathbb{Z}_{tr}(\mathbb{P})^{\otimes tr i} \xrightarrow{\tau^{\otimes i}} \mathbb{Z}(1)[2]^{\otimes tr i} \longrightarrow \mathbb{Z}(i)[2i].$$

Finally, we extend the structure map $\sigma_0 : \mathbb{Z}_{tr}(\mathbb{P}) \rightarrow \mathbb{Z}_{tr}(X)$ to a canonical family of maps in \mathbf{D}^-

$$\sigma_i : \mathbb{Z}_{tr}(\mathbb{P}) \xrightarrow{\Delta} \mathbb{Z}_{tr}(\mathbb{P}) \otimes^{tr} \mathbb{Z}_{tr}(\mathbb{P}) \xrightarrow{\sigma_0 \otimes \tau^i} \mathbb{Z}_{tr}(X) \otimes \mathbb{Z}(i)[2i].$$

EXERCISE 15.11. Show that the canonical map in 15.10 induces the isomorphism $\mathbb{Z}_{tr}(\mathbb{P}^n_k) \cong \bigoplus_{i=0}^n \mathbb{Z}(i)[2i]$ of 15.5. *Hint:* Use exercise 15.4.

THEOREM 15.12 (Projective Bundle Theorem). *Let $p : \mathbb{P}(\mathcal{E}) \rightarrow X$ be a projective bundle associated to a vector bundle \mathcal{E} of rank $n + 1$. Then the canonical map*

$$\bigoplus_{i=0}^n \mathbb{Z}_{tr}(X)(i)[2i] \rightarrow \mathbb{Z}_{tr}(\mathbb{P}(\mathcal{E}))$$

is an isomorphism in \mathbf{DM} , and p is the projection onto the factor $\mathbb{Z}_{tr}(X)$.

PROOF. Using induction on the number of open subsets in a trivialization of \mathcal{E} , together with the Mayer-Vietoris triangles (14.5.1), we are reduced to the case when $\mathbb{P}(\mathcal{E}) = X \times \mathbb{P}^n$. Since $\mathbb{Z}_{tr}(X \times \mathbb{P}^n) \cong \mathbb{Z}_{tr}(X) \otimes^{tr} \mathbb{Z}_{tr}(\mathbb{P}^n)$, we may even assume $X = \text{Spec}(k)$. This case is given by 15.5 and exercise 15.11. \square

COROLLARY 15.13. *Let X be a smooth scheme and Z a smooth subscheme of pure codimension c . Let $p : X' \rightarrow X$ be the blow-up along Z . Then*

$$C_*\mathbb{Z}_{tr}(X') \cong C_*\mathbb{Z}_{tr}(X) \oplus \left(\bigoplus_{i=1}^{c-1} C_*\mathbb{Z}_{tr}(Z)(i)[2i] \right).$$

Moreover, there is a natural ‘‘Gysin’’ map $\gamma : C_\mathbb{Z}_{tr}(X) \rightarrow C_*\mathbb{Z}_{tr}(Z)(c)[2c]$, which is zero on $C_*\mathbb{Z}_{tr}(X - Z)$.*

PROOF. Since Z is smooth, Z' is the projective bundle associated to the normal bundle of Z in X . We claim that the morphism $C_*\mathbb{Z}_{tr}(X') \rightarrow C_*\mathbb{Z}_{tr}(X)$ is a split surjection. By 13.26 and 15.12, this will prove the first assertion.

Let X'' be the blow-up of $X \times \mathbb{A}^1$ along $Z \times 0$, and let $Z'' = (Z \times 0) \times_{X \times \mathbb{A}^1} X''$. Consider the following diagram, whose rows are distinguished triangles by 13.26:

$$\begin{array}{ccccccc} C_*\mathbb{Z}_{tr}(Z') & \xrightarrow{a} & C_*\mathbb{Z}_{tr}(Z) \oplus C_*\mathbb{Z}_{tr}(X') & \xrightarrow{b} & C_*\mathbb{Z}_{tr}(X) & \longrightarrow & C_*\mathbb{Z}_{tr}(Z')[1] \\ \downarrow f & & \downarrow g & & \simeq \downarrow h = 1_X \times 0 & & \downarrow \\ C_*\mathbb{Z}_{tr}(Z'') & \xrightarrow{c} & C_*\mathbb{Z}_{tr}(Z \times 0) \oplus C_*\mathbb{Z}_{tr}(X'') & \xrightarrow{d} & C_*\mathbb{Z}_{tr}(X \times \mathbb{A}^1) & \longrightarrow & C_*\mathbb{Z}_{tr}(Z'')[1]. \end{array}$$

By 2.24, the map h is a quasi-isomorphism. But h is also \mathbb{A}^1 -homotopic to $1_X \times 1$, so h lifts to a map $C_*\mathbb{Z}_{tr}(X) \rightarrow C_*\mathbb{Z}_{tr}(X'')$. This splits d , and hence c .

Let \mathcal{N} be the normal bundle of Z in X . Then the morphism $Z' \rightarrow Z''$ is the canonical embedding of $\mathbb{P}(\mathcal{N})$ into $\mathbb{P}(\mathcal{N} \oplus \mathcal{O})$. By 15.12, f is a splitting

monomorphism. Composing g with the splittings of c and f , we see that a splits naturally. Since the first row is a distinguished triangle, b also splits naturally, which implies the claim.

Note that the cokernel of f is $C_*\mathbb{Z}_{tr}(Z)(c)[2c]$. Composing the splitting of b , g , and the splitting of c with the projection onto this cokernel yields the desired Gysin map γ . A diagram chase shows that γ vanishes on $C_*\mathbb{Z}_{tr}(X - Z)$. \square

EXERCISE 15.14. The Gysin map for $X = \mathbb{A}^n$ or \mathbb{P}^n and $Z = 0$ induces a map

$$C_*(\mathbb{Z}_{tr}(\mathbb{A}^n)/\mathbb{Z}_{tr}(\mathbb{A}^n - 0)) \xrightarrow{\simeq} C_*(\mathbb{Z}_{tr}(\mathbb{P}^n)/\mathbb{Z}_{tr}(\mathbb{P}^n - 0)) \xrightarrow{\bar{\gamma}} \mathbb{Z}(n)[2n].$$

(The first map is a quasi-isomorphism by 12.20.) Show that $\bar{\gamma}$ agrees with the quasi-isomorphism of 15.2.

THEOREM 15.15. *Let X be a smooth scheme over a perfect field and Z a smooth closed subscheme of X of codimension c . Then there is a ‘‘Gysin’’ triangle:*

$$C_*\mathbb{Z}_{tr}(X - Z) \rightarrow C_*\mathbb{Z}_{tr}(X) \xrightarrow{\gamma} C_*\mathbb{Z}_{tr}(Z)(c)[2c] \rightarrow C_*\mathbb{Z}_{tr}(X - Z)[1].$$

PROOF. We have to show that the Gysin map $\bar{\gamma}: C_*(\mathbb{Z}_{tr}(X)/\mathbb{Z}_{tr}(X - Z)) \rightarrow C_*\mathbb{Z}_{tr}(Z)(c)[2c]$ of 15.13 is a quasi-isomorphism. By 12.20 we may assume that $X = Y \times \mathbb{A}^c$ and $Z = Y \times 0$. But then $\bar{\gamma}$ is the isomorphism of 15.14, tensored with $C_*\mathbb{Z}_{tr}(Y)$. \square

Equidimensional cycles

In the first part of this lecture we introduce the notion of an equidimensional cycle, and use it to construct the Suslin-Friedlander chain complex $\mathbb{Z}^{SF}(i)$. We then show (in 16.7) that $\mathbb{Z}^{SF}(i)$ is quasi-isomorphic to $\mathbb{Z}(i)$. In lecture 19 (19.4) we shall compare $\mathbb{Z}^{SF}(i)$ to the complex defining higher Chow groups. In the second part of this lecture, we use equidimensional cycles to define motives with compact support and investigate their basic properties. Finally we use Friedlander-Voevodsky Duality (see 16.24) to prove the Cancellation Theorem 16.25.

Let Z be a scheme of finite type over S such that every irreducible component of Z dominates a component of S . We say that Z is **equidimensional over S** of relative dimension m if for every point s of S , either Z_s is empty or each of its components have dimension m . If $S' \rightarrow S$ is any map, the pullback $S' \times_S Z$ is equidimensional over S' of relative dimension m .

DEFINITION 16.1. Let T be any scheme of finite type over k and $m \geq 0$ an integer. The presheaf $z_{equi}(T, r)$ on Sm/k is defined as follows. For each smooth S , $z_{equi}(T, r)(S)$ is the free abelian group generated by the closed and irreducible subvarieties Z of $S \times T$ which are dominant and equidimensional of relative dimension r over a component of S . If $S' \rightarrow S$ is any map, the pullback of relative cycles (see 1A.6 and 1A.8) induces the required natural map $z_{equi}(T, r)(S) \rightarrow z_{equi}(T, r)(S')$.

It is not hard to see that $z_{equi}(T, r)$ is a Zariski sheaf, and even an étale sheaf, for each T and $r \geq 0$. One can also check that each $z_{equi}(T, r)$ is contravariant for flat maps in T , and covariant for proper maps in T , both with the appropriate change in the dimension index r , (see [RelCh, 3.6.2 and 3.6.4]); see [Blo86, 1.3]. In particular, if $T' \hookrightarrow T$ is a closed immersion, there are canonical inclusions $z_{equi}(T', r) \hookrightarrow z_{equi}(T, r)$ for all r .

EXAMPLE 16.2. The case $r = 0$ is of particular interest, since if U is irreducible the group $z_{equi}(T, 0)(U)$ is free abelian on the irreducible $Z \subset U \times T$ which are quasi-finite and dominant over U . Hence $\mathbb{Z}_{tr}(T)(U) \subseteq z_{equi}(T, 0)(U)$, because $\mathbb{Z}_{tr}(T)(U)$ is the free abelian group of cycles in $U \times T$ which are finite and surjective over U . In fact, $\mathbb{Z}_{tr}(T)$ is a subsheaf of $z_{equi}(T, 0)$ because the structure morphisms associated to $V \rightarrow U$ are compatible: $\mathbb{Z}_{tr}(T)(U) \rightarrow \mathbb{Z}_{tr}(T)(V)$ is also the pullback of relative cycles (see 1A.8 and 1A.11).

If T is projective, or proper, then $\mathbb{Z}_{tr}(T) = z_{equi}(T, 0)$. Indeed, each closed subvariety $Z \subset U \times T$ is proper over U , so Z is quasi-finite over U if and only if Z is finite over U (see [Har77, Ex. III.11.2]).

We now define transfer maps for $z_{\text{equi}}(T, r)$. Given an elementary correspondence V from X to Y and a cycle \mathcal{Z} in $z_{\text{equi}}(T, r)(Y)$, the pullback \mathcal{Z}_V is a well-defined cycle of $V \times T$ by 1A.6 and 1A.8. We define $\phi_V(\mathcal{Z}) \in z_{\text{equi}}(T, r)(X)$ to be the push-forward of \mathcal{Z}_V along the finite map $V \times T \rightarrow X \times T$. This gives a homomorphism $\phi_V : z_{\text{equi}}(T, r)(Y) \rightarrow z_{\text{equi}}(T, r)(X)$.

If $r = 0$, the restriction of ϕ_V to $\mathbb{Z}_{\text{tr}}(T)(Y)$ is the transfer map constructed in 1.1 and 1A.11, as we see from 1A.11 and 1A.13.

We leave the verification of the following to the reader; cf. [BivCy, 5.7].

EXERCISE 16.3. For all T and r , show that ϕ makes each $z_{\text{equi}}(T, r)$ into a presheaf with transfers. *Hint:* Use the identity 2.2(4) in [Ful75]. If $S \rightarrow T$ is flat of relative dimension d , show that the pullback $z_{\text{equi}}(T, r) \rightarrow z_{\text{equi}}(S, d+r)$ is a morphism of presheaves with transfers.

EXAMPLE 16.4. For each X , there is a natural map $z_{\text{equi}}(\mathbb{A}^i, 0)(X) \rightarrow CH^i(\mathbb{A}^i \times X) \cong CH^i(X)$, sending a subvariety Z of $\mathbb{A}^i \times X$, quasi-finite over X , to its cycle. Comparing the transfer map for $z_{\text{equi}}(\mathbb{A}^i, 0)$ to the transfer map for $CH^i(X)$ defined in 2.5, we see that $z_{\text{equi}}(\mathbb{A}^i, 0) \rightarrow CH^i(-)$ is a morphism of presheaves with transfers.

We define the **Suslin-Friedlander motivic complexes** $\mathbb{Z}^{SF}(i)$ by:

$$\mathbb{Z}^{SF}(i) = C_* z_{\text{equi}}(\mathbb{A}^i, 0)[-2i].$$

We regard $\mathbb{Z}^{SF}(i)$ as a bounded above cochain complex, whose top term is $z_{\text{equi}}(\mathbb{A}^i, 0)$ in cohomological degree $2i$. As in 3.1, $C_*(F)$ stands for the chain complex of presheaves associated to the simplicial presheaf $U \mapsto F(U \times \Delta^\bullet)$.

EXAMPLE 16.5. It follows from 16.2 and 16.3 that there is a morphism of presheaves with transfers from $\mathbb{Z}_{\text{tr}}(\mathbb{P}^i) = z_{\text{equi}}(\mathbb{P}^i, 0)$ to $z_{\text{equi}}(\mathbb{A}^i, 0)$, with kernel $\mathbb{Z}_{\text{tr}}(\mathbb{P}^{i-1})$. Applying C_* gives an exact sequence of complexes of presheaves with transfers $0 \rightarrow C_* \mathbb{Z}_{\text{tr}}(\mathbb{P}^{i-1}) \rightarrow C_* \mathbb{Z}_{\text{tr}}(\mathbb{P}^i) \rightarrow \mathbb{Z}^{SF}(i)[2i]$.

EXERCISE 16.6. Let E be the function field of a smooth variety over k . Show that the stalk at $\text{Spec } E$ of the sheaf $z_{\text{equi}}(\mathbb{A}_k^i, 0)$ on Sm/k is equal to the global sections $z_{\text{equi}}(\mathbb{A}_E^i, 0)(\text{Spec } E)$ of the sheaf $z_{\text{equi}}(\mathbb{A}_E^i, 0)$ on Sm/E . Similarly, show that the stalk of $C_m z_{\text{equi}}(\mathbb{A}_k^i, 0)$ at $\text{Spec } E$ equals $C_m z_{\text{equi}}(\mathbb{A}_E^i, 0)(\text{Spec } E)$.

Conclude that the stalk of $\mathbb{Z}^{SF}(i)$ at $\text{Spec } E$ is independent of the choice of k , since it equals $\mathbb{Z}^{SF}(i)(\text{Spec } E)$ evaluated in Sm/E .

Here are the two main results in this lecture. Figure 16.1 gives the scheme of the proof of 16.7. It shows how this result ultimately depends on theorem 13.12, whose proof will be completed in lectures 21-24 below.

THEOREM 16.7. *Assume that k is perfect. Then there is a quasi-isomorphism in the Zariski topology:*

$$\mathbb{Z}(i) \simeq \mathbb{Z}^{SF}(i).$$

In particular, $H^{n,i}(X, \mathbb{Z}) \cong \mathbb{H}^n(X, \mathbb{Z}^{SF}(i))$ for all n and i .

PROOF. As \mathbb{P}^i is proper, $z_{equi}(\mathbb{P}^i, 0) = \mathbb{Z}_{tr}(\mathbb{P}^i)$ by 16.2. Hence 16.7 follows from combining 15.2 and 16.8. \square




FIGURE 16.1. Scheme of the proof of 16.7

THEOREM 16.8. *There is a quasi-isomorphism in the Zariski topology:*

$$C_* [z_{equi}(\mathbb{P}^i, 0)/z_{equi}(\mathbb{P}^{i-1}, 0)] \xrightarrow{\simeq} C_* z_{equi}(\mathbb{A}^i, 0).$$

We now prepare for the proof of 16.8. We remark that an easier proof is available if k admits resolution of singularities, because then $C_* z_{equi}(-, 0)$ satisfies localization by [BivCy, 4.10.2]. This may be compared to the localization property for $C_* \mathbb{Z}_{tr}$ in 13.15.

Write $F_i(U)$ for the (free abelian) subgroup of $z_{equi}(\mathbb{A}^i, 0)(U)$ generated by the cycles in $U \times \mathbb{A}^i$ which do not touch $U \times 0$. The transfers $z_{equi}(\mathbb{A}^i, 0)(U) \rightarrow z_{equi}(\mathbb{A}^i, 0)(V)$ clearly send $F_i(U)$ to $F_i(V)$. Hence F_i is a sub-presheaf with transfers of $z_{equi}(\mathbb{A}^i, 0)$.

LEMMA 16.9. *There is a commutative diagram with exact rows in $\mathbf{PST}(k)$, in which all three vertical maps are injections:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}_{tr}(\mathbb{P}^i - 0)/\mathbb{Z}_{tr}(\mathbb{P}^{i-1}) & \longrightarrow & \mathbb{Z}_{tr}(\mathbb{P}^i)/\mathbb{Z}_{tr}(\mathbb{P}^{i-1}) & \longrightarrow & \text{coker}_1 \longrightarrow 0 \\ & & \downarrow & & \downarrow \varphi & & \downarrow \\ 0 & \longrightarrow & F_i & \longrightarrow & z_{equi}(\mathbb{A}^i, 0) & \longrightarrow & \text{coker}_2 \longrightarrow 0. \end{array}$$

PROOF. By example 16.2, there is a natural map from $\mathbb{Z}_{tr}(\mathbb{P}^i) = z_{equi}(\mathbb{P}^i, 0)$ to $z_{equi}(\mathbb{A}^i, 0)$ with kernel $\mathbb{Z}_{tr}(\mathbb{P}^{i-1})$. Thus φ is an injection; by exercise 16.3, φ is a morphism of presheaves with transfers. Now the inclusion $\mathbb{Z}_{tr}(\mathbb{P}^i - 0) \subset \mathbb{Z}_{tr}(\mathbb{P}^i)$ is a morphism in \mathbf{PST} by the Yoneda lemma; see 2.8. Since $\mathbb{Z}_{tr}(\mathbb{P}^i - 0)(U)$ consists of cycles $Z \subset U \times (\mathbb{P}^i - 0)$ finite over U , their restriction belongs to the subgroup $F_i(U)$, i.e., φ sends $\mathbb{Z}_{tr}(\mathbb{P}^i - 0)$ to F_i . Hence the diagram commutes.

By inspection, $\text{coker}_1(U)$ is free abelian on the elementary correspondences $Z \subset U \times \mathbb{P}^i$ which touch $U \times 0$ and $\text{coker}_2(U)$ is free abelian on the equidimensional $W \subset U \times \mathbb{A}^i$ which touch $U \times 0$. Since $Z \mapsto \varphi(Z)$ is a monomorphism on these generators, it follows that $\text{coker}_1(U) \rightarrow \text{coker}_2(U)$ is an injection for all U . \square

LEMMA 16.10. *$C_*(F_i)$ is chain contractible as a complex of presheaves.*

PROOF. Recall that $F_i(X)$ is a subgroup of the group of cycles on $X \times \mathbb{A}^i$. Let $h_X : F_i(X) \rightarrow F_i(X \times \mathbb{A}^1)$ be the pullback of cycles along $\mu : X \times \mathbb{A}^i \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^i$ defined by $(x, r, t) \rightarrow (x, r \cdot t)$. This is a good pullback because the map μ is flat over $X \times (\mathbb{A}^i - \{0\})$. By construction, the following diagram commutes.

$$\begin{array}{ccccc}
 X \times \mathbb{A}^i & \xrightarrow{t=1} & X \times \mathbb{A}^i \times \mathbb{A}^1 & \xleftarrow{t=0} & X \times \mathbb{A}^i \\
 & \searrow id & \downarrow \mu & \swarrow id \times 0 & \\
 & & X \times \mathbb{A}^i & &
 \end{array}$$

It follows that $F_i(t=1)h_X$ is the identity and $F_i(t=0)h_X = 0$. Thus the requirements of lemma 2.22 are satisfied for F_i . \square

LEMMA 16.11. *For every Hensel local scheme S , the map $\text{coker}_1(S) \rightarrow \text{coker}_2(S)$ in the diagram in 16.9 is an isomorphism.*

PROOF. Since $\text{coker}_1(S) \rightarrow \text{coker}_2(S)$ is injective by 16.9, it suffices to prove that it is surjective. Let Z be an equidimensional correspondence from S to \mathbb{A}^i . As Z is quasi-finite over a Hensel scheme, the projection decomposes Z into the disjoint union of Z_0 (which doesn't contain any point over the closed point of the Hensel scheme) and Z_1 (which is finite over the base). We claim that the Z_0 part comes from F_i . Take Z_0 and consider its irreducible components. The intersection $Z_0 \cap \{0\}$ must be empty, otherwise it would project to the closed point of the base. Hence Z_0 is zero in the cokernel. But now Z_1 comes from $\mathbb{Z}_{tr}(\mathbb{P}^i)/\mathbb{Z}_{tr}(\mathbb{P}^{i-1})$. \square

LEMMA 16.12. *Assume that k is perfect. Then the map $C_*(\text{coker}_1) \rightarrow C_*(\text{coker}_2)$ is a quasi-isomorphism of complexes of Zariski sheaves.*

PROOF. Let φ' be the map between the cokernels in 16.9. By 16.11, φ' is an isomorphism on all Hensel local schemes. By 13.14, φ' induces quasi-isomorphisms $C_* \text{coker}_1(X) \simeq C_* \text{coker}_2(X)$ for all local X . \square

PROOF OF 16.8. Applying C_* to the diagram in 16.9 yields a commutative diagram of chain complexes with exact rows. The left two complexes are acyclic by 15.1 and 16.10. The right two complexes are quasi-isomorphic by 16.12. Theorem 16.8 now follows from the 5-lemma. \square

Motives with compact support

By 16.1 and 16.3, $z_{\text{equi}}(X, 0)$ is a Nisnevich sheaf with transfers for every scheme X of finite type over k . As such, we can regard it as an element of $\mathbf{D}^- \text{Sh}_{\text{Nis}}(\text{Cor}_k)$.

DEFINITION 16.13. For any scheme X of finite type over k , let $M^c(X)$ denote $z_{\text{equi}}(X, 0)$, regarded as an object in $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k)$. By 14.4, $M^c(X) \cong C_* z_{\text{equi}}(X, 0)$ in $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k)$.

As pointed out in 16.1, $M^c(X)$ is contravariant in X for étale maps and covariant in X for proper maps. As observed in 16.2, there is a canonical map $M(X) \rightarrow M^c(X)$, induced by $\mathbb{Z}_{tr}(X) \subseteq z_{equi}(X, 0)$. Moreover, $M(X) = M^c(X)$ if X is proper over k .

EXAMPLE 16.14. We have an isomorphism $M^c(\mathbb{A}^i) \cong \mathbb{Z}(i)[2i]$ in $\mathbf{DM}_{Nis}^{\text{eff}, -}(k)$. To see this, recall from 16.7 that the Suslin-Friedlander motivic complex $\mathbb{Z}^{SF}(i) = C_* z_{equi}(\mathbb{A}^i, 0)[-2i]$ is quasi-isomorphic to $\mathbb{Z}(i)$.

THEOREM 16.15. *Assume that k admits resolution of singularities. If $Z \xrightarrow{i} X$ is a closed subscheme with complement $U \xrightarrow{j} X$, there is a distinguished triangle:*

$$M^c(Z) \xrightarrow{i_*} M^c(X) \xrightarrow{j^*} M^c(U) \rightarrow M^c(Z)[1].$$

PROOF. It is easy to see from the definitions that there is an exact sequence of sheaves with transfers:

$$0 \rightarrow z_{equi}(Z, 0) \xrightarrow{i_*} z_{equi}(X, 0) \xrightarrow{j^*} z_{equi}(U, 0) \rightarrow \mathcal{Q} \rightarrow 0.$$

By 13.25 it suffices to show that $\mathcal{Q}_{cdh} = 0$. By 12.30, it suffices to fix S in Sm/k and show that for any $q \in \mathcal{Q}(S)$ there is a composition of blow-ups $p: S' \rightarrow S$ such that $p^*(q) = 0$. A lift of q to $z_{equi}(U, 0)(S)$ is supported on a finite set of irreducible subschemes $W_\alpha|_U \subset U \times S$ which are quasi-finite and dominant over a component of S . We may assume that the closures W_α of $W_\alpha|_U$ in $X \times S$ are not quasi-finite over S . By platication (see [RG71] or 1A.1), there is a blow-up $p: S' \rightarrow S$ such that the proper transforms W'_α of W_α are flat and dominant over S' . By resolution of singularities, we may assume that S' is smooth and that p is a composition of blow-ups along smooth centers. But then $p^*(W_\alpha) = j^*(W'_\alpha)$ in $z_{equi}(U, 0)(S')$ and each $p^*(W_\alpha|_U)$ vanishes in $\mathcal{Q}(S')$. \square

COROLLARY 16.16. *For every X and Y , $M^c(X \times Y) \cong M^c(X) \otimes_{L, Nis}^{tr} M^c(Y)$. In particular, $M^c(X \times \mathbb{A}^i) \cong M^c(X)(i)[2i]$.*

PROOF. If X and Y are smooth and proper, this is just the identity $M(X \times Y) \cong M(X) \otimes_{L, Nis}^{tr} M(Y)$. The case when X and Y are proper follows formally from this using the axioms in 8A.1 for the tensor triangulated structure and the blow-up triangle in 13.26. Using the axioms and the closed subscheme triangle in 16.15, we obtain the general case. The last assertion comes from 16.14. \square

COROLLARY 16.17. *For every scheme X in Sch/k , $M^c(X)$ is in $\mathbf{DM}_{gm}^{\text{eff}}$.*

PROOF. If X is proper, so that $M^c(X) = M(X)$, this follows from (14.5.3), as pointed out in 14.1. The general case follows from this, using theorem 16.15. \square

EXERCISE 16.18. Let U, V be an open cover of X . Show that (assuming resolution of singularities) there is a distinguished triangle in $\mathbf{DM}_{Nis}^{\text{eff}, -}$:

$$M^c(X) \rightarrow M^c(U) \oplus M^c(V) \rightarrow M^c(U \cap V) \rightarrow M^c(X)[1].$$

EXERCISE 16.19. Assume that k admits resolution of singularities. If $Z \xrightarrow{i} X$ is a closed subscheme with complement $U \xrightarrow{j} X$, modify the proof of 16.15 to show that there is a distinguished triangle:

$$C_*\mathcal{Z}_{\text{equi}}(Z, r) \xrightarrow{i_*} C_*\mathcal{Z}_{\text{equi}}(X, r) \xrightarrow{j^*} C_*\mathcal{Z}_{\text{equi}}(U, r) \rightarrow C_*\mathcal{Z}_{\text{equi}}(Z, r)[1].$$

DEFINITION 16.20. Let X be any scheme of finite type over k and $i \geq 0$. We define the **motivic cohomology with compact supports** of X with coefficients in R to be:

$$H_c^{n,i}(X, R) = \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M^c(X), R(i)[n]).$$

Dually, we define the **(Borel-Moore) motivic homology with compact supports** $H_{n,i}^{BM}(X, R)$ to be

$$H_{n,i}^{BM}(X, R) = \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(R(i)[n], M^c(X)).$$

Applying Hom to the triangle in 16.15 yields the expected long exact localization sequences for motivic cohomology and homology with compact supports:

$$\begin{aligned} H_c^{n,i}(U, \mathbb{Z}) &\rightarrow H_c^{n,i}(X, \mathbb{Z}) \rightarrow H_c^{n,i}(Z, \mathbb{Z}) \rightarrow H_c^{n+1,i}(U, \mathbb{Z}), \\ H_{n,i}^{BM}(Z, \mathbb{Z}) &\rightarrow H_{n,i}^{BM}(X, \mathbb{Z}) \rightarrow H_{n,i}^{BM}(U, \mathbb{Z}) \rightarrow H_{n-1,i}^{BM}(Z, \mathbb{Z}). \end{aligned}$$

We will identify the motivic homology groups $H_{n,i}^{BM}(X, R)$ with higher Chow groups in lecture 20.

REMARK 16.21. Friedlander and Voevodsky introduced a bivariant cycle cohomology group $A_{r,i}(Y, X)$ in [BivCy, 4.3], as the cdh hypercohomology on Y of $C_*\mathcal{Z}_{\text{equi}}(X, r)$. Using 14.20, their definition is equivalent to:

$$A_{r,i}(Y, X) = \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M(Y)[i], C_*\mathcal{Z}_{\text{equi}}(X, r)).$$

In [BivCy, 8.3], they proved the following result:

THEOREM 16.22. *Let X be in Sch/k , where k admits resolution of singularities. Then for any $r \geq 0$ and any M in $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}$, there are natural isomorphisms:*

$$\begin{aligned} \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M(1)[2], C_*\mathcal{Z}_{\text{equi}}(X, r)) &\xrightarrow{\cong} \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M, C_*\mathcal{Z}_{\text{equi}}(X, r+1)); \\ \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M(r)[2r], M^c(X)) &\xrightarrow{\cong} \text{Hom}_{\mathbf{DM}_{\text{Nis}}^{\text{eff},-}}(M, C_*\mathcal{Z}_{\text{equi}}(X, r)). \end{aligned}$$

More precisely, they proved in [BivCy, 8.3] that there is a natural isomorphism $A_{r,i}(Y(1)[2], X) \cong A_{r+1,i}(Y, X)$ for every X, Y in Sch/k . Since the $Y[i]$ generate $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}$, this is equivalent to the first isomorphism. Since $M^c(X)$ is $C_*\mathcal{Z}_{\text{equi}}(X, 0)$, the second isomorphism follows from the first by induction on r .

COROLLARY 16.23. *Let $X \rightarrow Y$ be a flat map of relative dimension r . Then we have a morphism in $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}$:*

$$M^c(Y)(r)[2r] \rightarrow M^c(X).$$

PROOF. By 16.3, the pullback induces a morphism $C_{*Z_{\text{equi}}}(Y, 0) \rightarrow C_{*Z_{\text{equi}}}(X, r)$. Now take $M = C_{*Z_{\text{equi}}}(Y, 0)$ in 16.22. \square

The Duality Theorem below is also due to Friedlander and Voevodsky ([BivCy, 8.2]). We also cite it without proof.

THEOREM 16.24. (*Duality*) *Assume that k admits resolution of singularities. If T is a smooth scheme of dimension d over k then there are canonical isomorphisms*

$$\text{Hom}(M(X \times T)[n], M^c(Y)) \cong \text{Hom}(M(X)(d)[2d+n], M^c(T \times Y)).$$

THEOREM 16.25. (*Cancellation*) *Assume that k admits resolution of singularities. Let M and N be in $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}$. Then tensoring with $\mathbb{Z}(1)$ induces an isomorphism $\text{Hom}(M, N) \rightarrow \text{Hom}(M(1), N(1))$.*

PROOF. Suppose first that $M = M(X)[n]$ and $N = M(Y)$ for smooth proper schemes X and Y . Under this assumption, $M^c(Y) = M(Y)$ and $M^c(\mathbb{A}^1 \times Y) = M^c(\mathbb{A}^1) \otimes M^c(Y) = M(Y)(1)[2]$. Applying 16.24 with $T = \mathbb{A}^1$, homotopy invariance yields isomorphisms:

$$\begin{aligned} \text{Hom}(M, N) &= \text{Hom}(M(X)[n], M(Y)) \cong \text{Hom}(M(X \times \mathbb{A}^1)[n], M^c(Y)) \cong \\ &\text{Hom}(M(X)(1)[2+n], M^c(\mathbb{A}^1 \times Y)) \cong \text{Hom}(M(1)[2], N(1)[2]). \end{aligned}$$

Removing the shift yields $\text{Hom}(M, N) \cong \text{Hom}(M(1), N(1))$. To see that this isomorphism is induced by tensoring with $\mathbb{Z}(1)$, it suffices to trace through the explicit isomorphisms we used. We leave this to the reader.

Since these motives generate $\mathbf{DM}_{\text{gm}}^{\text{eff}}$, the theorem is true when M and N are in $\mathbf{DM}_{\text{gm}}^{\text{eff}}$. By 13.5, $\text{Hom}(M, \bigoplus_{\alpha} N_{\alpha}) \cong \bigoplus_{\alpha} \text{Hom}(M, N_{\alpha})$ for M in $\mathbf{DM}_{\text{gm}}^{\text{eff}}$. Since $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}$ is generated by $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ and direct sums, the theorem holds for all N in $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}$ when M is in $\mathbf{DM}_{\text{gm}}^{\text{eff}}$. Finally, $\text{Hom}(\bigoplus M_{\alpha}, N) \cong \bigoplus \text{Hom}(M_{\alpha}, N)$ so we may remove the restriction on M . \square

REMARK 16.26. The Cancellation Theorem 16.25 is also valid when k is perfect. This was proven in 2002 by Voevodsky in [Voe02].

The next three lectures will be devoted to a proof that Bloch’s higher Chow groups agree with motivic cohomology on smooth schemes. We will generalize this to all schemes of finite type at the end of lecture 19, replacing motivic cohomology with Borel-Moore motivic homology.

Part 5

Higher Chow Groups

Higher Chow groups

During the first part of this series of lectures we defined motivic cohomology and we studied its basic properties. We also established relations with some classical objects of algebraic geometry, such as Milnor K -Theory, 5.1, and étale cohomology, 10.2.

The goal of the next few lectures is to find a relation between motivic cohomology and the classical Chow groups CH^i , generalizing the isomorphism $H^{2,1}(X, \mathbb{Z}) \cong \text{Pic}(X) = CH^1(X)$ of 4.2. That is, we will prove that:

$$H^{2i,i}(X, \mathbb{Z}) \cong CH^i(X)$$

for any smooth variety X . There are at least three ways to prove this. The original approach, which needs resolution of singularities, was developed in the book “*Cycles, Transfers and Motivic Homology Theories*” [VSF00]. A second recent approach is to use the Cancellation Theorem 16.25 of [Voe02] and the Gersten resolution 24.11 for motivic cohomology sheaves.

A third approach, which is the one we shall develop here, uses Bloch’s higher Chow groups $CH^i(X, m)$ to establish the more general isomorphism $H^{n,i}(X, \mathbb{Z}) \cong CH^i(X, 2i - n)$. This approach uses the equidimensional cycle groups of the previous lecture, but does not use resolution of singularities.

The main goal of this lecture is to prove that the higher Chow groups are presheaves with transfers. (See theorem 17.21.) In particular, they are functorial for maps between smooth schemes. (We will give a second proof of this in 19.15.)

We begin with Bloch’s definition of higher Chow groups (see [Blo86]).

DEFINITION 17.1. Let X be an equidimensional scheme. We write $z^i(X, m)$ for the free abelian group generated by all codimension i subvarieties on $X \times \Delta^m$ which intersect all faces $X \times \Delta^j$ properly for all $j < m$ (in the sense of 17A.1).

Each face $X \times \Delta^j$ is defined by a regular sequence, and intersection of cycles defines a map $z^i(X, m) \rightarrow z^i(X, j)$ (see 17A.1, or [Ful84, Example 7.1.2]). We write $z^i(X, \bullet)$ for the resulting simplicial abelian group $m \mapsto z^i(X, m)$. We write $z^i(X, *)$ for the chain complex associated to $z^i(X, \bullet)$.

The **higher Chow groups** of X are defined to be the groups:

$$CH^i(X, m) = \pi_m(z^i(X, \bullet)) = H_m(z^i(X, *)).$$

If X is any scheme, it is easy to check that $CH^i(X, 0)$ is the classical Chow group $CH^i(X)$ (see 17.3). Indeed, $z^i(X, 0)$ is the group of all codimension i cycles

on X while $z^i(X, 1)$ is generated by those codimension i subvarieties Z on $X \times \mathbb{A}^1$ which intersect both $X \times \{0\}$ and $X \times \{1\}$ properly. Moreover the maps $\partial_j : z^i(X, 1) \rightrightarrows z^i(X, 0)$ send Z to $Z \cap (X \times \{j\})$.

EXAMPLE 17.2. If $i \leq d = \dim X$, then $z_{\text{equi}}(X, d - i)(\Delta^m) \subseteq z^i(X, m)$ for every m , because cycles in $X \times \Delta^m$ which are equidimensional over Δ^m must meet every face properly. By 1A.14, the inclusion is compatible with the face maps, which are defined in 16.1 and 17.1, so this yields an inclusion of simplicial groups, $z_{\text{equi}}(X, d - i)(\Delta^\bullet) \subseteq z^i(X, \bullet)$.

EXERCISE 17.3. (a) If $d = \dim X$, show that every irreducible cycle in $z^d(X, 1)$ is either disjoint from $X \times \{0, 1\}$ or else is quasi-finite over \mathbb{A}^1 . Use this to describe $z^d(X, 1) \rightarrow z^d(X, 0)$ explicitly and show that $CH^d(X, 0) \cong CH^d(X)$. (The group $CH^d(X)$ is defined in [Ful84, 1.6].)

(b) Show that $C_*\mathbb{Z}_{\text{tr}}(X)(\text{Speck})$ is a subcomplex of $z^d(X, *)$. On homology, this yields maps $H_m^{\text{sing}}(X/k) \rightarrow CH^d(X, m)$. For $m = 0$, show that this is the surjection $H_0^{\text{sing}}(X/k) \rightarrow CH_0(X) = CH^d(X)$ of 2.21, which is an isomorphism when X is projective. By 7.3, it is not an isomorphism when X is \mathbb{A}^1 or $\mathbb{A}^1 - 0$.

The push-forward of cycles makes the higher Chow groups covariant for finite morphisms (see 17A.10). It also makes them covariant for proper morphisms (with the appropriate change in codimension index i ; see [Blo86, 1.3]).

At the chain level, it is easy to prove that the complexes $z^i(-, *)$, and hence Bloch's higher Chow groups, are functorial for flat morphisms. However, the complexes $z^i(-, *)$ are not functorial for all maps. We will see in theorem 17.21 below that the higher Chow groups are functorial for maps between smooth schemes.

PROPERTIES 17.4. We will need the following non-trivial properties of higher Chow groups.

- (1) Homotopy Invariance: The projection $p : X \times \mathbb{A}^1 \rightarrow X$ induces an isomorphism

$$CH^i(X, m) \xrightarrow{\cong} CH^i(X \times \mathbb{A}^1, m)$$

for any scheme X over k . The proof is given in [Blo86, 2.1].

- (2) Localization Theorem: For any $U \subset X$ open, the cokernel of $z^i(X, \bullet) \rightarrow z^i(U, \bullet)$ is acyclic. This is proven by Bloch in [Blo94]. (Cf. [Blo86, 3.3].)

If the complement $Z = X - U$ has pure codimension c , it is easy to see that we have an exact sequence of simplicial abelian groups (and also of complexes of abelian groups):

$$0 \rightarrow z^{i-c}(Z, \bullet) \rightarrow z^i(X, \bullet) \rightarrow z^i(U, \bullet) \rightarrow \text{coker} \rightarrow 0.$$

Thus the localization theorem yields long exact sequences of higher Chow groups. The fact that we need to use Bloch's Localization Theorem is unfortunate, because its proof is very hard and complex.

Transfer maps associated to correspondences are not defined on all of $z^i(Y, *)$. We need to restrict to a subcomplex on which \mathscr{W}^* may be defined.

DEFINITION 17.5. Let \mathcal{W} be a finite correspondence from X to Y . Write $z^i(Y, m)_{\mathcal{W}}$ for the subgroup of $z^i(Y, m)$ generated by the irreducible subvarieties $T \subset Y \times \Delta^m$ such that the pullback $X \times T$ intersects $\mathcal{W} \times \Delta^j$ properly in $X \times Y \times \Delta^m$ for every face $\Delta^j \hookrightarrow \Delta^m$. By construction, $z^i(Y, *)_{\mathcal{W}}$ is a subcomplex of $z^i(Y, *)$.

The proof of the following proposition, which is a refinement of the results in [Lev98], is due to Marc Levine.

PROPOSITION 17.6. *Let \mathcal{W} be a finite correspondence from X to Y , with Y affine. Then the inclusion $z^i(Y, *)_{\mathcal{W}} \subset z^i(Y, *)$ is a quasi-isomorphism.*

PROOF. (Levine) Let $w : W \rightarrow Y$ be a morphism of schemes with Y smooth, and W locally equidimensional but not necessarily smooth. Write $z^i(Y, m)_w$ for the subgroup of $z^i(Y, m)$ generated by the irreducible subvarieties $T \subset Y \times \Delta^m$ for which every component of $w^{-1}(T)$ has codimension at least i in $W \times \Delta^m$ and intersects every face properly. By construction, $z^i(Y, *)_w$ is a subcomplex of $z^i(Y, *)$.

For example, if W is the support of a finite correspondence \mathcal{W} , let $w : W \rightarrow Y$ be the natural map. Then W is locally equidimensional, and the group $z^i(Y, m)_w$ is the same as the group $z^i(Y, m)_{\mathcal{W}}$ of 17.5.

Levine proves that $z^i(Y, m)_w \hookrightarrow z^i(Y, m)$ is a quasi-isomorphism on p. 102 of [Lev98] (in I.II.3.5.14), except that W is required to also be smooth in order to cite lemma I.II.3.5.2 of *op. cit.*. In *loc. cit.*, a finite set $\{C_j\}$ of locally closed irreducible subsets of Y and a sequence of integers $m_j \leq i$ is constructed, with the properties that T is in $z^i(Y, m)_w$ if and only if T is in $z^i(Y, m)$ and that the intersections of T with $C_j \times \Delta^p$ have codimension at least m_j for all j and for every face Δ^p of Δ^m . A reading of the proof of lemma I.II.3.5.2 shows that in fact W need only be locally equidimensional. \square

DEFINITION 17.7. Let \mathcal{W} be a finite correspondence between two smooth schemes X and Y . For each cycle \mathcal{Z} in $z^i(Y, m)_{\mathcal{W}}$, we define the cycle $\mathcal{W}^*(\mathcal{Z})$ on $X \times \Delta^m$ to be:

$$\mathcal{W}^*(\mathcal{Z}) = \pi_*((\mathcal{W} \times \Delta^m) \cdot (X \times \mathcal{Z})).$$

Here $\pi : X \times Y \times \Delta^m \rightarrow X \times \Delta^m$ is the canonical projection.

For each \mathcal{W} , it is clear that \mathcal{W}^* defines a homomorphism from the group $z^i(Y, m)_{\mathcal{W}}$ to the group of all cycles on $X \times \Delta^m$.

EXAMPLE 17.8. Let $f : X \rightarrow Y$ be a morphism of smooth varieties, and let Γ_f be its graph. For \mathcal{Z} in $z^i(Y, 0)_{\Gamma_f}$, $\Gamma_f^*(\mathcal{Z})$ is just the classical pullback of cycles $f^*(\mathcal{Z})$ defined in [Ser65, V-28] (see 17A.3).

REMARK 17.9. The map \mathcal{W}^* of 17.7 is compatible with the map \mathcal{W}^* defined in 17A.8 in the following sense. Given \mathcal{W} in $Cor(X, Y)$, $\mathcal{W} \times \text{diag}(\Delta^m)$ is a finite correspondence from $X \times \Delta^m$ to $Y \times \Delta^m$. If \mathcal{Z} is a cycle in $z^i(Y, m)_{\mathcal{W}}$, we may regard it as a cycle in $Y \times \Delta^m$. The projection formula 17A.11 says that the following

diagram commutes:

$$\begin{array}{ccc} z^i(Y, m)_{\mathcal{W}} & \hookrightarrow & z^i(Y \times \Delta^m)_{\mathcal{W}} \\ \mathcal{W}^* \downarrow & & \downarrow (\mathcal{W} \times \text{diag}(\Delta^m))^* \\ z^i(X, m) & \hookrightarrow & z^i(X \times \Delta^m). \end{array}$$

LEMMA 17.10. *The maps \mathcal{W}^* define a chain map $z^i(Y, *)_{\mathcal{W}} \rightarrow z^i(X, *)$.*

PROOF. Let $\partial_j : \Delta^m \hookrightarrow \Delta^{m+1}$ be a face, and consider the following diagram, whose vertical compositions are \mathcal{W}^* :

$$\begin{array}{ccc} z(Y \times \Delta^{m+1}) & \xrightarrow{\partial_j^*} & z(Y \times \Delta^m) \\ f^* \downarrow & & \downarrow f^* \\ z(X \times Y \times \Delta^{m+1}) & \xrightarrow{\partial_j^*} & z(X \times Y \times \Delta^m) \\ \mathcal{W} \times \Delta^{m+1} \cdot - \downarrow & & \downarrow \mathcal{W} \times \Delta^m \cdot - \\ z(X \times Y \times \Delta^{m+1}) & \xrightarrow{\partial_j^*} & z(X \times Y \times \Delta^m) \\ \pi_* \downarrow & & \downarrow \pi_* \\ z(X \times \Delta^{m+1}) & \xrightarrow{\partial_j^*} & z(X \times \Delta^m). \end{array}$$

The horizontal maps ∂_j^* are only defined for cycles meeting the face properly (see 17A.4) and the intersection products in the middle are only defined on cycles in good position for \mathcal{W} . The top square commutes because of the functoriality of Bloch's complex for flat maps, and the bottom square commutes by 17A.10.

Suppose that \mathcal{Z} is a cycle in $X \times Y \times \Delta^{m+1}$ which intersects the face $X \times Y \times \Delta^m$ as well as $\mathcal{W} \times \Delta^{m+1}$ and $\mathcal{W} \times \Delta^m$ properly. By 17A.2:

$$\mathcal{W} \times \Delta^m \cdot ((X \times Y \times \Delta^m) \cdot \mathcal{Z}) = X \times Y \times \Delta^m \cdot ((\mathcal{W} \times \Delta^{m+1}) \cdot \mathcal{Z}).$$

That is, the middle square commutes for \mathcal{Z} . Finally, if $\mathcal{Y} \in z^i(Y, m+1)_{\mathcal{W}}$, the cycle $(\mathcal{W} \times \Delta^{m+1}) \cdot f^* \mathcal{Y}$ is finite over $X \times \Delta^{m+1}$, so π_* may be applied to it. A diagram chase now shows that \mathcal{W}^* is a morphism of chain complexes. \square

COROLLARY 17.11. *If Y is affine, any finite correspondence \mathcal{W} from X to Y induces maps $\mathcal{W}^* : CH^i(Y, m) \rightarrow CH^i(X, m)$ for all m .*

PROOF. On homology, 17.6 and \mathcal{W}^* give: $CH^i(Y, m) \cong H_m(z^i(Y, *)_{\mathcal{W}}) \rightarrow H_m(z^i(X, *)) = CH^i(X, m)$. \square

EXAMPLE 17.12. If $f : X \rightarrow Y$ is a morphism of smooth varieties, and Y is affine, we will write f^* for the map Γ_f^* from $z^i(Y, m)_{\Gamma_f}$ to $z^i(X, m)$, and also for the induced map from $CH^i(Y, m)$ to $CH^i(X, m)$. It agrees with Levine's map f^* (see pp. 67 and 102 of [Lev98]). This is not surprising, since we are using proposition 17.6, which is taken from p. 102 of [Lev98]. The map f^* may also be obtained from [Blo86, 4.1] using [Blo94].

If f is flat, then f^* is just the flat pullback of cycles. That is, if $\mathcal{B} = [V]$ then $f^*(\mathcal{B})$ is the cycle associated to the scheme $f^{-1}(V)$. This fact is a special case of 17A.4.

We can now show that the higher Chow groups are functors on the subcategory of affine schemes in Cor_k .

LEMMA 17.13. *Let X, Y and Z be smooth affine schemes. Given finite correspondences \mathcal{W}_1 in $Cor(X, Y)$ and \mathcal{W}_2 in $Cor(Y, Z)$, then*

$$(\mathcal{W}_2 \circ \mathcal{W}_1)^* = \mathcal{W}_1^* \mathcal{W}_2^* : CH^i(Z, m) \rightarrow CH^i(X, m).$$

In particular, if $f_1 : X \rightarrow Y$ and $f_2 : Y \rightarrow Z$ are morphisms, then $(f_2 \circ f_1)^ = f_1^* f_2^*$.*

PROOF. By 17.6 and 17.12, it suffices to show that $(\mathcal{W}_2 \circ \mathcal{W}_1)^* = \mathcal{W}_1^* \mathcal{W}_2^*$ as maps from $z^i(Z, m)_{\mathcal{W}} \rightarrow z^i(X, m)$, where $\mathcal{W} \in Cor(Y \amalg X, Z)$ is the coproduct of \mathcal{W}_2 and $\mathcal{W}_2 \circ \mathcal{W}_1$. An element of $z^i(Z, m)_{\mathcal{W}}$ is a cycle in $z^i(Z, m)$ which is in good position with respect to both \mathcal{W}_2 and $\mathcal{W}_2 \circ \mathcal{W}_1$. Hence the result follows from theorem 17A.14, given the reinterpretation in 17.9. \square

We now extend the definition of the transfer map \mathcal{W}^* from affine varieties to all smooth varieties using Jouanolou's device [Jou73, 1.5] and [Wei89, 4.4]: over every smooth variety X there is a vector bundle torsor $X' \rightarrow X$ with X' affine.

LEMMA 17.14. *Let X be a variety and $p : X' \rightarrow X$ a vector bundle torsor. Then $p^* : CH^*(X, *) \rightarrow CH^*(X', *)$ is an isomorphism.*

PROOF. By definition, there is a dense open U in X so that $p^{-1}(U) \cong U \times \mathbb{A}^n$. There is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & z^*(X' - p^{-1}(U)) & \longrightarrow & z^*(X', *) & \longrightarrow & z^*(p^{-1}(U)) \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \longrightarrow & z^*(X - U) & \longrightarrow & z^*(X, *) & \longrightarrow & z^*(U). \end{array}$$

By homotopy invariance of the higher Chow groups (see 17.4), the right vertical arrow is a quasi-isomorphism. By Noetherian induction, the result is true for $X - U$, i.e., the left vertical arrow is a quasi-isomorphism. By the Localization Theorem and the 5-lemma, $p^* : CH^*(X, *) \rightarrow CH^*(X', *)$ is an isomorphism. \square

LEMMA 17.15. *Let $p : Y' \rightarrow Y$ be a vector bundle torsor and let X be affine.*

- *Every morphism $f : X \rightarrow Y$ has a lift $f' : X \rightarrow Y'$ such that $pf' = f$.*
- *Every finite correspondence has a lift, i.e., $p_* : Cor(X, Y') \rightarrow Cor(X, Y)$ is surjective.*

PROOF. Clearly, $X \times_Y Y' \rightarrow X$ is a vector bundle torsor. But X is affine and therefore every vector bundle torsor over X is a vector bundle (see [Wei89, 4.2]). Define $f' : X \rightarrow Y'$ to be the composition of the zero-section of $X \times_Y Y'$ followed by the projection. Clearly, $pf' = f$.

Now suppose that $W \subset X \times Y$ is an elementary correspondence. Since W is finite over X , it is affine. By the first part of this proof, the projection $p : W \rightarrow Y$ lifts to a map $p' : W \rightarrow Y'$. Together with the projection $W \rightarrow X$, p' induces a lift $i : W \rightarrow X \times Y'$ of $W \subset X \times Y$. Then $i(W)$ is an elementary correspondence from X to Y' whose image under p_* is W . \square

LEMMA 17.16. *Let X and Y be two smooth varieties over k and let $p : X' \rightarrow X$ and $q : Y' \rightarrow Y$ be vector bundle torsors with X' and Y' affine. Then for every finite correspondence \mathcal{W} from X to Y , there exists a correspondence \mathcal{W}' from X' to Y' so that $q \circ \mathcal{W}' = \mathcal{W} \circ p$ in $\text{Cor}_k(X', Y)$.*

$$\begin{array}{ccc} X' & \xrightarrow{\mathcal{W}'} & Y' \\ p \downarrow & & \downarrow q \\ X & \xrightarrow{\mathcal{W}} & Y \end{array}$$

PROOF. Since $\text{Cor}(X', Y') \rightarrow \text{Cor}(X', Y)$ is onto by 17.15, $\mathcal{W} \circ p$ has a lift \mathcal{W}' . \square

DEFINITION 17.17. Let X and Y be two smooth varieties over k and let \mathcal{W} be a finite correspondence from X to Y . We define $\mathcal{W}^* : CH^i(Y, m) \rightarrow CH^i(X, m)$ as follows.

By Jouanolou's device [Jou73, 1.5], there exist vector bundle torsors $p : X' \rightarrow X$ and $q : Y' \rightarrow Y$ where X' and Y' are affine. Both X' and Y' are smooth, because X and Y are. By lemma 17.14, p^* and q^* are isomorphisms. By 17.16 there is a finite correspondence \mathcal{W}' from X' to Y' so that $q \circ \mathcal{W}' = \mathcal{W} \circ p$ in $\text{Cor}_k(X', Y)$. Since Y' is affine, the map $\mathcal{W}'^* : CH^*(Y', m) \rightarrow CH^*(X', m)$ was defined in 17.11. We set $\mathcal{W}^* = (p^*)^{-1} \mathcal{W}'^* q^* : CH^*(Y, m) \rightarrow CH^*(X, m)$.

$$\begin{array}{ccc} CH^*(X', *) & \xleftarrow{\mathcal{W}'^*} & CH^*(Y', *) \\ p^* \uparrow \cong & & q^* \uparrow \cong \\ CH^*(X, *) & \xleftarrow{\mathcal{W}^*} & CH^*(Y, *) \end{array}$$

If $f : X \rightarrow Y$ is a morphism, we define $f^* : CH^i(Y, m) \rightarrow CH^i(X, m)$ to be Γ_f^* , that is, $f^* = (p^*)^{-1} (f')^* q^*$, where $f' : X' \rightarrow Y'$ lies over f .

LEMMA 17.18. *If X and Y are affine, the map defined in 17.17 agrees with the map \mathcal{W}^* defined in 17.11.*

PROOF. By 17.13, the map defined in 17.17 equals:

$$(p^*)^{-1} \mathcal{W}'^* q^* = (p^*)^{-1} (q \circ \mathcal{W}')^* = (p^*)^{-1} (\mathcal{W} \circ p)^* = (p^*)^{-1} p^* \mathcal{W}^* = \mathcal{W}^*. \quad \square$$


LEMMA 17.19. *The definition of \mathcal{W}^* in 17.17 is independent of the choices.*

PROOF. Suppose given affine torsors $X'' \rightarrow X$ and $Y'' \rightarrow Y$ and a lift $\mathcal{W}'' \in \text{Cor}(X'', Y'')$ of \mathcal{W} . We have to show that \mathcal{W}' and \mathcal{W}'' induce the same map.

We begin by making two reductions. First, we may assume that $X' = X''$ and $Y' = Y''$, by passing to $X' \times_X X''$ and $Y' \times_Y Y''$ and choosing lifts of \mathcal{W}' and \mathcal{W}'' . (This reduction uses 17.18.)

We may also assume that X is affine and that $X' = X$, by replacing X by X' . Thus we need to show that for any two lifts \mathcal{W}_0 and \mathcal{W}_1 of \mathcal{W} , $\mathcal{W}_0^* q^* = \mathcal{W}_1^* q^*$.

By lemma 17.20, there is a finite correspondence $\widetilde{\mathcal{W}}$ so that the following diagram commutes:




Since s_0 and s_1 are both inverses to the projection $p : X \times \mathbb{A}^1 \rightarrow X$, we have $s_0^* p^* = s_1^* p^*$ by 17.13. Since higher Chow groups are homotopy invariant, p^* is an isomorphism and we get $s_0^* = s_1^*$. Since X and Y' are affine, and $\mathcal{W}_i = \widetilde{\mathcal{W}} \circ s_i$, we may apply 17.13 again to get

$$\mathcal{W}_0^* = s_0^* \widetilde{\mathcal{W}}^* = s_1^* \widetilde{\mathcal{W}}^* = \mathcal{W}_1^*. \quad \square$$

Recall from 2.25 that two correspondences \mathcal{W}_0 and \mathcal{W}_1 from X to Y are said to be \mathbb{A}^1 -homotopic, written $\mathcal{W}_0 \simeq \mathcal{W}_1$, if they are the restrictions of an element of $Cor(X \times \mathbb{A}^1, Y)$ along $X \times 0$ and $X \times 1$.

LEMMA 17.20. *Let \mathcal{W} be a finite correspondence between a smooth affine scheme X and a smooth Y . If $q : Y' \rightarrow Y$ is a vector bundle torsor, then any two lifts \mathcal{W}_0 and \mathcal{W}_1 are \mathbb{A}^1 -homotopic.*

PROOF. Let V be the image of the union of the supports of \mathcal{W}_0 and \mathcal{W}_1 in $X \times Y$, and let V' denote the fiber product of V and Y' over Y ; $p : V' \rightarrow V$ is a vector bundle torsor. Since X is affine and the induced map $V \rightarrow X$ is finite, V is affine too. Hence $p : V' \rightarrow V$ is a vector bundle. Fix a section $s : V \rightarrow V'$.



Clearly, p is an \mathbb{A}^1 -homotopy equivalence (in the sense of 2.25) with inverse s , that is, sp is \mathbb{A}^1 -homotopic to the identity.

Both \mathcal{W}_0 and \mathcal{W}_1 induce correspondences $\widetilde{\mathcal{W}}_0$ and $\widetilde{\mathcal{W}}_1$ from X to V' . Now the composition $g \circ (\widetilde{\mathcal{W}}_i \times \mathbb{A}^1) \in Cor(X \times \mathbb{A}^1, V')$ is an \mathbb{A}^1 -homotopy from $sp\widetilde{\mathcal{W}}_i$ to $\widetilde{\mathcal{W}}_i$, for $i = 0, 1$. Since $p\widetilde{\mathcal{W}}_0 = p\widetilde{\mathcal{W}}_1$, we have

$$\widetilde{\mathcal{W}}_0 \simeq sp\widetilde{\mathcal{W}}_0 = sp\widetilde{\mathcal{W}}_1 \simeq \widetilde{\mathcal{W}}_1.$$

Since \mathcal{W}_i is the composition of $\widetilde{\mathcal{W}}_i$ with the map $V' \rightarrow Y$, \mathcal{W}_0 is \mathbb{A}^1 -homotopic to \mathcal{W}_1 . \square

At last, we have the tools to show that the higher Chow groups are presheaves with transfers, i.e., functors on the category Cor_k of smooth separated schemes over k .

THEOREM 17.21. *The maps \mathcal{W}^* defined in 17.17 give the higher Chow groups $CH^i(-, m)$ the structure of presheaves with transfers.*

That is, for any two finite correspondences \mathcal{W}_1 and \mathcal{W}_2 from X to Y and from Y to Z , respectively, and for all $\alpha \in CH^i(Z, m)$:

$$\mathcal{W}_1^*(\mathcal{W}_2^*(\alpha)) = (\mathcal{W}_2 \circ \mathcal{W}_1)^*(\alpha).$$

In particular, if $f_1 : X \rightarrow Y$ and $f_2 : Y \rightarrow Z$ are morphisms, then $(f_2 \circ f_1)^ = f_1^* f_2^*$.*

PROOF. By 17.16, there is a commutative diagram in Cor_k of the form

$$\begin{array}{ccccc} X' & \xrightarrow{\mathcal{W}'_1} & Y' & \xrightarrow{\mathcal{W}'_2} & Z' \\ p \downarrow & & q \downarrow & & r \downarrow \\ X & \xrightarrow{\mathcal{W}_1} & Y & \xrightarrow{\mathcal{W}_2} & Z \end{array}$$

where the vertical maps are affine vector bundle torsors. By 17.13, we have $\mathcal{W}'_1^* \mathcal{W}'_2^* = (\mathcal{W}'_2 \circ \mathcal{W}'_1)^*$. Since the definitions of \mathcal{W}_i^* and $(\mathcal{W}_2 \circ \mathcal{W}_1)^*$ are independent of the choices by 17.19, the statement now follows from an unwinding of 17.17:

$$\mathcal{W}_1^* \mathcal{W}_2^* = (p^*)^{-1} \mathcal{W}'_1^* q^* (q^*)^{-1} \mathcal{W}'_2^* r^* = (p^*)^{-1} (\mathcal{W}'_2 \circ \mathcal{W}'_1)^* r^* = (\mathcal{W}_2 \circ \mathcal{W}_1)^*. \quad \square$$

APPENDIX 17A

Cycle maps

If \mathscr{W} is a finite correspondence from X to Y , we can define a map \mathscr{W}^* from “good” cycles on Y to cycles on X . The formula is to pull the cycle back to $X \times Y$, intersect it with \mathscr{W} , and push forward to X . In this appendix, we will make this precise, in 17A.8. First we must explain what makes a cycle “good”.

DEFINITION 17A.1. Two subvarieties Z_1 and Z_2 of X are said to **intersect properly** if every component of $Z_1 \cap Z_2$ has codimension $\text{codim} Z_1 + \text{codim} Z_2$ in X . This is vacuously true if $Z_1 \cap Z_2 = \emptyset$.

If the ambient variety X is regular, the intersection cycle $Z_1 \cdot Z_2$ is defined to be the sum $\sum n_j [W_j]$, where the indexing is over the irreducible components W_j of $Z_1 \cap Z_2$, and the n_j are their (local) intersection multiplicities. Following Serre [Ser65], the multiplicity n_j is defined as follows. If A is the local ring of X at the generic point of W_j , and I_i are the ideals of A defining Z_i , then

$$n_j = \sum_i (-1)^i \text{length Tor}_i^A(A/I_1, A/I_2).$$

If X is not regular, the multiplicity will make sense only when at most finitely many Tor-terms are non-zero.

We say that two equidimensional cycles $\mathscr{V} = \sum m_i V_i$ and $\mathscr{W} = \sum n_j W_j$ intersect properly if each V_i and W_j intersect properly. In this case, the intersection cycle $\mathscr{V} \cdot \mathscr{W}$ is defined to be $\sum m_j n_j (V_i \cdot W_j)$.

EXERCISE 17A.2. Let $\mathscr{V}_1, \mathscr{V}_2$ and \mathscr{V}_3 be three cycles on a smooth scheme X . Show that $(\mathscr{V}_1 \cdot \mathscr{V}_2) \cdot \mathscr{V}_3 = \mathscr{V}_1 \cdot (\mathscr{V}_2 \cdot \mathscr{V}_3)$ whenever both sides are defined. (This is proven in [Ser65, V-24].)

DEFINITION 17A.3. Suppose that $f : X \rightarrow Y$ is a morphism with X and Y regular, and that \mathscr{Z} is a codimension i cycle on Y . We say that $f^*(\mathscr{Z})$ is **defined** if each component of $f^{-1}(\text{Supp}(\mathscr{Z}))$ has codimension $\geq i$ in X . As in [Ser65, V-28], we define the cycle $f^*(\mathscr{Z})$ to be $\Gamma_f \cdot (X \times \mathscr{Z})$ (see 17A.1), identifying the graph Γ_f with X .

As noted in [Ser65, V-29], the intersection cycle makes sense even if X is not regular, since the multiplicities may be computed over Y by flat base change for Tor (see [Wei94, 3.2.9]).

EXAMPLE 17A.4. If f is flat and $\mathscr{Z} = [V]$, then $f^*(\mathscr{Z})$ is the cycle associated to the scheme $f^{-1}(V)$. If X is a subvariety of Y , then the cycle $f^*(\mathscr{Z})$ on X is

the same as the cycle $X \cdot \mathcal{Y}$ considered as a cycle on X . If $X \hookrightarrow Y$ is a regular embedding, the coefficients of $f^*(\mathcal{Y})$ agree with the intersection multiplicities defined in [Ful84, 7.1.2].

REMARK 17A.5. Here is a variant of definition 17A.3 we will need in the next lecture. Suppose that S and T are smooth, that X is a scheme of finite type, and that $f : X \times S \rightarrow X \times T$ is a morphism over X . If V is a codimension i cycle in $X \times T$, we say that $f^*(V)$ is defined if every component of $f^{-1}(V)$ has codimension i in $X \times S$. (It is not hard to see that they have codimension $\leq i$.)

As in 17A.3, $f^*(V)$ is defined to be the intersection product $\Gamma_f \cdot (S \times V)$ of 17A.1, where Γ_f is the image of the graph embedding of $X \times S$ into $X \times S \times T$. The Tor formula of 17A.1 makes sense because the inclusion $\Gamma_f \subset X \times S \times T$ is locally defined by a regular sequence, and hence has finite Tor dimension.

DEFINITION 17A.6. Let $f : Y' \rightarrow Y$ be a morphism of smooth varieties and \mathcal{W} a cycle on Y' . We say that a cycle \mathcal{Y} on Y is in **good position** for \mathcal{W} (relative to f) if the cycle $f^*(\mathcal{Y})$ is defined, and intersects \mathcal{W} properly on Y' . If \mathcal{Y} is in good position for \mathcal{W} , the intersection product $\mathcal{W} \cdot f^*\mathcal{Y}$ is defined (see 17A.1). If the map f is flat, the cycle $f^*(\mathcal{Y})$ is always defined.

Let W be an irreducible subvariety of Y' and let w be the composition $W \rightarrow Y' \rightarrow Y$. By 17A.1 and 17A.3, a codimension i cycle \mathcal{Y} is in good position for \mathcal{W} if and only if $\text{codim}_W w^{-1}(\text{Supp}(\mathcal{Y})) \geq i$, that is, if $w^*(\mathcal{Y})$ is defined.

As a special case, we will say that a cycle \mathcal{Y} is in good position for a finite correspondence \mathcal{W} from X to Y if \mathcal{Y} is in good position for the cycle underlying \mathcal{W} , relative to the projection $X \times Y \rightarrow Y$.

REMARK 17A.7. Let $f : X \rightarrow Y$ be a morphism of smooth varieties and let \mathcal{Z} be a cycle on X , supported on a closed subscheme Z so that the composition $Z \rightarrow X \rightarrow Y$ is a proper map. It is clear that $f_*(\mathcal{Z})$ is well-defined even though f is not proper.

DEFINITION 17A.8. Let \mathcal{W} be a finite correspondence between two smooth schemes X and Y . For every cycle \mathcal{Y} on Y in good position for \mathcal{W} , we define

$$\mathcal{W}^*(\mathcal{Y}) = \pi_*(\mathcal{W} \cdot f^*\mathcal{Y}),$$

where $f : X \times Y \rightarrow Y$ and $\pi : X \times Y \rightarrow X$ are the canonical projections. The intersection and the push-forward are well-defined by 17A.6 and 17A.7. The map \mathcal{W}^* induces the transfer map for Chow groups, see 17.11 and 17.17.

For any smooth T , $\mathcal{W} \times T$ is a finite correspondence from $X \times T$ to $Y \times T$ over T . By abuse of notation, we shall also write \mathcal{W}^* for $(\mathcal{W} \times T)^*$.

EXAMPLE 17A.9. We can now reinterpret the composition of correspondences. If \mathcal{W}_1 and \mathcal{W}_2 are finite correspondences from X to Y and from Y to Z , respectively, we have:

$$\mathcal{W}_2 \circ \mathcal{W}_1 = (\mathcal{W}_1 \times Z)^*(\mathcal{W}_2) = (X \times \mathcal{W}_2)^*(\mathcal{W}_1).$$

Here are two formulas which are useful in the study of \mathcal{W}^* .

LEMMA 17A.10. *Consider the following diagram of varieties*

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{g} & Y \end{array}$$

where the square is fiber and both X and Y are smooth. Let \mathcal{X} be a cycle on X whose support is finite over Y and for which $(g')^* \mathcal{X}$ is defined. Then $g^* f_* \mathcal{X}$ is defined and $g^* f_* \mathcal{X} = f'_*(g')^* \mathcal{X}$.

PROOF. If V is a component of \mathcal{X} , then the map $f : V \rightarrow f(V)$ is finite. Hence $f' : (g')^{-1}(V) \rightarrow g^{-1}(f(V))$ is finite too, so that $\text{codim}(g')^{-1}(V) = \text{codim} g^{-1}(f(V))$. By hypothesis, $\text{codim}(g')^{-1}(V) \geq i$, which proves that $g^* f_* \mathcal{X}$ is defined. The equality now follows from [Ful75, 2.2(4)]. \square

LEMMA 17A.11 (Projection Formula). *Let $f : X \rightarrow Y$ be a morphism of smooth schemes. Suppose given a cycle \mathcal{X} on X , whose support is finite over Y , and a cycle \mathcal{Y} on Y which is in good position for \mathcal{X} (see 17A.6). Then $f_* \mathcal{X}$ and \mathcal{Y} intersect properly, and the projection formula holds:*

$$f_*(\mathcal{X} \cdot f^* \mathcal{Y}) = f_* \mathcal{X} \cdot \mathcal{Y}.$$

PROOF. Since the restriction of f to the support of \mathcal{X} is finite, it is clear that $f_*(\mathcal{X})$ and \mathcal{Y} intersect properly too. The result is now a consequence of the basic identity 2.2(2) of [Ful75], or [Ser65, V-30]. \square

EXERCISE 17A.12. Let i be the inclusion of a closed subvariety W in a smooth scheme X and let $f : X \rightarrow Y$ be a map of smooth schemes. Prove that if \mathcal{Y} is a cycle on Y so that both $f^* \mathcal{Y}$ and $(fi)^*(\mathcal{Y})$ are defined, then $i_*(fi)^*(\mathcal{Y}) = W \cdot f^* \mathcal{Y}$. *Hint:* Use [Ser65, V-30] or [Ful75, 2.2(2)].

Recall from 1A.10 that if $V \rightarrow Y$ is a morphism with Y regular, then the pullback \mathcal{Z}_V of a relative cycle \mathcal{Z} in $T \times Y$ is a well defined cycle on $T \times V$ with integer coefficients.

LEMMA 17A.13. *Let T and Y be regular and let \mathcal{Z} be a cycle in $T \times Y$ which is dominant equidimensional over Y . If $f : V \rightarrow Y$ is a morphism, then the pullback \mathcal{Z}_V agrees with the pullback cycle $(f \times T)^*(\mathcal{Z})$.*

PROOF. Note that \mathcal{Z} is a relative cycle by 1A.6, so that \mathcal{Z}_V is defined. Its coefficients are characterized by the equalities $(\mathcal{Z}_V)_v = \mathcal{Z}_{f(v)}$ for every $v \in V$. By [ReiCh, 3.5.8 and 3.5.9], the coefficients of \mathcal{Z}_V are the same as the multiplicities in 17A.1, i.e., the coefficients of $(f \times T)^*(\mathcal{Z})$ given by 17A.3. \square


THEOREM 17A.14. *Let \mathcal{W}_1 and \mathcal{W}_2 be two finite correspondences from X to Y and from Y to Z , respectively. Suppose that \mathcal{Z} is a cycle on Z which is in good position with respect to both \mathcal{W}_2 and $\mathcal{W}_2 \circ \mathcal{W}_1$. Then*

$$(\mathcal{W}_2 \circ \mathcal{W}_1)^*(\mathcal{Z}) = \mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{Z})).$$

The term $\mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{Z}))$ makes sense by the following lemma.

LEMMA 17A.15. *Let \mathcal{Z} be in good position for \mathcal{W}_2 and $\mathcal{W}_2 \circ \mathcal{W}_1$. Then $\mathcal{W}_2^*(\mathcal{Z})$ is in good position with respect to \mathcal{W}_1 .*

PROOF. We may assume that the correspondences are elementary, i.e., \mathcal{W}_1 and \mathcal{W}_2 are subvarieties W_1 and W_2 of $X \times Y$, and $Y \times Z$, respectively. In this spirit, we will write $W_2 \circ W_1$ for the subvariety of $X \times Z$ which is the support of the composition of correspondences, $\mathcal{W}_2 \circ \mathcal{W}_1$. Consider the following diagram.



By hypothesis, $\text{codim } d^{-1}(\mathcal{Z}) \geq \text{codim } \mathcal{Z}$ and $\text{codim } c^{-1}(\mathcal{Z}) \geq \text{codim } \mathcal{Z}$.

We claim that $\text{codim } b^{-1}pd^{-1}\mathcal{Z} \geq \text{codim } \mathcal{Z}$. Since the central square is cartesian, $b^{-1}p = qe^{-1}$. Since q is finite, this yields

$$\text{codim } b^{-1}pd^{-1}\mathcal{Z} = \text{codim } qe^{-1}d^{-1}\mathcal{Z} = \text{codim } e^{-1}d^{-1}\mathcal{Z}.$$

But $e^{-1}d^{-1} = u^{-1}c^{-1}$, and u is finite, so:

$$\text{codim } e^{-1}d^{-1}\mathcal{Z} = \text{codim } u^{-1}c^{-1}\mathcal{Z} = \text{codim } c^{-1}\mathcal{Z}.$$

But $\text{codim } c^{-1}\mathcal{Z} \geq \text{codim } \mathcal{Z}$ by hypothesis, as claimed. □

PROOF OF 17A.14. The right side is defined by 17A.15. We will follow the notation established in figure 17A.1, where we have omitted the factor Δ^n in every entry to simplify notation. Note that the central square is cartesian.

By definition 17.7, we have

$$\mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{Z})) = r_*(\mathcal{W}_1 \cdot b^*(p_*(\mathcal{W}_2 \cdot d^*\mathcal{Z}))).$$

Since the central square is cartesian, we have $b^*p_* = q_*e^*$ by 17A.10. Since the pullback e^* is a ring homomorphism, we have

$$b^*(p_*(\mathcal{W}_2 \cdot d^*\mathcal{Z})) = q_*(e^*(\mathcal{W}_2 \cdot d^*\mathcal{Z})) = q_*(e^*(\mathcal{W}_2) \cdot e^*d^*\mathcal{Z}).$$




FIGURE 17A.1. Composition of correspondences

Consider the two cycles $\mathcal{X} = e^*(\mathcal{W}_2) \cdot (de)^*(\mathcal{L})$ and $\mathcal{Y} = \mathcal{W}_1$ and the function q . The intersection $\mathcal{X} \cdot q^*\mathcal{Y} = e^*(\mathcal{W}_2) \cdot (de)^*(\mathcal{L}) \cdot q^*(\mathcal{W}_1)$ is proper because \mathcal{L} is in good position with respect to $\mathcal{W}_2 \circ \mathcal{W}_1$. Therefore the conditions for 17A.11 are satisfied, and the projection formula yields $\mathcal{Y} \cdot q_*\mathcal{X} = q_*(q^*\mathcal{Y} \cdot \mathcal{X})$, i.e.,

$$\mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{L})) = r_*q_*(q^*(\mathcal{W}_1) \cdot (e^*(\mathcal{W}_2) \cdot e^*d^*\mathcal{L})).$$

Since the push-forward and pullback are functorial, we have $r_*q_* = v_*u_*$ and $e^*d^* = u^*c^*$. Our cycle then becomes

$$v_*u_*(q^*(\mathcal{W}_1) \cdot e^*(\mathcal{W}_2) \cdot u^*c^*\mathcal{L}).$$

We may use the projection formula (17A.11) once again, this time for u^* , with $\mathcal{X} = q^*(\mathcal{W}_1) \cdot e^*(\mathcal{W}_2)$ and $\mathcal{Y} = c^*\mathcal{L}$ (the conditions are satisfied by the same argument we used above). This yields $u_*(\mathcal{X} \cdot u^*\mathcal{Y}) = (u_*\mathcal{X}) \cdot \mathcal{Y}$, i.e.,

$$\mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{L})) = v_*(u_*(q^*(\mathcal{W}_1) \cdot e^*(\mathcal{W}_2)) \cdot c^*\mathcal{L}).$$

Since the composition of \mathcal{W}_1 and \mathcal{W}_2 as correspondences is exactly $u_*(q^*(\mathcal{W}_1) \cdot e^*(\mathcal{W}_2))$, the last equation becomes

$$\mathcal{W}_1^*(\mathcal{W}_2^*(\mathcal{L})) = v_*((\mathcal{W}_2 \circ \mathcal{W}_1) \cdot c^*\mathcal{L}) = (\mathcal{W}_2 \circ \mathcal{W}_1)^*(\mathcal{L}).$$

This concludes the proof of 17A.14. □

Higher Chow groups and equidimensional cycles

The next step in the proof of theorem 19.1 (that motivic cohomology and higher Chow groups agree) is the reduction to equidimensional cycles. The main references for this lecture are [HighCh] and [FS02].

DEFINITION 18.1. For an equidimensional X , and $i \leq \dim X$, we write $z_{equi}^i(X, m)$ for $z_{equi}(X, \dim X - i)(\Delta^m)$, the free abelian group generated by all codimension i subvarieties on $X \times \Delta^m$ which are dominant and equidimensional over Δ^m (of relative dimension $\dim X - i$). We write $z_{equi}^i(X, \bullet)$ and $z_{equi}^i(X, *)$ for the simplicial abelian group $m \mapsto z_{equi}^i(X, m)$ and its associated chain complex, respectively.

By 17.2, $z_{equi}^i(X, m)$ is a subgroup of $z^i(X, m)$ and $z_{equi}^i(X, \bullet)$ is a simplicial subgroup of $z^i(X, \bullet)$.

EXAMPLE 18.2. The inclusion $z_{equi}^i(X, *) \subset z^i(X, *)$ will not be a quasi-isomorphism in general. Indeed, if $i > d$ then $z_{equi}^i(X, m) = 0$ while $z^i(X, m)$ is not generally zero. For example, consider $X = \text{Spec } k$. If $i > 0$ we have $z_{equi}^i(\text{Spec } k, *) = 0$. In contrast, $z^i(\text{Spec } k, i)$ is the group of points on Δ^i which do not lie on any proper face. We will see in 19.7 that $H_i z^i(\text{Spec } k, *) = H^{i,i}(\text{Spec } k) = K_i^M(k)$.

THEOREM 18.3. (Suslin [HighCh, 2.1]) *Let X be an equidimensional affine scheme of finite type over k , then the inclusion map:*

$$z_{equi}^i(X, *) \hookrightarrow z^i(X, *)$$

is a quasi-isomorphism for $i \leq \dim X$.

COROLLARY 18.4. *Let X be an affine variety, then for all $i \geq 0$*

$$CH^i(X, m) = H_m(z_{equi}^i(X \times \mathbb{A}^i, *)).$$

*In particular, $CH^i(\text{Spec } k, m) = H_m(z_{equi}^i(\mathbb{A}^i, *))$.*

PROOF. This is an immediate corollary of 18.3, definition 17.1 and the homotopy invariance of the higher Chow groups; see 17.4. \square

COROLLARY 18.5. *Let X be an equidimensional quasi-projective scheme over a field k which admits resolution of singularities. For all $i \leq \dim X$, the natural*

inclusion $z_{equi}^i(X, *) \hookrightarrow z^i(X, *)$ is a quasi-isomorphism, i.e., it induces isomorphisms

$$H_m z_{equi}^i(X, *) \xrightarrow{\cong} H_m z^i(X, *) = CH^i(X, m).$$

PROOF. If U is affine, $z_{equi}^i(U, *) \hookrightarrow z^i(U, *)$ is a quasi-isomorphism by 18.3. We proceed by induction on $\dim X$. Let U be a dense open affine subscheme of X with complement Z of codimension 1. The commutative diagram

$$\begin{array}{ccccc} z_{equi}^{i-1}(Z, *) & \longrightarrow & z_{equi}^i(X, *) & \longrightarrow & z_{equi}^i(U, *) \\ \downarrow \simeq & & \downarrow & & \downarrow \simeq \\ z^{i-1}(Z, *) & \longrightarrow & z^i(X, *) & \longrightarrow & z^i(U, *) \end{array}$$

becomes a morphism of triangles by 16.19 and the Localization Theorem for higher Chow groups (see 17.4). The result follows from the 5-lemma. \square

We need 18.9, 18.12, and 18.14 to prove theorem 18.3. All of their proofs rely on a technical theorem 18A.1, which will be proven in the appendix.

We begin by introducing some auxiliary notions. Let X be a scheme over S .

DEFINITION 18.6. An **N-skeletal map** φ over X , relative to $X \rightarrow S$, is a collection $\{\varphi_n : X \times \Delta^n \rightarrow X \times \Delta^n\}_{n=0}^N$ of S -morphisms, such that φ_0 is the identity 1_X and for every face map $\partial_j : \Delta^{n-1} \rightarrow \Delta^n$ with $n \leq N$ the following diagram commutes.

$$\begin{array}{ccc} X \times \Delta^{n-1} & \xrightarrow{\varphi_{n-1}} & X \times \Delta^{n-1} \\ 1_X \times \partial_j \downarrow & & \downarrow 1_X \times \partial_j \\ X \times \Delta^n & \xrightarrow{\varphi_n} & X \times \Delta^n \end{array}$$

Note that φ_N determines φ_n for all $n < N$. When $S = X$, we shall just call φ an n -skeletal map over X .

The condition that an $(N-1)$ -skeletal map over X can be extended to an N -skeletal map is a form of the homotopy extension property, and follows from the Chinese Remainder Theorem when X is affine.

For example, a 1-skeletal map over $X = \text{Spec } R$ (relative to $S = X$) is determined by a polynomial $f \in R[t]$ such that $f(0) = 0$ and $f(1) = 1$; φ_1 is Spec of the R -algebra map $R[t] \rightarrow R[t]$ sending t to f .

DEFINITION 18.7. Given an N -skeletal map φ over X and $n \leq N$, we define $\varphi z^i(X, n)$ to be the subgroup of $z^i(X, n)$ generated by all V in $X \times \Delta^n$ such that $\varphi_n^*(V)$ is defined (in the sense of 17A.5) and is in $z^i(X, n)$. If $n > N$ we set $\varphi z^i(X, n) = 0$. In other words, $\varphi z^i(X, n)$ is the group of cycles in $X \times \mathbb{A}^n$ which intersect all the faces properly and whose pullbacks along φ_n intersect all the faces properly.

By definition 18.6 we know that the face map $\partial_j : z^i(X, n) \rightarrow z^i(X, n-1)$ sends $\varphi z^i(X, n)$ to $\varphi z^i(X, n-1)$. Thus $\varphi z^i(X, *)$ is a chain subcomplex of $z^i(X, *)$.

Moreover it follows from 18.6 that the φ_n^* assemble to define a chain map $\varphi^* : \varphi z^i(X, *) \rightarrow z^i(X, *)$.

Similarly, we can define $\varphi z_{equi}^i(X, n)$ to be the subgroup of $z_{equi}^i(X, n)$ generated by all V such that $\varphi_n^*(V)$ is defined and is in $z_{equi}^i(X, n)$. The same argument shows that $\varphi z_{equi}^i(X, *)$ is a subcomplex of $z_{equi}^i(X, *)$ and that the φ_n form a chain map $\varphi^* : \varphi z_{equi}^i(X, *) \rightarrow z_{equi}^i(X, *)$.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \varphi z^i(X, 1) & \longrightarrow & \varphi z^i(X, 0) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & i \downarrow & \varphi_1^* \downarrow & \downarrow = & & \\
 z^i(X, 2) & \longrightarrow & z^i(X, 1) & \longrightarrow & z^i(X, 0) & \longrightarrow & 0.
 \end{array}$$

FIGURE 18.1. A 1-skeletal map φ and its chain map φ^* .

EXAMPLE 18.8. If $N = 1$, and $\alpha \in k - \{0, 1\}$, the subvariety $V = X \times \{\alpha\}$ of $X \times \mathbb{A}^1$ is in $z^1(X, 1)$ but not $z_{equi}^1(X, 1)$. If $X = \text{Spec} R$, fix $r \in R$ and let $\varphi_1 : X \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1$ be the 1-skeletal map defined by the R -algebra map $R[t] \rightarrow R[t]$ sending t to $f(t) = t + r(t^2 - t)$. The condition that $\varphi_1^*(V)$ is in $z_{equi}^1(X, 1)$, i.e., dominant and equidimensional over \mathbb{A}^1 , is equivalent to the condition that the map $r : X \rightarrow \mathbb{A}^1$ is equidimensional, i.e., that $r - \beta$ is nonzero in the domain R for all $\beta \in k$. Indeed, the fiber of $\varphi_1^{-1}(V)$ over $t \neq 0, 1$ is supported on $R/(r - (\alpha - t/t^2 - t))$, and is empty if $t = 0, 1$. Such r always exists when $\dim X \geq 1$.

LEMMA 18.9. (See [HighCh, 2.8]) Let C_* be a finitely generated subcomplex in $z^i(X, *)$ with $i \leq \dim X$. Choose N so that $C_n = 0$ for $n > N$. Then there is an N -skeletal map φ over X such that $C_* \subseteq \varphi z^i(X, *)$, and the chain map $\varphi^* : \varphi z^i(X, *) \rightarrow z^i(X, *)$ satisfies

$$\varphi^* C_* \subseteq z_{equi}^i(X, *).$$

PROOF. Suppose that C_n is generated by $\{V_n^k\} \subseteq z^i(X, n)$. Set $d = \dim X - i$ and note that $d \geq 0$ since $i \leq \dim X$. Then $V_n = \cup V_n^k$ is closed in $X \times \Delta^n$ of dimension $n + d$.

We proceed by induction on N . Since N is finite, we may assume that the $\partial_j(V_n^k)$ are supported in V_{n-1} . Inductively, we may suppose that we have constructed an $(N - 1)$ -skeletal map $\{\varphi_n\}$ such that the fibers of the projections $\varphi_n^{-1}(V_n) \rightarrow \Delta^n$ have dimension $\leq d$. Let $\partial \Delta^N$ be the union of the faces Δ^N . The compatibility granted by definition 18.6 implies that these maps fit together to form a map from $X \times \partial \Delta^N$ to itself such that the fibers of $\varphi^{-1}(X \times \partial \Delta^N) \cap V_N \rightarrow \partial \Delta^N$ have dimension $\leq d$. By Generic Equidimensionality 18A.1, this map extends to an N -skeletal map $\varphi_N : X \times \Delta^N \rightarrow X \times \Delta^N$ over X such that the fibers of $\varphi_N^{-1}(V_N) \rightarrow \mathbb{A}^N$ have dimension $\leq d$. Because each component W of $\varphi^{-1}(V_n^k)$ satisfies the inequality $\dim W \leq n + d = \dim V_n^k$, each cycle $\varphi_n^*(V_n^k)$ is defined and

lies in $z_{equi}^i(X, n)$. Since C_n is generated by the V_n^k , it lies in $\varphi^i(X, n)$ and satisfies $\varphi^*(C_n) \subset z_{equi}^i(X, n)$. \square

DEFINITION 18.10. Let φ^0 and φ^1 be N -skeletal maps over X . An **N -skeletal homotopy Φ** between φ^0 and φ^1 is an N -skeletal map $\{\Phi_n : X \times \Delta^n \times \mathbb{A}^1 \rightarrow X \times \Delta^n \times \mathbb{A}^1\}_{n=0}^N$ over $X \times \mathbb{A}^1$ relative to the projection $X \times \mathbb{A}^1 \rightarrow X$, which is compatible with the φ^j in the sense that the following diagram commutes for every n .

$$\begin{array}{ccccc} X \times \Delta^n & \xrightarrow{i_0} & X \times \Delta^n \times \mathbb{A}^1 & \xleftarrow{i_1} & X \times \Delta^n \\ \varphi_n^0 \downarrow & & \downarrow \Phi_n & & \downarrow \varphi_n^1 \\ X \times \Delta^n & \xrightarrow{i_0} & X \times \Delta^n \times \mathbb{A}^1 & \xleftarrow{i_1} & X \times \Delta^n \end{array}$$

Recall from 2.17 that the simplicial decomposition of $\Delta^n \times \mathbb{A}^1$ is given by isomorphisms $\theta_j : \Delta^{n+1} \rightarrow \Delta^n \times \mathbb{A}^1$, $j = 0, \dots, n$. Each θ_j identifies the subgroup $z^i(X, n+1)$ of cycles in $X \times \Delta^{n+1}$ with a subgroup of cycles in $X \times \Delta^n \times \mathbb{A}^1$.

The subgroup $\Phi z^i(X, n)$ of $z^i(X, n)$ is defined to be the subgroup generated by all V in $X \times \Delta^n$ such that: (a) $(\varphi^0)^*(V)$ and $(\varphi^1)^*(V)$ are defined and in $z^i(X, n)$; (b) each $\Phi_n^*(V \times \mathbb{A}^1)$ is defined (see 17A.5); and (c) each isomorphism θ_j identifies $\Phi_n^*(V \times \mathbb{A}^1)$ with an element of $z^i(X, n+1)$. As in definition 18.7, $\Phi z^i(X, *)$ is a subcomplex of $z^i(X, *)$. In fact, Φz^i lies in $(\varphi^0 z^i) \cap (\varphi^1 z^i)$.

We define the subgroup $\Phi z_{equi}^i(X, n)$ of $z_{equi}^i(X, n)$ similarly, replacing z^i with z_{equi}^i in the definition of $\Phi z^i(X, n)$.

LEMMA 18.11. *If Φ is an N -skeletal homotopy between φ^0 and φ^1 , then the maps $(\varphi^0)^*$ and $(\varphi^1)^*$ are chain homotopic, both from $\Phi z^i(X, *)$ to $z^i(X, *)$ and from $\Phi z_{equi}^i(X, *)$ to $z_{equi}^i(X, *)$.*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Phi z^i(X, 2) & \xrightarrow{\partial} & \Phi z^i(X, 1) & \xrightarrow{\partial} & \Phi z^i(X, 0) \\ \downarrow & & \downarrow & \searrow h & \downarrow & \searrow h & \downarrow \\ \downarrow & & \downarrow \varphi_2^1 & & \downarrow \varphi_1^1 & & \downarrow \varphi_0^1 \\ \downarrow & & \downarrow \varphi_2^0 & & \downarrow \varphi_1^0 & & \downarrow \varphi_0^0 \\ z^i(X, 3) & \xrightarrow{\partial} & z^i(X, 2) & \xrightarrow{\partial} & z^i(X, 1) & \xrightarrow{\partial} & z^i(X, 0) \\ & & \downarrow \varphi_2^0 & & \downarrow \varphi_1^0 & & \downarrow \varphi_0^0 \\ & & & & & & \varphi_0^0 = 1_X \end{array}$$

FIGURE 18.2. The chain homotopy between id and φ^* when $N = 2$.

PROOF. For $0 \leq j \leq n$, let h_j denote the composite

$$X \times \Delta^{n+1} \xrightarrow{1_X \times \theta_j} X \times \Delta^n \times \mathbb{A}^1 \xrightarrow{\Phi_n} X \times \Delta^n \times \mathbb{A}^1 \xrightarrow{pr} X \times \Delta^n,$$

where pr is the projection. That is, for V in $\Phi z^i(X, n)$ we define

$$h_j^*[V] = (\Phi_n \circ (1_X \times \theta_j))^*[V \times \mathbb{A}^1] \in z^i(X, n+1).$$

The h_j^* form a simplicial homotopy (see [Wei94, 8.3.11]) from $\partial_0 h_0^* = (\varphi^1)^*$ to $\partial_{n+1} h_n^* = (\varphi^0)^*$. Hence their alternating sum $h = \sum (-1)^j h_j^*$ satisfies $h\partial + \partial h = (\varphi^1)^* - (\varphi^0)^*$. (This is illustrated in figure 18.2 when $N = 2$.) \square

PROPOSITION 18.12. *Let φ be an N -skeletal map, and $\{V_n^k\}$ a finite set of varieties in $\varphi z^i(X, n)$, $n \leq N$. Then there exists an N -skeletal homotopy Φ between φ and the identity map, such that each V_n^k lies in $\Phi z^i(X, n)$.*

If the $\{V_n^k\}$ lie in $\varphi z_{equi}^i(X, n)$, then the $\Phi_n^(V_n^k \times \mathbb{A}^1)$ lie in $z_{equi}^i(X \times \mathbb{A}^1, n)$.*

For the construction of Φ , we may assume without loss of generality that the set of V_n^k is closed under taking components of restrictions to faces.

PROOF. Set $d = \dim(X) - i$, and let $\partial(\Delta^n \times \mathbb{A}^1)$ denote the union of $(\partial\Delta^n) \times \mathbb{A}^1$ and $\Delta^n \times \{0, 1\}$. As in the proof of 18.9, we shall construct an N -skeletal homotopy Φ by induction on N satisfying the “fiber condition” that (for each k and $n \leq N$) the fibers of the projections $\Phi_n^{-1}(V_n^k \times \mathbb{A}^1) \rightarrow \Delta^n \times \mathbb{A}^1$ have dimension $\leq d$ over all points not in $\partial(\Delta^n \times \mathbb{A}^1)$.

Inductively, we are given Φ_n ($n < N$) forming an $(N - 1)$ -skeletal map which satisfies the fiber condition. The compatibility with the faces of Δ^N and with i_0, i_1 granted by definition 18.10 implies that the Φ_n and φ_N fit together to form a map $\partial\Phi_N$ from $X \times \partial(\Delta^N \times \mathbb{A}^1)$ to itself. By Generic Equidimensionality 18A.1, with $\mathbb{A}^n = \Delta^N \times \mathbb{A}^1$, this map extends to a map Φ_N from $X \times \Delta^N \times \mathbb{A}^1$ to itself which extends $\partial\Phi_N$ (i.e., Φ_N is an N -skeletal homotopy from the identity to φ over X), such that the fibers of $\Phi_N^{-1}(V_N^k \times \mathbb{A}^1) \rightarrow \Delta^N \times \mathbb{A}^1$ have dimension $\leq d$ over all points of $\Delta^N \times \mathbb{A}^1$ not on $\partial(\Delta^N \times \mathbb{A}^1)$.

To complete the proof of 18.12, we need to show that each $\Phi_N^*(V_N^k)$ is defined and that each isomorphism θ_j identifies them with elements in $z^i(X, N + 1)$ (resp., in $z_{equi}^i(X, N + 1)$ when $V_N^k \in \varphi z_{equi}^i(X, n)$). Set $W = \bigcup_k \Phi_N^{-1}(V_N^k \times \mathbb{A}^1)$.

Because the V_N^k belong to $\varphi z^i(X, N)$ (resp., to $z_{equi}^i(X, N)$), the part of W lying over $\Delta^N \times \{0, 1\}$ has dimension $d + N$ (resp., is equidimensional). The inductive hypothesis implies that the part of W lying over $\partial(\Delta^N \times \mathbb{A}^1)$ has dimension $\leq d + N$ (resp., is equidimensional). Let $F \subseteq \Delta^N \times \mathbb{A}^1$ correspond to a face of Δ^{N+1} under one of the isomorphisms θ_j . The fiber condition on Φ_n implies that the part of W lying over F but not over $\partial(\Delta^N \times \mathbb{A}^1)$ is equidimensional, and so has dimension $\leq d + \dim(F)$. Hence W has codimension at least i in $X \times F$ (resp., is equidimensional). \square

In order to prove that $z_{equi}^i(X, *) \rightarrow z^i(X, *)$ is a quasi-isomorphism in theorem 18.3, we introduce the “topological” notion of weak homotopy.

DEFINITION 18.13. Two maps $f, g : K \rightarrow L$ of complexes of abelian groups are called **weakly homotopic** if for every finitely generated subcomplex C of K , the restrictions $f|_C$ and $g|_C$ are chain homotopic.

It is easy to check that weakly homotopic maps induce the same maps on homology. If K and L are bounded complexes of free abelian groups, this notion is

equivalent to the usual notion of chain homotopy between maps. To see that this notion is weaker than chain homotopy, consider a pure subgroup A of B which is not a summand, such as $\bigoplus_1^\infty \mathbb{Z} \subset \prod_1^\infty \mathbb{Z}$. Then the canonical map from $(A \rightarrow B)$ to $(A \rightarrow 0)$ is weakly homotopic to zero but not chain contractible.

LEMMA 18.14. (See [HighCh, 2.3 and 2.6]) *Let φ be an N -skeletal map over X . Then the inclusion map ι and the map φ^* are weakly homotopic on φz^i :*

$$\varphi z^i(X, *) \xrightarrow[\varphi^*]{\iota} z^i(X, *),$$

and also on φz_{equi}^i :

$$\varphi z_{equi}^i(X, *) \xrightarrow[\varphi^*]{\iota} z_{equi}^i(X, *).$$

PROOF. Consider a subcomplex $C_* \hookrightarrow \varphi z^i(X, *)$ generated by some closed irreducible subvarieties V_n^k so that $\partial_j(V_n^k)$ is a linear combination of generators. By 18.12, there is an N -skeletal homotopy Φ such that $C_* \subset \Phi z^i(X, *)$, and if C_* lies in $\varphi z_{equi}^i(X, *)$ then $C_* \subset \Phi z_{equi}^i(X, *)$. By 18.11, Φ induces a chain homotopy between φ^* and ι .

Note that the following diagram commutes:

$$\begin{array}{ccc} \varphi z_{equi}^i(X, -) & \hookrightarrow & \varphi z^i(X, -) \\ \downarrow \iota & & \downarrow \iota \\ \varphi^* & & \varphi^* \\ \downarrow & & \downarrow \\ z_{equi}^i(X, -) & \hookrightarrow & z^i(X, -). \end{array}$$

Moreover if $a \in \varphi z^i(X, n) \cap z_{equi}^i(X, n)$, and $\varphi^* a \in z_{equi}^i(X, n)$, then $a \in \varphi z_{equi}^i(X, n)$. \square

PROOF OF THEOREM 18.3. We have to prove that the induced map on homology classes is an isomorphism:

$$(18.14.1) \quad H_n(z_{equi}^i(X, *)) \rightarrow H_n(z^i(X, *)).$$

First we prove surjectivity. Let $a \in z^i(X, n)$ be such that $d(a) = 0$. Lemma 18.9 provides an integer N and an N -skeletal map $\{\varphi_n\}$ such that $a \in \varphi z^i(X, n)$ and $\varphi^*(a) \in z_{equi}^i(X, n)$. By 18.14, $a - \varphi^*a$ is a boundary in $z^i(X, n)$, i.e., a and $\varphi^*(a)$ represent the same class in homology. Hence the map 18.14.1 is surjective.

For injectivity we need to consider $a \in z_{equi}^i(X, n)$ so that $d(a) = 0$ and $b \in z^i(X, n+1)$ with $d(b) = a$. Apply lemma 18.9 to b and a . We find an $(n+1)$ -skeletal map φ such that $a, b \in \varphi z^i(X, *)$ and $\varphi^*a, \varphi^*b \in z_{equi}^i(X, *)$. But now we have:

$$\varphi^*a = \varphi^*(db) = d(\varphi^*b) = 0.$$

From lemma 18.14, a and $\varphi^*a = 0$ represent the same class in the homology of $z_{equi}^i(X, *)$. Therefore a is a boundary in $z_{equi}^i(X, *)$. Hence the map (18.14.1) is also injective. \square

Generic equidimensionality

This appendix is devoted to a proof of the following Generic Equidimensionality Theorem, due to Suslin. (See [HighCh] 1.1.)

THEOREM 18A.1. *Let S be an affine scheme of finite type over a field. Let V be a closed subscheme of $S \times \mathbb{A}^n$, Z an effective divisor of \mathbb{A}^n and $\varphi : S \times Z \rightarrow S \times \mathbb{A}^n$ any morphism over S . For every $t \geq 0$ so that $\dim V \leq n + t$, there exists a map $\Phi : S \times \mathbb{A}^n \rightarrow S \times \mathbb{A}^n$ over S so that:*

- (1) $\Phi|_{S \times Z} = \varphi$;
- (2) *the fibers of the projection $\Phi^{-1}(V) \rightarrow \mathbb{A}^n$ have dimension $\leq t$ over the points of $\mathbb{A}^n - Z$.*

The S -morphism $\varphi : S \times Z \rightarrow S \times \mathbb{A}^n$ is determined by its component $\varphi' : S \times Z \rightarrow \mathbb{A}^n$. If $S \subset \mathbb{A}^m$, we can extend φ' to a morphism $\psi' : \mathbb{A}^m \times Z \rightarrow \mathbb{A}^n$. If we knew the theorem for \mathbb{A}^m , there would exist an extension $\Psi' : \mathbb{A}^m \times \mathbb{A}^n \rightarrow \mathbb{A}^n$ of ψ' such that, setting $\Psi(X, Y) = (X, \Psi'(X, Y))$, the fibers of $\Psi^{-1}(V) \rightarrow \mathbb{A}^n$ over points of $\mathbb{A}^n - Z$ have dimension $\leq t$, and the restriction Φ of Ψ to $S \times \mathbb{A}^n$ would satisfy the conclusion of the theorem. Thus we may suppose that $S = \mathbb{A}^m$.

Write $\mathbb{A}^m = \text{Spec } k[x_1, \dots, x_m]$ and $\mathbb{A}^n = \text{Spec } k[y_1, \dots, y_n]$. If the divisor Z is defined by a polynomial $h \in k[Y]$ then the component $\varphi' : \mathbb{A}^m \times Z \rightarrow \mathbb{A}^n$ of φ extends to $f = (f_1, \dots, f_n) : \mathbb{A}^m \times \mathbb{A}^n \rightarrow \mathbb{A}^n$ for polynomials $f_i \in k[X, Y]$ defined up to a multiple of h . For each n -tuple $F = (F_1, \dots, F_n)$ of homogeneous forms in $k[X]$ of degree N , consider the maps

$$\Phi_F : \mathbb{A}^m \times \mathbb{A}^n \rightarrow \mathbb{A}^n$$

$$\Phi_F(X, Y) = (f_1(X, Y) + h(Y)F_1(X), \dots, f_n(X, Y) + h(Y)F_n(X)).$$

By construction, the restriction of Φ_F to $Z \times S$ is φ' , i.e., property (1) holds. It suffices to show that if $N \gg 0$ and the F_i are in general position then $\Phi(X, Y) = (X, \Phi_F(X, Y))$ has the desired property (2).

If $I = (g_1, \dots, g_s)$ is the ideal of $k[X, Y]$ defining V , then the ideal J of $k[X, Y]$ defining $\Phi^{-1}(V)$ is generated by the polynomials

$$g_j(X, \Phi_F) = g_j(x_1, \dots, x_m, \Phi_1, \Phi_2, \dots, \Phi_n), \quad \Phi_i = f_i(X, Y) + h(Y)F_i(X).$$

If b is a k -point of \mathbb{A}^n , the ideal J_b of $k[X]$ defining the fiber over b is generated by the $g_j(X, \Phi_F(X, b))$. We need to show that if $b \notin Z$, then J_b has height $\geq m - t$.

EXAMPLE 18A.2. Suppose that $m = 1$ and $t = 0$. We may assume that $\dim V = n$, and that V is defined by $g(x, Y) = 0$. Then $\Phi^{-1}(V)$ is defined by $g(x, \Phi_F)$, $F_i(x) = a_i x^N$, and the fiber over $b \in \mathbb{A}^n - Z$ is defined by

$$g(x, f_1(x, b) + h(b)a_1 x^N, \dots) = 0.$$

Since $b \notin Z$, $h(b) \neq 0$. Hence the left side of this equation is a nonzero polynomial in $k[x]$ for almost all choices of a_1, \dots, a_n when $N \gg 0$. Hence the fiber over b is finite.

The same argument works more generally when $t = m - 1$; we may assume that V is defined by $g = 0$, and the fiber over b is defined by $g(X, \Phi_F(X, b)) = 0$. In order to see that the left side is nonzero for almost all choices of F_1, \dots, F_n one just needs to analyze the leading form of $g(X, \Phi_F)$ with respect to X .

For any ring R we grade the polynomial ring $R[X] = R[x_1, \dots, x_m]$ with all x_i in degree 1. Any polynomial of degree d is the sum $f = F_d + \dots + F_0$ where F_i is a homogeneous form of degree i ; F_d is called the **leading form** of f with respect to X . If I is an ideal in $R[X]$ the leading forms of elements of I generate a homogeneous ideal I' of $R[X]$.

LEMMA 18A.3. *Let R be a catenary Noetherian ring, $I \subset R[X]$ an ideal, and I' the ideal of leading forms in I with respect to X . Then $\text{ht}(I') = \text{ht}(I)$.*

PROOF. Let $I_h \subset S = R[x_0, \dots, x_m]$ be the homogeneous ideal defining the closure \bar{V} of $V(I)$ in \mathbb{P}_R^m . Then $\text{ht}(I) = \text{ht}_S(I_h) = \text{ht}_S(I_h, x_0) - 1$. But $I' = (I_h, x_0)S/x_0S$, so $\text{ht}(I') = \text{ht}_S(I_h, x_0) - 1$. \square

Now the ring $k[X, Y]$ is bigraded, with each x_i of bidegree $(0, 1)$ and each y_i of bidegree $(1, 0)$. Thus each polynomial can be written as a sum $g = \sum G_{ij}$, where the G_{ij} have bidegree (i, j) . Ordering the bidegrees lexicographically allows us to talk about the bidegree of g , namely the largest (p, q) with $G_{pq} \neq 0$; this G_{pq} is the bi-homogeneous leading form of g .

Without loss of generality, we assume that the generators g_1, \dots, g_s of I have the following property: the bi-homogeneous leading forms $G_j(X, Y)$ of g_j generate the ideal of the leading forms of I .

LEMMA 18A.4. *If F_1, \dots, F_n are homogeneous forms in $k[X]$ of degree $N > \max\{\deg_X(f_i), \deg_X(g_j)\}$ then the ideal J' of leading forms in J with respect to X contains forms $h^r G_j(X, F_1, \dots, F_n)$, for $r \gg 0$.*

PROOF. (See [HighCh] 1.6.1.) Recall that J is generated by the $g_j(X, \Phi_F)$. For any choice of the N -forms F_i it is easy to see that $\deg_X g_j(X, \Phi_F) = \deg_X G_j(X, \Phi_F) = N \deg_Y G_j + \deg_X G_j$, and that the leading form in $g_j(X, \Phi_F)$ with respect to X is $h^{\deg_Y G_j} G_j(X, F_1, \dots, F_n)$. \square

PROPOSITION 18A.5. *Let $T \subset \mathbb{A}^m \times \mathbb{A}^n$ be a closed subscheme of dimension $\leq n + t$, $t \geq 0$. If k is infinite, then for any $N \geq 0$ we can find forms F_1, \dots, F_n in $k[X]$ of degree N so that $W = \{w \in \mathbb{A}^m : (w, F_1(w), \dots, F_n(w)) \in T\}$ has dimension at most t .*

PROOF. The vector space of n -tuples $F = (F_1, \dots, F_n)$ of homogeneous forms of degree N in $k[X]$ is finite-dimensional, say of dimension D . We identify it with the set of k -rational points of the affine space \mathbb{A}^D . Consider the evaluation map

$$\eta : \mathbb{A}^m \times \mathbb{A}^D \rightarrow \mathbb{A}^{m+n}, \quad \eta(w, F) = (w, F(w)).$$

If $w \neq 0$, the fibers of $\eta : w \times \mathbb{A}^D \rightarrow w \times \mathbb{A}^n$ are isomorphic to \mathbb{A}^{D-n} , because the linear homomorphism $\eta(w, -) : \mathbb{A}^D \rightarrow \mathbb{A}^n$ is surjective. By inspection, $\eta^{-1}(0 \times \mathbb{A}^n) = 0 \times \mathbb{A}^D$. It follows that $\eta^{-1}(T)$ has dimension at most $D+t$.

Now consider the projection $\pi : \eta^{-1}(T) \rightarrow \mathbb{A}^D$. The theorem on dimension of the fibers [Har77] III.9.6 implies that there is a nonempty $U \subset \mathbb{A}^D$ whose fibers have dimension $\leq t$. Choosing a rational point in U , the corresponding homogeneous forms (F_1, \dots, F_n) satisfy $\dim\{w \in \mathbb{A}^m : (w, F(w)) \in T\} \leq t$. \square

REMARK 18A.6. The case $N = 0$ is easy to visualize, since $D = n$. There is an open subset U of \mathbb{A}^n so that for each $b \in U$ the fiber $T \cap (\mathbb{A}^m \times b)$ of the projection $T \rightarrow \mathbb{A}^n$ over b has dimension at most t .

If T is defined by bi-homogeneous polynomials, then W is defined by homogeneous polynomials. Suslin states 18A.5 for the corresponding projective varieties in [HighCh, 1.7].

We are now ready to complete the proof of theorem 18A.1. By 18A.3, $J_b \subset k[X]$ has the same height as the ideal J'_b of its leading forms. Suppose that $N > \max\{\deg_X(f_i), \deg_X(g_j)\}$. Since $h(b) \neq 0$, J'_b contains all the $G_j(X, F)$ by 18A.4. Let $T \subset \mathbb{A}^{m+n}$ be the variety defined by the ideal of bi-homogeneous forms of I , i.e., the $G_j(X, Y)$. Hence the variety $W = \{w \in \mathbb{A}^m : (w, F(w)) \in T\}$ is defined by the $G_j(X, F)$. By two applications of 18A.3, $\dim T = \dim V \leq n+t$. Thus $\dim W \leq t$ by 18A.5. But the height of J'_b is at least the height of the ideal generated by the $G_j(X, F)$, i.e., the codimension of W , which is at least $m-t$.

Motivic cohomology and higher Chow groups

With the preparation of the last three lectures, we are ready to prove the fundamental comparison theorem:

THEOREM 19.1. *Let X be a smooth separated scheme over a perfect field k , then for all n and $i \geq 0$ there is a natural isomorphism:*

$$H^{n,i}(X, \mathbb{Z}) \xrightarrow{\cong} CH^i(X, 2i - n).$$

At the end of this lecture, we will generalize this to all schemes of finite type, replacing motivic cohomology by Borel-Moore motivic homology. Assuming resolution of singularities we will prove in 19.18 that $CH^{d-i}(X, n) \cong H_{2i+n,i}^{BM}(X, \mathbb{Z})$.

Because $CH^i(X, 0)$ is the classical Chow group $CH^i(X)$ we obtain:

COROLLARY 19.2. $H^{2i,i}(X, \mathbb{Z}) \cong CH^i(X)$.

It is clear from definition 17.1 that $CH^i(X, m) = 0$ for $m < 0$. We immediately deduce the:

VANISHING THEOREM 19.3. *For every smooth variety X and any abelian group A , we have $H^{n,i}(X, A) = 0$ for $n > 2i$.*

The proof of 19.1 will proceed in two stages. First we will show (in theorem 19.8) that $\mathbb{Z}(i)[2i]$ is quasi-isomorphic to $U \mapsto z^i(U \times \mathbb{A}^i, *)$ as a complex of Zariski sheaves. Then we will show (in 19.12) that the hypercohomology of $z^i(- \times \mathbb{A}^i, *)$ is $CH^i(-, *)$.

We saw in 16.7 that $\mathbb{Z}(i)$ is quasi-isomorphic to the Suslin-Friedlander motivic complex $\mathbb{Z}^{SF}(i)$. Recall from page 126 that the shift $\mathbb{Z}^{SF}(i)[2i]$ is the chain complex $C_{*z_{equi}}(\mathbb{A}^i, 0)$ associated to the simplicial abelian presheaf with transfers $C_{\bullet z_{equi}}(\mathbb{A}^i, 0)$, which sends X to $m \mapsto z_{equi}(\mathbb{A}^i, 0)(X \times \Delta^m)$. The following result generalizes example 17.2.

LEMMA 19.4. *Let T be smooth of dimension d . If $0 \leq i \leq d$ then for all X there is an embedding of simplicial abelian groups:*

$$C_{\bullet z_{equi}}(T, d - i)(X) \hookrightarrow z^i(X \times T, \bullet).$$

*In particular (for $T = \mathbb{A}^i$), $\mathbb{Z}^{SF}(i)[2i](X)$ is a subcomplex of $z^i(X \times \mathbb{A}^i, *)$.*

PROOF. The cycles in $C_{mz_{equi}}(T, d - i)(X)$ are equidimensional over $X \times \Delta^m$ at all points, while the ones in $z^i(X \times T, m)$ need only be equidimensional at the generic points of the faces of $X \times T \times \Delta^m$. Hence the first group is contained in the

second group of cycles. Moreover, the face maps of the two simplicial groups are compatible by 1A.14. \square

EXAMPLE 19.5. The complex $\mathbb{Z}^{SF}(i)[2i](Y)$ is a subcomplex of $z^i(Y \times \mathbb{A}^i, *)_{\mathcal{W}}$ (see 17.6) for every finite correspondence \mathcal{W} from X to Y . Indeed, $z_{equi}(\mathbb{A}^i, 0)(Y \times \Delta^m)$ lies in $z^i(Y \times \mathbb{A}^i, m)_{\mathcal{W} \times \mathbb{A}^i}$ because every generating cycle is quasi-finite over $Y \times \Delta^m$.

In contrast, it is easy to see that $z_{equi}(Y \times \mathbb{A}^i, \dim Y)(\Delta^m)$ need not lie in $z^i(Y \times \mathbb{A}^i, m)_{\mathcal{W} \times \mathbb{A}^i}$, by letting X be a point of Y .

For any schemes X and T , consider the simplicial presheaf on X :

$$U \mapsto z^i(U \times T, \bullet).$$

This can be regarded as a simplicial sheaf on the flat site over X and hence on both the (small) étale site and the Zariski site of X as well. We will write $z^i(- \times T, *)$ for the associated complex of sheaves. The homology of $z^i(- \times T, *)$ has the more general structure of a presheaf with transfers by 17.21.

PROPOSITION 19.6. *The homology of the embedding in 19.4 is a morphism of presheaves with transfers:*

$$(19.6.1) \quad H_m C_* z_{equi}(\mathbb{A}^i, 0)(-) \rightarrow H_m z^i(- \times \mathbb{A}^i, *) = CH^i(- \times \mathbb{A}^i, m).$$

PROOF. The source and target are presheaves with transfers by 16.3 and 17.21, respectively. It suffices to show that their transfer maps are compatible.

Let W be an elementary correspondence from X to Y . We need to verify that ϕ_W and W^* are compatible with the map (19.6.1). If W is the graph of a flat map from X to Y , then ϕ_W and W^* are compatible because both are just the flat pullback of cycles. Since W^* is defined in 17.17 by passing to an affine vector bundle torsor $Y' \rightarrow Y$, a simple diagram chase (which we leave to the reader) shows that it suffices to prove the statement when Y is affine.

Let Y be affine. Since $H_n z^i(Y \times \mathbb{A}^i, m)_W = H_n z^i(Y \times \mathbb{A}^i, m)$ by 17.6, the result will follow once we show that the following diagram commutes.

$$\begin{array}{ccc} z_{equi}(\mathbb{A}^i, 0)(Y \times \Delta^m) & \xrightarrow{\phi_W} & z_{equi}(\mathbb{A}^i, 0)(X \times \Delta^m) \\ \text{19.5} \downarrow & & \downarrow \text{19.4} \\ z^i(Y \times \mathbb{A}^i, m)_W & \xrightarrow{W^*} & z^i(X \times \mathbb{A}^i, m) \end{array}$$

Let i, f and π , respectively, denote the products with $\mathbb{A}^i \times \Delta^m$ of the inclusion $W \hookrightarrow X \times Y$, and the canonical projections $X \times Y \rightarrow Y$ and $X \times Y \rightarrow X$. The transfer map W^* was defined as $W^*(\mathcal{Z}) = \pi_*((W \times \mathbb{A}^i \times \Delta^m) \cdot f^* \mathcal{Z})$ in 17.7. According to 16.3, the transfer map on $z_{equi}(\mathbb{A}^i, 0)(Y \times \Delta^m)$ is $\phi_W(\mathcal{Z}) = (i\pi)_*(\mathcal{Z}_{W \times \Delta^m})$, where the pullback $\mathcal{Z}_{W \times \Delta^m}$ was defined on page 10. By 17A.13, $\mathcal{Z}_{W \times \Delta^m} = (fi)^*(\mathcal{Z})$, so we have:

$$\phi_W(\mathcal{Z}) = (i\pi)_*(fi)^*(\mathcal{Z}) = \pi_* i_*(fi)^*(\mathcal{Z}).$$

By 17A.12, $i_*(fi)^*(\mathcal{L}) = (W \times \mathbb{A}^i \times \Delta^m) \cdot f^* \mathcal{L}$ and therefore for every \mathcal{L} in $z_{\text{equi}}(\mathbb{A}^i, 0)(Y \times \Delta^m)$ we have:

$$\phi_W(\mathcal{L}) = \pi_*((W \times \mathbb{A}^i \times \Delta^m) \cdot f^* \mathcal{L}) = W^*(\mathcal{L}). \quad \square$$

EXAMPLE 19.7. If E is a field over k , then the map of 19.6 evaluated at $\text{Spec } E$ is an isomorphism:

$$H_m C_* z_{\text{equi}}(\mathbb{A}^i, 0)(\text{Spec } E) \xrightarrow{\cong} H_m z^i(\text{Spec } E \times \mathbb{A}^i, *).$$

This follows from Suslin's theorem 18.3 with $X = \mathbb{A}_E^i$, since we may identify $z_{\text{equi}}(\mathbb{A}_k^i, 0)(\Delta_E^m)$ and $z_{\text{equi}}(\mathbb{A}_E^i, 0)(\Delta^m)$ by 16.6.

This implies that theorem 19.1 is true when evaluated on fields. To see this, set $S = \text{Spec } E$ and recall that $\mathbb{H}^m(S, C^*) = H^m C^*(S)$ for any complex of sheaves C^* . By 16.7, the above map fits into the sequence of isomorphisms:

$$\begin{aligned} H^{n,i}(S, \mathbb{Z}) &\cong H^n \mathbb{Z}(i)(S) \cong H^n \mathbb{Z}^{SF}(i)(S) \\ &= H_{2i-n} C_* z_{\text{equi}}(\mathbb{A}^i, 0)(S) \xrightarrow{\cong} H_{2i-n} z^i(\mathbb{A}_E^i, *) \\ &= CH^i(\mathbb{A}_E^i, 2i-n) \cong CH^i(S, 2i-n). \end{aligned}$$

THEOREM 19.8. *The map $\mathbb{Z}^{SF}(i)[2i] = C_* z_{\text{equi}}(\mathbb{A}^i, 0) \rightarrow z^i(- \times \mathbb{A}^i, *)$ is a quasi-isomorphism of complexes of Zariski sheaves.*

PROOF. The induced homomorphisms on homology presheaves,

$$(19.8.1) \quad H_m C_* z_{\text{equi}}(\mathbb{A}^i, 0) \rightarrow H_m z^i(- \times \mathbb{A}^i, *),$$

are morphisms of presheaves with transfers by 19.6. The left side is homotopy invariant by 2.19 and the right side is homotopy invariant because the higher Chow groups are homotopy invariant (see 17.4). By 19.7, this is an isomorphism for all fields. By 11.2, the sheafification of the map (19.8.1) is an isomorphism. Hence $C_* z_{\text{equi}}(\mathbb{A}^i, 0) \rightarrow z^i(- \times \mathbb{A}^i, *)$ is a quasi-isomorphism for the Zariski topology. \square

COROLLARY 19.9. *For any smooth scheme X , the inclusion of 19.4 induces an isomorphism:*

$$H^{n,i}(X, \mathbb{Z}) \xrightarrow{\cong} \mathbb{H}^{n-2i}(X, z^i(- \times \mathbb{A}^i, *)).$$

PROOF. By 16.7 and 19.8, we have the sequence of isomorphisms:

$$\begin{aligned} H^{n,i}(X, \mathbb{Z}) &= \mathbb{H}^n(X, \mathbb{Z}(i)) \cong \mathbb{H}^n(X, \mathbb{Z}^{SF}(i)) \\ &= \mathbb{H}^{n-2i}(X, \mathbb{Z}^{SF}(i)[2i]) \xrightarrow{\cong} \mathbb{H}^{n-2i}(X, z^i(- \times \mathbb{A}^i, *)). \quad \square \end{aligned}$$

Corollary 19.9 is the first half of the proof of 19.1. The rest of this lecture is dedicated to proving the second half, that $\mathbb{H}^{-m}(X, z^i(- \times \mathbb{A}^i, *)) \cong CH^i(X, m)$. To do this, we shall use Bloch's Localization Theorem (see 17.4) to reinterpret the higher Chow groups as the hypercohomology groups of a complex of sheaves.

A chain complex of presheaves C is said to satisfy **Zariski descent** on X if $H^*(C(U)) \rightarrow \mathbb{H}^*(U, C_{\text{Zar}})$ is an isomorphism for every open U in X .

DEFINITION 19.10. Let C be a complex of presheaves on X_{Zar} (the small Zariski site of X). We say that C has the (Zariski) **Mayer-Vietoris property** if for every $U \subset X$, and any open covering $U = V_1 \cup V_2$, the diagram

$$\begin{array}{ccc} C(U) & \longrightarrow & C(V_1) \\ \downarrow & & \downarrow \\ C(V_2) & \longrightarrow & C(V_1 \cap V_2) \end{array}$$

is homotopy cartesian (i.e., the total complex is an acyclic presheaf). This implies that there is a long exact sequence

$$\cdots \rightarrow H^i(C(U)) \rightarrow H^i(C(V_1)) \oplus H^i(C(V_2)) \rightarrow H^i(C(V_1 \cap V_2)) \rightarrow \cdots .$$

For example, any chain complex of flasque sheaves has the Mayer-Vietoris property. This is an easy consequence of the fact that $C(U) \rightarrow C(V)$ is onto for each $V \subset U$.

The following result is proven in [BG73].

THEOREM 19.11 (Brown-Gersten). *Let C be a complex of presheaves on X with the Mayer-Vietoris property. Then C satisfies Zariski descent. That is, the maps $H^*(C(U)) \rightarrow \mathbb{H}^*(U, C_{Zar})$ are all isomorphisms.*

Our main application of the Brown-Gersten theorem is to prove that Bloch's complexes satisfy Zariski descent.

PROPOSITION 19.12. *Let X be any scheme of finite type over a field. For any scheme T , each $z^i(- \times T)$ satisfies Zariski descent on X . That is, for all m and i , we have:*

$$CH^i(X \times T, m) \cong \mathbb{H}^{-m}(X, z^i(- \times T)).$$

In particular (for $T = \mathbb{A}^i$),

$$CH^i(X, m) \xrightarrow{\cong} CH^i(X \times \mathbb{A}^i, m) \xrightarrow{\cong} \mathbb{H}^{-m}(X, z^i(- \times \mathbb{A}^i)).$$

PROOF. (Bloch [Blo86, 3.4]) By 19.11, we have to show that $C(U) = z^i(U \times T)$ has the Mayer-Vietoris property. For each cover $\{V_1, V_2\}$ of each U we set $V_{12} = V_1 \cap V_2$ and consider the diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C(U - V_1) & \longrightarrow & C(U) & \longrightarrow & C(V_1) & \longrightarrow & \text{coker}_1 & \longrightarrow & 0 \\ & & \downarrow = & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C(V_2 - V_{12}) & \longrightarrow & C(V_2) & \longrightarrow & C(V_{12}) & \longrightarrow & \text{coker}_2 & \longrightarrow & 0. \end{array}$$

By Bloch's Localization Theorem, the cokernels are both acyclic. A diagram chase shows that the middle square is homotopy cartesian, i.e., the Mayer-Vietoris condition is satisfied. \square

We are now ready to prove the main result of this section, theorem 19.1.

PROOF OF 19.1. Using 19.9 and 19.12, we define the map to be the compositions of isomorphisms:

$$H^{n,i}(X, \mathbb{Z}) \cong \mathbb{H}^n(X, \mathbb{Z}(i)) \xrightarrow{\cong} \mathbb{H}^{n-2i}(X, z^i(- \times \mathbb{A}^i)) \cong CH^i(X, 2i - n). \quad \square$$

Zariski descent has also been used by Bloch and Levine to show that the higher Chow groups are functorial for morphisms between smooth schemes. We first recall their definition and then show in proposition 19.16 below that it agrees with ours.

DEFINITION 19.13. (Bloch-Levine) Let f be a morphism from X to Y . Natural maps $f^* : CH^i(Y, m) \rightarrow CH^i(X, m)$ for all m and i are defined as follows. As in the proof of 17.6, write $z^i(Y, *)_f$ for $z^i(Y, *)_{\Gamma_f}$.

If $U \subset Y$ is open, $z^i(Y, *)_f$ restricts to $z^i(U, *)_f$, and z^i_f is a complex of sheaves. Since Y is locally affine, $z^i_f \simeq z^i$ by 17.6 and there is a map $z^i_f \rightarrow f_* z^i$ of complexes of sheaves on Y . The map is now defined using Zariski descent 19.12 as the composite:

$$CH^i(Y, m) \cong \mathbb{H}^{-m}(Y, z^i) \cong \mathbb{H}^{-m}(Y, z^i_f) \xrightarrow{f^*} \mathbb{H}^{-m}(X, z^i) \cong CH^i(X, m).$$

EXAMPLE 19.14. If $q : Y' \rightarrow Y$ is flat, then $z^i_q = z^i$, and the map q^* defined in 19.13 is just the flat pullback of cycles map q^* , described in 17.12.

LEMMA 19.15. If $X \xrightarrow{g} Y \xrightarrow{f} Z$ are morphisms of smooth schemes, then the maps defined in 19.13 satisfy $(fg)^* = g^* f^*$.

PROOF. If $fg \amalg f : X \amalg Y \rightarrow Z$, we can restrict $(fg)^*$ and f^* to the subgroup $z^i(Z, m)_{fg \amalg f}$. Since $(fg)^* = g^* f^*$ on cycles (see [Ser65, V-30]), f^* maps this subgroup into $z^i(Y, m)_g$. By construction, the diagram of groups

$$\begin{array}{ccc} z^i(Z, m)_{fg \amalg f} & \hookrightarrow & z^i(Z, m)_{fg} \\ \downarrow f^* & & \downarrow (fg)^* \\ z^i(Y, m)_g & \xrightarrow{g^*} & z^i(X, m) \end{array}$$

commutes. Sheafifying and applying hypercohomology, 17.6 and Zariski descent 19.12 show that the composite

$$CH^i(Z, m) \cong \mathbb{H}^{-m}(Z, z^i_{fg \amalg f}) \xrightarrow{f^*} \mathbb{H}^{-m}(Y, z^i_g) \xrightarrow{g^*} \mathbb{H}^{-m}(X, z^i) \cong CH^i(X, m)$$

is just $(fg)^*$, as required. □

PROPOSITION 19.16. The map $f^* : CH^i(Y, m) \rightarrow CH^i(X, m)$ defined in 19.13 agrees with the map $f^* = \Gamma_f^*$ defined in 17.17.

PROOF. Suppose first that X and Y are affine, and consider the commutative diagram

$$\begin{array}{ccccc}
 CH^i(Y, m) = H^{-m}z^i(Y, *) & \xleftarrow{\cong} & H^{-m}z^i(Y, *)_f & \longrightarrow & H^{-m}z^i(X, *) = CH^i(X, m) \\
 \downarrow \cong & & \downarrow & & \downarrow \cong \\
 \mathbb{H}^{-m}(Y, z^i) & \xleftarrow{\cong} & \mathbb{H}^{-m}(Y, z^i_f) & \longrightarrow & \mathbb{H}^{-m}(X, z^i).
 \end{array}$$

The arrows marked ‘ \cong ’ are isomorphisms by 17.6 and 19.12. The top composite is the map of 17.12, which by 17.18 is the map Γ_f^* of 17.17. The bottom composite is the map f^* of 19.13, proving that $f^* = \Gamma_f^*$ in this case.

In the general case, 17.15 gives a diagram

$$\begin{array}{ccc}
 X' & \xrightarrow{g} & Y' \\
 p \downarrow & & \downarrow q \\
 X & \xrightarrow{f} & Y,
 \end{array}$$

where $X' \rightarrow X$ and $Y' \rightarrow Y$ are affine vector bundle torsors. By definition 17.17, Γ_f^* is $(p^*)^{-1}\Gamma_g^*q^*$, where p^* and q^* are flat pullbacks of cycles. By 19.14, these are the same as the maps p^* and q^* defined in 19.13. Since $\Gamma_g^* = g^*$ by the first part of the proof and $g^*q^* = (qg)^* = (pf)^* = p^*f^*$ by 19.15, we have:

$$\Gamma_f^* \stackrel{17.17}{=} (p^*)^{-1}\Gamma_g^*q^* = (p^*)^{-1}g^*q^* \stackrel{19.15}{=} (p^*)^{-1}p^*f^* = f^*. \quad \square$$

We conclude this lecture by reinterpreting theorem 19.1 in terms of the Borel-Moore motivic homology groups $H_{n,i}^{BM}(X, \mathbb{Z}) = \text{Hom}_{\mathbf{DM}_{Nis}^{\text{eff.}}}(\mathbb{Z}(i)[n], M^c(X))$, assuming resolution of singularities. We begin with the smooth case.

EXAMPLE 19.17. When X is smooth of dimension d , the identification follows from the isomorphism $CH^i(X, n) \cong H^{2i-n,i}(X, \mathbb{Z})$ of 19.1. To see this we set $j = d - i$ and compute:

$$\begin{aligned}
 CH^j(X, n) &\cong H^{2j-n,j}(X, \mathbb{Z}) && \text{by 19.1,} \\
 &= \text{Hom}(M(X), \mathbb{Z}(j)[2j-n]) && \text{by 14.16} \\
 &= \text{Hom}(\mathbb{Z}(d)[2d], M^c(X)(j)[2j-n]) && \text{by 16.24.} \\
 &= \text{Hom}(\mathbb{Z}(i)[2i+n], M^c(X)) && \text{by 16.25.} \\
 &= H_{2i+n,i}^{BM}(X, \mathbb{Z}) && \text{by definition 16.20.}
 \end{aligned}$$

We now establish this isomorphism when X is not smooth, using 16.22.

PROPOSITION 19.18. *Assume that k admits resolution of singularities. Let X be a quasi-projective equidimensional scheme over k of dimension d . Then for every positive $i \leq d$ and n there is a canonical isomorphism:*

$$CH^{d-i}(X, n) \cong H_{2i+n,i}^{BM}(X, \mathbb{Z}) = \text{Hom}(\mathbb{Z}(i)[2i+n], M^c(X)).$$

PROOF. By 16.22, the right-hand side is isomorphic to $\text{Hom}(\mathbb{Z}[n], C_*z_{\text{equi}}(X, i))$. But by 13.5 this is isomorphic to $H_n C_*z_{\text{equi}}(X, i)(\text{Spec } k)$, i.e., the n -th homology of the complex of abelian groups $z_{\text{equi}}^{d-i}(X, *)$. We conclude using corollary 18.5. \square

COROLLARY 19.19. *If $i \geq 0$ there are canonical isomorphisms:*

$$CH^{d+i}(X, n) \cong \text{Hom}(\mathbb{Z}, M^c(X)(i)[2i - n]).$$

PROOF. By homotopy invariance (see 17.4), $CH^{d+i}(X, n) = CH^{d+i}(X \times \mathbb{A}^i, n)$. By 19.18, this is $\text{Hom}(\mathbb{Z}[n], M^c(X \times \mathbb{A}^i))$, and $M^c(X \times \mathbb{A}^i) \cong M^c(X)(i)[2i]$ by 16.16. \square

LECTURE 20

Geometric motives

In lectures 14 and 16 we introduced the category $\mathbf{DM}_{gm}^{\text{eff}}$ of effective geometric motives, and the category \mathbf{DM}_{gm} of all geometric motives. In this lecture we complete our investigation of the properties of these categories.

We begin by embedding Grothendieck's classic category *Chow* of Chow motives into \mathbf{DM}_{gm} . We then construct the dual of any object in \mathbf{DM}_{gm} , based on the *RHom* of 14.12. This allows us to construct internal Hom objects $\underline{\text{Hom}}(X, Y)$. We will conclude this lecture by proving that the tensor triangulated subcategory \mathbf{DM}_{gm} of \mathbf{DM}^- is rigid.

Recall that Grothendieck's category of effective Chow motives Chow^{eff} is the idempotent completion of the category whose objects are smooth projective varieties over k , and whose morphisms are given by: $\text{Hom}_{\text{Chow}}(Y, X) = \text{CH}^{\dim X}(X \times Y)$. There is a canonical decomposition $\mathbb{P}^1 = (\text{Spec } k) \oplus \mathbb{L}$, where \mathbb{L} is the Lefschetz motive. The category *Chow* of Chow motives is obtained by inverting \mathbb{L} and Chow^{eff} is a full subcategory of *Chow*.

In this lecture k will always be a perfect field which admits resolution of singularities and the coefficients will be taken over \mathbb{Z} .

PROPOSITION 20.1. *Assume that k is a perfect field which admits resolution of singularities. Then Grothendieck's category of effective Chow motives embeds contravariantly into $\mathbf{DM}_{gm}^{\text{eff}}(k, \mathbb{Z})$, and hence into $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}(k, \mathbb{Z})$, in the sense that if X and Y are two smooth projective schemes, then*

$$\text{Hom}_{\text{Chow}}(Y, X) \cong \text{Hom}(M(X), M(Y)).$$

PROOF. We set $d = \dim X$ and compute in $\mathbf{DM}_{\text{Nis}}^{\text{eff}, -}$:

$$\begin{aligned} \text{CH}^d(X \times Y) &= H^{2d, d}(X \times Y, \mathbb{Z}) && \text{by 19.2,} \\ &= \text{Hom}(M(X \times Y), \mathbb{Z}(d)[2d]) && \text{by 14.16,} \\ &= \text{Hom}(M(X)(d)[2d], M(Y)(d)[2d]) && \text{by 16.24 and } Y \text{ proper} \\ &= \text{Hom}(M(X), M(Y)) && \text{by 16.25. } \square \end{aligned}$$

REMARK 20.2. The Lefschetz motive \mathbb{L} is mapped to $\mathbb{Z}(1)[2]$ by 13.17. So from 16.25 and 20.1 we have the following diagram of fully faithful tensor functors:

$$\begin{array}{ccccc} (\mathit{Chow}^{\text{eff}})^{\text{op}} & \longrightarrow & \mathbf{DM}_{gm}^{\text{eff}} & \longrightarrow & \mathbf{DM}_{Nis}^{\text{eff},-} \\ \downarrow & & \downarrow & & \downarrow \\ \mathit{Chow}^{\text{op}} & \longrightarrow & \mathbf{DM}_{gm} & \longrightarrow & \mathbf{DM}_{Nis}^- \end{array}$$

The category \mathbf{DM}_{gm} also has dual objects. We can construct the dual of any object in \mathbf{DM}_{gm} , based on the \underline{RHom} of 14.12. Recall that if B is in $\mathbf{DM}_{gm}^{\text{eff}}$ and A, C are in $\mathbf{DM}_{Nis}^{\text{eff},-}$ then in $\mathbf{DM}_{Nis}^{\text{eff},-}$ we have:

$$\text{Hom}(A \otimes B, C) \cong \text{Hom}(A, \underline{RHom}(B, C)).$$

By construction, the functor $\underline{RHom}(B, C)$ is triangulated in both variables.

PROPOSITION 20.3. *If X is smooth of dimension d , the diagonal $X \rightarrow X \times X$ induces isomorphisms for $r \geq 0$:*

$$\Delta_r : M^c(X)(r)[-2d] \cong \underline{RHom}(M(X), \mathbb{Z}(d))(r) \cong \underline{RHom}(M(X), \mathbb{Z}(d+r)).$$

PROOF. If $A = M(U)[n]$ for a smooth scheme U , we have:

$$\begin{aligned} \text{Hom}(A, M^c(X)(r)[-2d]) &= \text{Hom}(A(d)[2d], M^c(X)(d+r)) && \text{by 16.25} \\ &\cong \text{Hom}(A \otimes M(X), \mathbb{Z}(d+r)) && \text{by 16.24} \\ &\cong \text{Hom}(A, \underline{RHom}(M(X), \mathbb{Z}(d+r))) && \text{by 14.12.} \end{aligned}$$

When $A = M^c(X)[-2d]$, the graph of the identity on X (the diagonal) is the correspondence inducing the identity on $M(X)$ and on $M^c(X)$, so it induces natural maps Δ_r from $M^c(X)(r)[-2d]$ to $\underline{RHom}(M(X), \mathbb{Z}(d+r))$.

The subcategory of objects A for which $\text{Hom}(A, \Delta_r)$ is an isomorphism is triangulated, and contains the $M(U)[n]$, so it is all of $\mathbf{DM}_{Nis}^{\text{eff},-}$. The Yoneda lemma implies that each Δ_r is a natural isomorphism. \square

COROLLARY 20.4. *If X is a scheme in Sch/k , then $\underline{RHom}(M(X), \mathbb{Z}(i))$ is in $\mathbf{DM}_{gm}^{\text{eff}}$ for all $i \geq \dim(X)$.*

PROOF. It suffices to recall from 16.17 that each $M^c(X)$ is in $\mathbf{DM}_{gm}^{\text{eff}}$. \square

EXERCISE 20.5. Show that $\underline{RHom}(M(X), L) \cong \underline{RHom}(M(X)(1), L(1))$ for every smooth X and every L in $\mathbf{DM}_{Nis}^{\text{eff},-}$, by mimicking the proof of 20.3.

DEFINITION 20.6. If X is in Sm/k and $d = \dim X$, we define the dual to be:

$$M(X)^* = \underline{RHom}(M(X), \mathbb{Z}(d))(-d).$$

By 20.3, $M(X)^*$ is the same as $\underline{RHom}(M(X), \mathbb{Z}(i))(-i)$ for all $i \geq d$.

If M is any object of \mathbf{DM}_{gm} , some twist $M(r)$ is effective. We define the dual M^* to be $\underline{RHom}(M(r), \mathbb{Z}(i))(r-i)$ for large i . Note that M^* is independent of i and r by 20.5 and 20.3, and belongs to \mathbf{DM}_{gm} by 20.4. This independence implies that for every r there is a canonical isomorphism $M(r)^* \cong M^*(-r)$.

LEMMA 20.7. *The assignment $M \mapsto M^*$ is a contravariant triangulated functor from \mathbf{DM}_{gm} to itself.*

PROOF. By construction, each contravariant functor $\underline{RHom}(-, \mathbb{Z}(i))$ and each covariant functor $M \mapsto M(-i)$ is triangulated. Given any diagram in $\mathbf{DM}_{gm}^{\text{eff}}$, there is an i such that the dual coincides with the triangulated functor $\underline{RHom}(-, \mathbb{Z}(i))(-i)$ on the diagram. \square

The following proposition justifies the terminology “dual”. For simplicity, we write $\text{Hom}_{gm}(A, B)$ for $\text{Hom}_{\mathbf{DM}_{gm}}(A, B)$.

PROPOSITION 20.8. *The dual M^* of an object M in \mathbf{DM}_{gm} represents the functor $A \mapsto \text{Hom}_{gm}(A \otimes M, \mathbb{Z})$, in the sense that there is a natural isomorphism:*

$$\text{Hom}_{gm}(A, M^*) \cong \text{Hom}_{gm}(A \otimes M, \mathbb{Z}).$$

PROOF. Since $M(r)^*(r) \cong M^*$, $\text{Hom}_{gm}(A(-r), M(r)^*) \cong \text{Hom}_{gm}(A, M^*)$. Hence we may assume that M is effective. But then $\text{Hom}_{gm}(A, M^*) \cong \text{Hom}(A(i), \underline{RHom}(M, \mathbb{Z}(i)))$ for large i . By adjunction, this is $\text{Hom}(A(i) \otimes M, \mathbb{Z}(i))$. By 16.25, it is $\text{Hom}_{gm}(A \otimes M, \mathbb{Z})$. \square

COROLLARY 20.9. *There is a natural morphism $\varepsilon_M : M^* \otimes M \rightarrow \mathbb{Z}$ for every M in \mathbf{DM}_{gm} , adjoint to the identity of M^* .*

REMARK 20.10. The dual M^* is not the same as $\underline{RHom}(M, \mathbb{Z})$ in general. For the Lefschetz motive $\mathbb{L} = M(\mathbb{P}^1)/M(\text{Spec } k)$, for example, $\mathbb{L}^* \cong \underline{RHom}(\mathbb{L}, \mathbb{Z}(1))(-1) \cong \mathbb{Z}(-1)[-2]$, while exercise 14.13 implies that $\underline{RHom}(\mathbb{L}, \mathbb{Z}) = 0$.

EXAMPLE 20.11. If X is smooth of dimension d , the dual $M(X)^*$ is just an untwisting of $M^c(X)$. To see this, we combine 20.3 with definition 20.6:

$$M^c(X) \cong \underline{RHom}(M(X), \mathbb{Z}(d))[2d] \cong M(X)^*(d)[2d].$$

In particular, if X is projective then $M(X) \cong M(X)^*(d)[2d]$.

PROPOSITION 20.12. *There is a natural isomorphism $\iota_M : M \xrightarrow{\cong} M^{**}$ for M in \mathbf{DM}_{gm} .*

PROOF. The identity of M^* gives a natural map $\iota_M : M \rightarrow M^{**}$ via 20.8, adjoint to the map ε_M of 20.9:

$$\text{Hom}_{gm}(M, M^{**}) \cong \text{Hom}_{gm}(M \otimes M^*, \mathbb{Z}).$$

To prove that ι_M is an isomorphism for all M , it suffices to prove it when $M = M(X)$, where X is a smooth projective scheme of dimension d . Since $M(X)^*(d)$ is effective, we see by 20.6 and 20.3 that for all $i \geq d$:

$$\begin{aligned} (M(X)^*)^* &= \underline{RHom}(M(X)^*(d), \mathbb{Z}(i))(d-i) \\ &\cong \underline{RHom}(M(X)[-2d], \mathbb{Z}(i))(d-i) \cong M(X). \end{aligned}$$

A careful comparison of $t_{M(X)}$ with this isomorphism shows that they are inverse to each other. \square

PROPOSITION 20.13. *There is a natural isomorphism $M^* \otimes N^* \xrightarrow{\cong} (M \otimes N)^*$ for every M and N in \mathbf{DM}_{gm} .*

PROOF. There is a natural map $M^* \otimes N^* \rightarrow (M \otimes N)^*$ arising from $\varepsilon_M \otimes \varepsilon_N$ via the isomorphism of 20.8:

$$\begin{aligned} \mathrm{Hom}_{gm}(M^* \otimes N^*, (M \otimes N)^*) &\cong \mathrm{Hom}_{gm}(M^* \otimes N^* \otimes (M \otimes N), \mathbb{Z}) \\ &\cong \mathrm{Hom}_{gm}((M^* \otimes M) \otimes (N^* \otimes N), \mathbb{Z}). \end{aligned}$$

To show that it is an isomorphism we may assume that $M = M(X)$ and $N = M(Y)$, where X and Y are smooth projective varieties of dimensions d and e . Using 20.11 three times, and writing \mathbb{L} for $\mathbb{Z}(1)[2]$, we have:

$$M^* \otimes N^* \otimes \mathbb{L}^{d+e} \cong (M^* \otimes \mathbb{L}^d) \otimes (N^* \otimes \mathbb{L}^e) \cong M \otimes N \cong (M \otimes N)^* \otimes \mathbb{L}^{d+e}.$$

Since this isomorphism is our natural map, we are done. \square

Using the dual, we can now show that \mathbf{DM}_{gm} has an internal Hom functor.

PROPOSITION 20.14. *Let L , M , and N be three objects of \mathbf{DM}_{gm} . Then there is a natural isomorphism*

$$\mathrm{Hom}_{gm}(L \otimes M, N) \cong \mathrm{Hom}_{gm}(L, M^* \otimes N).$$

PROOF. Using proposition 20.12, which states that $M \cong M^{**}$ and $N \cong N^{**}$, 20.8 and 20.13, we have:

$$\begin{aligned} \mathrm{Hom}_{gm}(L \otimes M, N) &\cong \mathrm{Hom}_{gm}(L \otimes M \otimes N^*, \mathbb{Z}) \\ &\cong \mathrm{Hom}_{gm}(L \otimes (M^* \otimes N)^*, \mathbb{Z}) \cong \mathrm{Hom}_{gm}(L, M^* \otimes N). \quad \square \end{aligned}$$

Proposition 20.14 says that $M^* \otimes N$ represents the functor $L \mapsto \mathrm{Hom}_{gm}(L \otimes M, N)$. This justifies the following definition.

DEFINITION 20.15. If M and N are two objects of \mathbf{DM}_{gm} , we define their internal Hom to be:

$$\underline{Hom}(M, N) = M^* \otimes N.$$

By 20.6, $\underline{Hom}(M, N)$ is a geometric motive, i.e., an object of \mathbf{DM}_{gm} . Moreover, it is clear that $\underline{Hom}(M, \mathbb{Z}) = M^*$.

EXERCISE 20.16. To see the relation between \underline{Hom} and \underline{RHom} , let M and N be two effective geometric motives. First show that $\underline{RHom}(M, N(i))$ is in $\mathbf{DM}_{gm}^{\mathrm{eff}}$ for large i . Then show that $\underline{Hom}(M, N) = \underline{RHom}(M, N(i))(-i)$ for large i .

Recall from [DMOS82, p. 111] that a tensor category \mathcal{A} is said to be **rigid** if it has an internal Hom, bi-distributive for the tensor, and if $A \rightarrow (A^*)^*$ is an isomorphism for every A .

THEOREM 20.17. *The tensor category \mathbf{DM}_{gm} is rigid.*

PROOF. We have already shown in 20.15 that \mathbf{DM}_{gm} has an internal Hom and in 20.12 that every object is isomorphic to its double dual. It remains to check bi-distributivity. But this is just the routine calculation:

$$\begin{aligned} \underline{Hom}(M_1 \otimes M_2, N_1 \otimes N_2) &= (M_1 \otimes M_2)^* \otimes (N_1 \otimes N_2) \\ &\cong (M_1^* \otimes N_1) \otimes (M_2^* \otimes N_2) = \underline{Hom}(M_1, N_1) \otimes \underline{Hom}(M_2, N_2). \quad \square \end{aligned}$$

Part 6

Zariski Sheaves with Transfers

Covering morphisms of triples

The main goal of the rest of the lectures will be to prove that if F is a homotopy invariant presheaf with transfers, then the presheaf $H_{Nis}^n(-, F)$ is homotopy invariant. This was stated in theorem 13.8 and it was used in lectures 13-20. The remaining lectures depend upon lectures 11, 12, and the first part of 13 (13.1–13.5), but not on the material from 13.7 to the end of lecture 20.

DEFINITION 21.1. Let $T_Y = (\bar{Y}, Y_\infty, Z_Y)$ and $T_X = (\bar{X}, X_\infty, Z_X)$ be standard triples (as defined in 11.5). For convenience, set $Y = \bar{Y} - Y_\infty$ and $X = \bar{X} - X_\infty$. A **covering morphism** $f : T_Y \rightarrow T_X$ of standard triples is a finite morphism $f : \bar{Y} \rightarrow \bar{X}$ such that:

- $f^{-1}(X_\infty) \subset Y_\infty$ (or equivalently, $f(Y) \subset X$);
- $f|_Y : Y \rightarrow X$ is étale;
- f induces an isomorphism $Z_Y \xrightarrow{\cong} Z_X$, where $Z_Y = f^{-1}(Z_X) \cap Y$.

Note that f need not induce a finite morphism $f : Y \rightarrow X$.




FIGURE 21.1. A covering morphism $f : \bar{Y} \rightarrow \bar{X}$

By definition, the square $Q = Q(X, Y, X - Z_X)$ induced by a covering morphism of standard triples is upper distinguished (see 12.5):

$$\begin{array}{ccc} (Y - Z_Y) & \longrightarrow & Y \\ \downarrow & & \downarrow f \\ (X - Z_X) & \longrightarrow & X. \end{array}$$

We say that this upper distinguished square **comes from** the covering morphism of standard triples.

EXAMPLE 21.2. Suppose that an affine X has a covering $X = U \cup V$ and a good compactification (\bar{X}, X_∞) over some smooth S . Then the Zariski square

$$Q(X, U, V) : \begin{array}{ccc} U \cap V & \longrightarrow & U \\ \downarrow & & \downarrow \\ V & \longrightarrow & X \end{array}$$

comes from a morphism of triples, provided that $\bar{X} - (U \cap V)$ lies in an affine open neighborhood in \bar{X} .

Indeed, if $Z = X - V$ then $T = (\bar{X}, X_\infty, Z)$ is a standard triple and $T' = (\bar{X}, \bar{X} - U, Z)$ is also a standard triple. The identity on \bar{X} induces a covering morphism $T' \rightarrow T$ and the above square comes from this morphism.

Recall from 11.11 that a splitting of a standard triple (\bar{X}, X_∞, Z) over $V \subset X$ is a trivialization of \mathcal{L}_{Δ_X} on $V \times_S Z$.

LEMMA 21.3. *Let $f : T_Y \rightarrow T_X$ be a covering morphism of standard triples. A splitting of T_X over V induces a splitting of T_Y over $f^{-1}(V) \cap Y$.*

PROOF. Since T_X was split over $V \subseteq \bar{X}$, we are given $t : \mathcal{L}_{\Delta_X}|_{V \times_S Z} \cong \mathcal{O}$. We need a trivialization

$$f^{-1}(t) : \mathcal{L}_{\Delta_Y}|_{f^{-1}(V) \times_S Z_Y} \cong \mathcal{O}.$$

Now $(f \times f)^{-1}(\Delta_X)$ is the disjoint union of Δ_Y and some Q , so $(f \times f)^*(\mathcal{L}_{\Delta_X})$ is $\mathcal{L}_{\Delta_Y} \otimes \mathcal{L}_Q$, where \mathcal{L}_Q is the associated line bundle. Since f induces an isomorphism $Z_Y \rightarrow Z_X$, Q is disjoint from $Y \times_S Z_Y$. Since \mathcal{L}_Q has a canonical trivialization outside Q , we have $\mathcal{L}_Q \cong \mathcal{O}$ on $Y \times_S Z_Y$. Since $(f \times f)^*(t)$ is a trivialization of $\mathcal{L}_{\Delta_Y} \otimes \mathcal{L}_Q$ on $(f \times f)^{-1}(V \times_S Z_X)$, we may regard $(f \times f)^*(t)$ as a trivialization of \mathcal{L}_{Δ_Y} on $(f^{-1}(V) \cap Y) \times_S Z_Y$. \square

EXAMPLE 21.4. Let $\bar{Y} \rightarrow \bar{X}$ be a finite separable morphism of smooth projective curves, $X_\infty \subset \bar{X}$ a finite nonempty set containing the branch locus, and $y \in \bar{Y}$ a k -rational point so that $x = f(y)$ is not in X_∞ . Set $Y_\infty = f^{-1}(X_\infty) \amalg f^{-1}(x) - \{y\}$. Then $(\bar{Y}, Y_\infty, \{y\}) \rightarrow (\bar{X}, X_\infty, \{x\})$ is a covering morphism of standard triples. If $X = \text{Spec } A$ and P is the prime ideal of A defining x , then PB is prime in the coordinate ring B of Y . If $a \in A$ then by 11.13, lemma 21.3 states that if $P[1/a]$ is principal, then so is $PB[1/a]$.

DEFINITION 21.5. Let Q be any commutative square of the form

$$\begin{array}{ccc} B & \xrightarrow{i} & Y \\ \downarrow f & & \downarrow f \\ A & \xrightarrow{i} & X. \end{array}$$

We write $MV(Q)$ for the following chain complex in Cor_k :

$$MV(Q) : 0 \longrightarrow B \xrightarrow{(-f,i)} A \oplus Y \xrightarrow{(i,f)} X \longrightarrow 0.$$

If F is a presheaf, then $F(MV(Q))$ is the complex of abelian groups:

$$0 \longrightarrow F(X) \xrightarrow{(i,f)} F(A) \oplus F(Y) \xrightarrow{(-f,i)} F(B) \longrightarrow 0.$$

The general theorem below will involve an intricate set of data which we now describe. Let f be a covering morphism of standard triples, from $T_Y = (\bar{Y}, Y_\infty, Z_Y)$ to $T_X = (\bar{X}, X_\infty, Z_X)$. Let Q denote the square that comes from f . Let $Q' = (X', Y', A')$ be another upper distinguished square with Y' and X' affine so that Q and Q' are of the form:

$$(21.5.1) \quad Q' : \begin{array}{ccc} B' & \longrightarrow & Y' \\ \downarrow f' & & \downarrow f' \\ A' & \xrightarrow{i'} & X' \end{array} \quad Q : \begin{array}{ccc} B & \longrightarrow & Y \\ \downarrow f & & \downarrow f \\ A & \xrightarrow{i} & X. \end{array}$$

THEOREM 21.6. *Let $j : Q' \rightarrow Q$ be a morphism of upper distinguished squares of the form 21.5.1 such that:*

- Q comes from a covering morphism $T_Y \rightarrow T_X$ of standard triples;
- $X' \rightarrow X$ is an open embedding, and (\bar{X}, X_∞, Z_X) splits over X' ;
- X' and Y' are affine.

Then for any homotopy invariant presheaf with transfers F , the map of complexes $F(MV(Q)) \rightarrow F(MV(Q'))$ is chain homotopic to zero.

$$\begin{array}{ccccccc} 0 & \longrightarrow & F(X) & \xrightarrow{(i,f)} & F(A) \oplus F(Y) & \xrightarrow{(-f,i)} & F(B) \longrightarrow 0 \\ & & \downarrow j_X & & \downarrow \begin{pmatrix} j_A \\ j_Y \end{pmatrix} & & \downarrow j_B \\ 0 & \longrightarrow & F(X') & \xrightarrow{(i',f')} & F(A') \oplus F(Y') & \xrightarrow{(-f',i')} & F(B') \longrightarrow 0 \end{array}$$

The proof of 21.6 will be assembled from lemmas 21.7, 21.8 and 21.9 below.

We say that a diagram in Cor_k is **homotopy commutative** if every pair of composites $f, g : X \rightarrow Y$ with the same source and target are \mathbb{A}^1 -homotopic. Any homotopy invariant presheaf with transfers identifies \mathbb{A}^1 -homotopic maps, and converts a homotopy commutative diagram into a commutative diagram.

LEMMA 21.7. *Let $j : Q' \rightarrow Q$ be as in the statement of 21.6. Then there are maps $\lambda_A \in Cor(X', A)$ and $\lambda_B \in Cor(Y', B)$, well-defined up to \mathbb{A}^1 -homotopy, such*

that the following diagram is homotopy commutative.

$$\begin{array}{ccccc}
 & & Y' & \xrightarrow{f'} & X' \\
 & \swarrow j_Y & \downarrow \exists \lambda_B & & \downarrow \exists \lambda_A & \searrow j_X \\
 Y & \xleftarrow{i} & B & \xrightarrow{f} & A & \xrightarrow{i} & X
 \end{array}$$

Applying a homotopy invariant presheaf with transfers F gives a commutative diagram:

$$\begin{array}{ccccccc}
 F(X) & \xrightarrow{i} & F(A) & \xrightarrow{f} & F(B) & \xleftarrow{i} & F(Y) \\
 & \searrow j_X & \downarrow \exists \lambda_A & & \downarrow \exists \lambda_B & & \swarrow j_Y \\
 & & F(X') & \xrightarrow{f'} & F(Y') & &
 \end{array}$$

The assertion in the lemma that λ_A is only well defined up to \mathbb{A}^1 -homotopy equivalence reflects the identification

$$\text{Cor}(X', A) / \mathbb{A}^1\text{-h.e.} = H_0^{\text{sing}}(X' \times_S A / X') \xrightarrow{\cong} \text{Pic}(X' \times_S \bar{X}, X' \times_S (X_\infty \amalg Z_X))$$

arising from 7.2 and 7.16. A similar remark applies to the indeterminacy of λ_B .

PROOF. By 21.3, both triples T_X and T_Y split over an affine. Hence the maps in question exist and the outer triangles commute up to \mathbb{A}^1 -homotopy by 11.15. The construction of the relative Picard classes representing λ_A and λ_B from the compatible splittings in the proof of 11.15 shows that the middle square is homotopy commutative. \square

Since $\text{Cor}(X, Y) / \mathbb{A}^1\text{-homotopy} = H_0^{\text{sing}}(X \times Y / X)$ by 7.2, two elements of $\text{Cor}(X, Y)$ are \mathbb{A}^1 -homotopic exactly when they agree in $H_0^{\text{sing}}(X \times Y / X)$. This allows us to apply the relative Picard techniques of lecture 7.

LEMMA 21.8. *Let h be a rational function on $\bar{X} \times_S \bar{Y}$ which is invertible in a neighborhood U of $A' \times_S Y_\infty$ and $A' \times_S Z_Y$, and equals 1 on $A' \times_S Y_\infty$. Then the Weil divisor D defined by h defines an element ψ of $\text{Cor}(A', B)$ such that the composition $i\psi \in \text{Cor}(A', Y)$ is \mathbb{A}^1 -homotopic to zero.*

PROOF. As a divisor on the normal variety $A' \times_S \bar{Y}$, we can write $D = \sum n_i D_i$ with each D_i integral and supported off of U . Since each D_i misses $A' \times_S Y_\infty$, it is quasi-finite over A' . Since D_i is proper over A' , and has the same dimension as A' , it is finite and surjective over A' . As such, each D_i and hence D defines an element of $C_0(A' \times_S B / A')$ which is a subgroup of $C_0(A' \times_S B / A') = \text{Cor}(A', B)$. By construction (see 7.15), the image of D in $\text{Pic}(A' \times_S \bar{Y}, A' \times_S (Y_\infty \amalg Z))$ is given by (\mathcal{O}, h) , the trivial line bundle with trivialization 1 on $A' \times_S Y_\infty$, and h on $A' \times_S Z_Y$. The composition with $i : B \rightarrow Y$ sends D to an element of $C_0(A' \times_S Y / A')$

whose image in $\text{Pic}(A' \times_S \bar{Y}, A' \times_S Y_\infty)$ is the class of (\mathcal{O}, h) . By 7.16, this group is isomorphic to $H_0^{\text{sing}}(A' \times_S B/A')$. But in this group $(\mathcal{O}, h) = (\mathcal{O}, 1)$ is the zero element. This implies that the image is zero in $H_0^{\text{sing}}(A' \times B/A')$. \square

LEMMA 21.9. *Let $j : Q' \rightarrow Q$ be as in the statement of 21.6. Then there are $\lambda_A \in \text{Cor}(X', A)$, $\lambda_B \in \text{Cor}(Y', B)$, satisfying the conditions of lemma 21.7, and a ψ in $\text{Cor}_k(A', B)$ fitting into a homotopy commutative diagram in Cor_k :*

$$\begin{array}{ccc}
 B' & \xrightarrow{\lambda_B \circ i' - j_B} & B \\
 \downarrow f' & \nearrow \psi & \downarrow f \\
 A' & \xrightarrow{\lambda_A \circ i' - j_A} & A.
 \end{array}$$

Moreover the composition $A' \xrightarrow{\psi} B \xrightarrow{i} Y$ is \mathbb{A}^1 -homotopic to 0.

Applying a homotopy invariant presheaf with transfers F gives a commutative diagram:

$$\begin{array}{ccc}
 F(A) & \xrightarrow{f} & F(B) \\
 \downarrow i' \circ \lambda_A - j_A & \nearrow \psi & \downarrow i' \circ \lambda_B - j_B \\
 F(A') & \xrightarrow{f'} & F(B'),
 \end{array}$$

and the composite $F(Y) \xrightarrow{i} F(B) \xrightarrow{\psi} F(A')$ is zero.

PROOF OF 21.9. In order to streamline notation, we write \times for \times_S .

Let $\mathcal{L}_{\Delta X'}$ be the line bundle on $X' \times \bar{X}$ corresponding to the graph $\Delta X'$ of $X' \hookrightarrow \bar{X}$, and $\mathcal{L}_{\Delta Y'}$ for the line bundle on $Y' \times \bar{Y}$ corresponding to the graph $\Delta Y'$ of $Y' \hookrightarrow \bar{Y}$. In between these, we have the line bundle \mathcal{M} on $X' \times \bar{Y}$, obtained by pulling back $\mathcal{L}_{\Delta X'}$.

Since these three line bundles come from effective divisors, they have canonical global sections. We will write s_X for the canonical global section of $\mathcal{L}_{\Delta X'}$ on $X' \times \bar{X}$, $s_{\mathcal{M}}$ for \mathcal{M} on $X' \times \bar{Y}$, and s_Y for $\mathcal{L}_{\Delta Y'}$ on $Y' \times \bar{Y}$. Each global section determines a section on $X' \times Z_X$, $X' \times Z_Y$, and $Y' \times Z_Y$, respectively. Since $A' \subseteq X' - Z_X$ and $B' \subseteq Y' - Z_Y$, the restrictions of $s_X, s_{\mathcal{M}}, s_Y$ also determine trivializations in each case, of $\mathcal{L}_{\Delta X'}$ on $A' \times Z_X$, of \mathcal{M} on $A' \times Z_Y$, and of $\mathcal{L}_{\Delta Y'}$ on $B' \times Z_Y$.

Because $Z_Y \cong Z_X$, the inclusion of $X' \times Z_X$ in $X' \times \bar{X}$ lifts to $X' \times \bar{Y}$, and we may identify the pullbacks of $\mathcal{L}_{\Delta X'}$ and \mathcal{M} to $X' \times Z_Y$, together with their respective trivializations s_X and $s_{\mathcal{M}}$ on $A' \times Z_Y$.

Since the standard triple (\bar{X}, X_∞, Z_X) splits over X' , we are given a fixed trivialization t_X of $\mathcal{L}_{\Delta X'}$ on $X' \times Z_X$. As with s_X , we may identify t_X with a trivialization

$t_{\mathcal{M}}$ of \mathcal{M} on $X' \times Z_Y$. By 21.3, t_X also induces a trivialization t_Y of $\mathcal{L}_{\Delta Y}$ on $Y' \times Z_Y$. Since Z_X lives in an affine neighborhood U_X in \bar{X} , we extend t_X to $X' \times U_X$ and we fix this particular extension. Pulling back, the same is true for $t_{\mathcal{M}}$ and t_Y and we fix those two extensions too.

Because $t_X, t_{\mathcal{M}}, t_Y$ are trivializations, there are regular functions $r_X, r_{\mathcal{M}}, r_Y$ so that:

$$s_X = r_X t_X \text{ on } X' \times Z_X; \quad s_{\mathcal{M}} = r_{\mathcal{M}} t_{\mathcal{M}} \text{ on } X' \times Z_Y; \quad s_Y = r_Y t_Y \text{ on } Y' \times Z_Y.$$

Because s_X is a trivialization on $A' \times Z_X$, r_X is invertible on $A' \times Z_X$. Similarly, $r_{\mathcal{M}}$ is invertible on $A' \times Z_Y$, and r_Y is invertible on $B' \times Z_Y$. (See figure 21.2.)




FIGURE 21.2. The covering morphism $f : \bar{Y} \rightarrow \bar{X}$ over A'

Because $(\bar{Y}, Y_{\infty}, Z_Y)$ is a standard triple, there is an affine open neighborhood U of $Y_{\infty} \amalg Z_Y$ in \bar{Y} . Hence $X' \times U$ is an affine open neighborhood of $X' \times Z_Y$ and $X' \times Y_{\infty}$ in $X' \times \bar{Y}$. Since Z_Y and Y_{∞} are disjoint, the Chinese Remainder Theorem yields a regular function h on $X' \times U$ which equals 1 on $X' \times Y_{\infty}$ and equals $r_{\mathcal{M}}$ on $X' \times Z_Y$. Let $D \subset X' \times \bar{Y}$ denote the principal divisor corresponding to h . By lemma 21.8, the divisor $-D$ defines an element ψ of $Cor(A', B)$ such that the composition $i\psi \in Cor(A', Y)$ is homotopically trivial. By 7.15, the map $Cor(A', B) \rightarrow Pic(A' \times \bar{Y}, A' \times (Y_{\infty} \amalg Z_Y))$ sends ψ to the class of $(\mathcal{O}_{A' \times \bar{Y}}, 1_{\infty} \amalg r_{\mathcal{M}}^{-1})$.

It remains to verify that the diagram in 21.9 is homotopy commutative.

We first interpret the horizontal maps in 21.9. By the construction of λ_A and λ_B in 11.15 and 21.7, the compositions $\lambda_A \circ i' \in Cor(A', A)$ and $\lambda_B \circ i' \in Cor(B', B)$ represent the classes of $(\mathcal{L}_{\Delta A'}, s_{\infty} \amalg t_X)$ and $(\mathcal{L}_{\Delta B'}, s_{\infty} \amalg t_Y)$ in $Pic(A' \times \bar{X}, A' \times (X_{\infty} \amalg Z_Y))$ and $Pic(B' \times \bar{Y}, B' \times (Y_{\infty} \amalg Z_Y))$, respectively. On the other hand, the inclusions j_A and j_B represent the classes of $(\mathcal{L}_{\Delta A'}, s_{\infty} \amalg s_X)$ and $(\mathcal{L}_{\Delta B'}, s_{\infty} \amalg s_Y)$, respectively. It follows that the differences $j_A - \lambda_A \circ i' \in Cor(A', A)$ and $j_B - \lambda_B \circ i' \in Cor(B', B)$ represent the classes of $(\mathcal{O}_{A' \times \bar{X}}, 1_{\infty} \amalg r_X)$ and $(\mathcal{O}_{B' \times \bar{Y}}, 1_{\infty} \amalg r_Y)$, respectively (cf. exercise 11.16).

The composition $\psi f' \in \text{Cor}(B', B)$ represents $(\mathcal{O}_{B' \times \bar{Y}}, f'^* h^{-1})$. Since $f^* h$ is a rational function on $B' \times \bar{Y}$ which is 1 on $B' \times Y_\infty$ and r_Y on $B' \times Z_Y$, we have $\psi f' = \lambda_B \circ i' - j_B$ in $\text{Pic}(B' \times \bar{Y}, B' \times (Y_\infty \amalg Z_Y))$.

Now the composition $f\psi \in \text{Cor}(A', A)$ represents the push-forward of ψ along $H_0(A' \times B/A') \rightarrow H_0(A' \times A/A')$. By 7.24, this represents the class of $(\mathcal{O}_{A' \times \bar{X}}, f_*(1_\infty \amalg r_{\mathcal{M}}^{-1}))$. By definition 7.22, the norm of h is a rational function which extends the trivialization $f_*(1_\infty \amalg r_{\mathcal{M}})$ to an affine neighborhood. Since h is identically 1 on $f^{-1}(X_\infty) \subset Y_\infty$, $N(h) = 1$ on $A' \times X_\infty$ by 7.23. We will show that $N(h) = r_X$ on $A' \times Z_X$ in lemma 21.10 below. Hence $f\psi = \lambda_{A'} i' - j_A$ in $\text{Cor}(A', A)$, as desired. \square

LEMMA 21.10. *Let $f : U \rightarrow V$ be a finite map with U and V normal. Suppose that $Z \subset V$ and $Z' \subset U$ are reduced closed subschemes such that the induced map $Z' \rightarrow Z$ is an isomorphism, and $U \rightarrow V$ is étale in a neighborhood of Z' .*

If $h \in \mathcal{O}^(U)$ is 1 on $f^{-1}(Z) - Z'$, then $N(h)|_Z$ and $h|_{Z'}$ are identified by $Z' \cong Z$.*

PROOF. Suppose first that f has a section $s : V \rightarrow U$ sending Z to Z' . Then $U \cong s(U) \amalg U'$ and h is 1 on $f^{-1}(Z) \cap U'$. In this case, the assertion follows from the componentwise calculation of the norm $N(h)$, together with 7.23.

In the general case, let $U' \subset U$ be a neighborhood of Z' which is étale over V , and let $h' \in \mathcal{O}^*(U' \times_V U)$ be the pullback of h . The graph $Z'' \subset U' \times_V U$ of $Z' \rightarrow Z$ is isomorphic to Z' , and $U' \times_V U'$ is an étale neighborhood of Z'' in $U' \times_V U$. By construction, h' is 1 on $U' \times_V (f^{-1}(Z) - Z'')$ and $U' \times_V U \rightarrow U'$ has a canonical section sending Z' to Z'' ; in this case we have shown that $N(h')|_{Z''}$ is identified with $h|_{Z'}$. Since norms commute with base change, we can identify $N(h)$ with $N(h')$ under $\mathcal{O}^*(V) \subseteq \mathcal{O}^*(U')$. This proves the lemma. \square

PROOF OF 21.6. From 21.7 and 21.8, we have maps $s_1 = (\lambda_A, 0) : F(A) \oplus F(Y) \rightarrow F(X')$ and $s_2 = (\psi, \lambda_B) : F(B) \rightarrow F(A') \oplus F(Y')$. In order for these maps to form a chain homotopy from j to zero, we must have $sd + ds = j$. This amounts to six equations, three of which come from the commutativity of the trapezoid in 21.7. The other three, which involve ψ are: $\psi i \simeq 0$, $j_A \simeq i' \lambda_A - \psi f$ and $j_B \simeq i' \lambda_B - f' \psi$. These are provided by 21.9. \square

We isolate a special case of theorem 21.6 as a corollary, which will be needed in the proof of theorem 22.2.

COROLLARY 21.11. *Let $Q = Q(X, Y, A)$ be an upper distinguished square of smooth schemes coming from a covering morphism of standard triples and let Σ be a finite set of points in Y . Then there exist affine neighborhoods X' of $f(\Sigma)$ in X and Y' of Σ in $Y \cap f^{-1}(X')$ such that:*

- *The induced square $Q' = Q(X', Y', A')$ is upper distinguished, where $A' = A \cap X'$ and $B' = B \cap Y'$;*

- For any homotopy invariant presheaf with transfers F , the map $F(MV(Q)) \rightarrow F(MV(Q'))$ is chain homotopic to zero.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F(X) & \longrightarrow & F(A) \oplus F(Y) & \longrightarrow & F(B) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & F(X') & \longrightarrow & F(A') \oplus F(Y') & \longrightarrow & F(B') \longrightarrow 0
 \end{array}$$

PROOF. By 11.14, $f(\Sigma)$ has an affine neighborhood X' over which the triple (\bar{X}, X_∞, Z) splits. Set $\bar{Y}' = X' \times_{\bar{X}} \bar{Y}$, $Z'_X = Z_X \cap X'$ and $Z'_Y = Z_Y \cap \bar{Y}'$, and note that $Z'_Y \rightarrow Z'_X$ is an isomorphism by 21.1. Since $\bar{Y}' \rightarrow X'$ is finite, \bar{Y}' is affine. The subsets $Y_\infty \cap \bar{Y}'$ and $Z'_Y \cup \Sigma$ of \bar{Y}' are closed in \bar{Y}' , and disjoint by 11.5. Hence there is an affine open subscheme Y' of \bar{Y}' which contains both Z'_Y and Σ but is disjoint from $Y_\infty \cap \bar{Y}'$. Since Y' is open in Y , it is étale over X' . Since $B' = B \cap Y'$ is the complement in Y' of $Z'_Y = Z_Y \cap Y'$, and $B' = A \times_X Y' = A' \times_{X'} Y'$, the square $Q' = Q(X', Y', A')$ is upper distinguished (see 12.5). Thus the hypotheses of theorem 21.6 are satisfied for $Q' \rightarrow Q$, and the final part of the corollary is the conclusion of 21.6. \square

Zariski sheaves with transfers

With the technical results of the last lecture in hand, we are ready to prove the following results.

THEOREM 22.1. *Let F be a homotopy invariant presheaf with transfers. Then the Zariski sheaf F_{Zar} is homotopy invariant.*

THEOREM 22.2. *Let F be a homotopy invariant presheaf with transfers. Then $F_{Zar} = F_{Nis}$.*

Combining 22.1 and 22.2, we obtain theorem 22.3 below, which is the case $n = 0$ of theorem 13.8. This theorem does not require k to be perfect.

THEOREM 22.3. *If F is a homotopy invariant presheaf with transfers, then the Nisnevich sheaf F_{Nis} is homotopy invariant.*

We will prove theorems 22.1 and 22.2 in order, using a sequence of lemmas. We make the running assumption that F is a homotopy invariant presheaf with transfers. The Mayer-Vietoris sequence $F(MV(Q))$ associated to a commutative square Q is defined in 21.5.

LEMMA 22.4. *Let U be an open subset of \mathbb{A}^1 and $U = U_1 \cup U_2$ be a Zariski covering of U . Then the complex $F(MV(Q))$ is split exact, where $Q = Q(U, U_1, U_2)$.*

$$F(MV(Q)) : 0 \longrightarrow F(U) \longrightarrow F(U_1) \oplus F(U_2) \longrightarrow F(U_1 \cap U_2) \longrightarrow 0.$$

In particular, F is a Zariski sheaf on \mathbb{A}^1 .

PROOF. Setting $Y_\infty = \mathbb{P}^1 - U$, $Y'_\infty = \mathbb{P}^1 - U_1$ and $Z = U - U_2$, the identity of \mathbb{P}^1 is a covering morphism $(\mathbb{P}^1, Y'_\infty, Z) \rightarrow (\mathbb{P}^1, Y_\infty, Z)$ of standard triples as in example 21.2. Both triples are split over U itself by 11.13, so by theorem 21.6 with $Q' = Q$, the complex $F(MV(Q))$ is chain contractible, i.e., split exact. \square

LEMMA 22.5. *If F is a homotopy invariant Zariski sheaf with transfers, and U is an open subset of \mathbb{A}^1 , then $H_{Zar}^n(U, F) = 0$ for $n > 0$.*

PROOF. If $\mathcal{U} = \{U_1, \dots, U_n\}$ is a finite cover of U , it follows from 22.4 and induction on n that the following sequence is exact.

$$0 \rightarrow F(U) \rightarrow \oplus_i F(U_i) \rightarrow \oplus_{i,j} F(U_i \cap U_j) \rightarrow \dots \rightarrow F(\cap_i U_i) \rightarrow 0$$

Hence the Čech cohomology of F satisfies $\check{H}^i(\mathcal{U}, F) = 0$ for $i > 0$. But then $H^1(U, F) = \check{H}^1(U, F) = 0$ by [Har77, Ex III.4.4]. Since $\dim U = 1$, we must also have $H^i(U, F) = 0$ for $i > 1$ (see [Har77, III.2.7]). \square

EXERCISE 22.6. Show that 22.4 and 22.5 fail for $F = \mathcal{O}^*$ if \mathbb{A}^1 is replaced by an affine elliptic curve.

LEMMA 22.7. *If F is a homotopy invariant Nisnevich sheaf with transfers, and U is an open subset of \mathbb{A}^1 , then $H_{Nis}^n(U, F) = 0$ for $n > 0$.*

PROOF. Since $\dim U = 1$, we have $H_{Nis}^n(U, F) = 0$ for $n > 1$. By [Mil80, III.2.10], $H_{Nis}^1(U, F) = \check{H}^1(U, F)$. Therefore we only need to show that $\check{H}^1(U, F) = 0$.

Since F takes disjoint unions to direct sums, the Čech cohomology can be computed using covering families $V \rightarrow X$, instead of the more general $\{V_i \rightarrow X\}$. By 12.6, any such cover of U has a refinement $\mathcal{U} = \{A, V\}$, where $A \subset U$ is dense open, $V \rightarrow U$ is étale, and the square $Q = Q(U, V, A)$ is upper distinguished (see 12.5). Embed V in a smooth projective curve \bar{V} finite over \mathbb{P}^1 , and set $V_\infty = \bar{V} - V$. By construction (see 21.1), Q comes from the covering morphism of standard triples $(\bar{V}, V_\infty, Z) \rightarrow (\mathbb{P}^1, U_\infty, Z)$, where $U_\infty = \mathbb{P}^1 - U$ and $Z = U - A$. Since $(\mathbb{P}^1, U_\infty, Z)$ splits over U by 11.13, theorem 21.6 with $Q' = Q$ implies that the complex $F(MV(Q))$ is split exact. That is $\check{H}^1(\mathcal{U}, F) = 0$. Passing to the limit over all such covers yields $\check{H}^1(U, F) = 0$. \square

LEMMA 22.8. *Let F be a homotopy invariant presheaf with transfers. If X is smooth and $U \subset X$ is dense open, then $F_{Zar}(X) \rightarrow F_{Zar}(U)$ is injective.*

PROOF. As F_{Zar} is a sheaf it suffices to verify this locally. Let $f \in F_{Zar}(X)$ be a nonzero section which vanishes in $F_{Zar}(U)$. Pick a point $x \in X$ so that f is nonzero in the stalk $F_x = F(\text{Spec } \mathcal{O}_{X,x})$. By shrinking X around x we may assume that $f \in F(X)$. By shrinking U , we may assume that f vanishes in $F(U)$ and hence in $F(V)$ for $V = \text{Spec}(\mathcal{O}_{X,x}) \cap U$. By 11.1, f is nonzero in $F(V)$, and this is a contradiction. \square

PROOF OF 22.1. We have to prove that $i^* : F_{Zar}(X \times \mathbb{A}^1) \rightarrow F_{Zar}(X)$ is an isomorphism, where $i : X \rightarrow X \times \mathbb{A}^1$. It is enough to prove that i^* is injective. We may assume that X is connected and therefore irreducible. Let $\gamma : \text{Spec } K \rightarrow X$ be the generic point. We get a diagram:

$$\begin{array}{ccc} F_{Zar}(X \times \mathbb{A}^1) & \xrightarrow{i^*} & F_{Zar}(X) \\ (\gamma \times 1)^* \downarrow & & \downarrow \gamma^* \\ F_{Zar}(\text{Spec } K \times \mathbb{A}^1) & \xrightarrow{\cong} & F_{Zar}(\text{Spec } K) \end{array}$$

where the vertical maps are injective by 22.8. The bottom map is an isomorphism by 22.4 since we may regard F as a homotopy invariant presheaf with transfers over the field K by 2.10:

$$F_{Zar}(\mathbb{A}_K^1) = F(\mathbb{A}_K^1) \xrightarrow{\cong} F(\text{Spec } K) = F_{Zar}(\text{Spec } K).$$

Thus i^* is injective. \square

Let $s_{Zar}(F)$ be the separated presheaf (with respect to the Zariski topology) associated to the presheaf F . It is defined by the formula:

$$s_{Zar}(F)(X) = F(X)/F_0(X), \quad F_0(X) = \operatorname{colim}_{\substack{\text{covers} \\ \{U_i \rightarrow X\}}} \ker F(X) \rightarrow \prod F(U_i).$$

LEMMA 22.9. $s_{Zar}(F)$ is a homotopy invariant presheaf with transfers.

PROOF. The homotopy invariance of $s_{Zar}F$ is immediate from the fact that homotopy invariance is preserved by quotient presheaves. The existence of transfers is more difficult. Let $Z \subset S \times X$ be an elementary correspondence from S to X . We must show that the corresponding transfer $F(X) \rightarrow F(S)$ sends $F_0(X)$ to $F_0(S)$, i.e., that the image of $F_0(X)$ vanishes at each stalk $F(\operatorname{Spec} \mathcal{O}_{S,s})$. It suffices to suppose S local, so that Z is semilocal. Hence there is a semilocal subscheme X' of X with $Z \subset S \times X'$. But by 11.1, $F(X')$ injects into $F(U)$ for each dense $U \subset X'$, so $F_0(X') = 0$. Hence $F_0(X) \rightarrow F(S)$ is zero, because it factors through $F_0(X') = 0$. \square

For the next few lemmas, S will be the semilocal scheme of a smooth quasi-projective variety X at a finite set of points. Since any finite set of points lies in an affine neighborhood, we may even assume that X is affine. Clearly, S is the intersection of the filtered family of its affine open neighborhoods X_α in X .

LEMMA 22.10. Suppose that F is a homotopy invariant presheaf with transfers. Then for any open covering $S = U_0 \cup V$ there is an open $U \subset U_0$ such that $S = U \cup V$ and the sequence $F(MV(Q))$ is exact, where $Q = Q(S, U, V)$:

$$0 \rightarrow F(S) \rightarrow F(U) \oplus F(V) \rightarrow F(U \cap V) \rightarrow 0.$$

PROOF. We may assume that S is connected, since we can work separately with each component. By assumption, there are open \tilde{U}_0, \tilde{V} in X such that $U_0 = S \cap \tilde{U}_0, V = S \cap \tilde{V}$. Since \tilde{U}_0 is open in X , there is an affine open \tilde{U} contained in \tilde{U}_0 which contains the finite set of closed points of U_0 . Setting $U = S \cap \tilde{U}$, we have $S = U \cup V$. We will show that $F(MV(Q))$ is exact for the square $Q = Q(S, U, V)$.

We first suppose that k is an infinite field. For each α , set $U_\alpha = X_\alpha \cap \tilde{U}$ and $V_\alpha = X_\alpha \cap \tilde{V}$. The canonical map from Q to the square $Q_\alpha = Q(X_\alpha, U_\alpha, V_\alpha)$ induces a morphism of Mayer-Vietoris sequences, $F(MV(Q_\alpha)) \rightarrow F(MV(Q))$. It suffices to show that these morphisms are chain homotopic to zero, because $F(MV(Q))$ is the direct limit of the $F(MV(Q_\alpha))$.

Let $Z \subset X$ denote the union of $X - (\tilde{U} \cap \tilde{V})$ and the closed points of S . For each X_α , we know by 11.17 that there is an affine neighborhood X'_α of S in X_α and a standard triple $T_\alpha = (\bar{X}_\alpha, X_{\infty, \alpha}, Z_\alpha)$ with $X'_\alpha \cong \bar{X}_\alpha - X_{\infty, \alpha}$ and $Z_\alpha = X_\alpha \cap Z$. Set $U'_\alpha = X'_\alpha \cap \tilde{U}$ and $V'_\alpha = X'_\alpha \cap \tilde{V}$. Since $\bar{X}_\alpha - (U'_\alpha \cap V'_\alpha)$ lies in $X_{\infty, \alpha} \cup Z_\alpha$, it lies in an affine open subset of \bar{X}_α (by definition 11.5). By 21.2, the Zariski square $Q'_\alpha = Q(X'_\alpha, U'_\alpha, V'_\alpha)$ comes from a covering morphism of triples $T'_\alpha \rightarrow T_\alpha$.

By 11.14, the triple T_α is split over an affine neighborhood X''_α of S in X'_α . Set $U''_\alpha = X''_\alpha \cap \tilde{U}$ and $V''_\alpha = X''_\alpha \cap \tilde{V}$, and form the square $Q''_\alpha = Q(X''_\alpha, U''_\alpha, V''_\alpha)$. Since

X''_α and \tilde{U} are affine, so is U''_α . By theorem 21.6, the morphism $F(MV(Q'_\alpha)) \rightarrow F(MV(Q''_\alpha))$ is chain homotopic to zero. Since $F(MV(Q_\alpha)) \rightarrow F(MV(Q))$ factors through this morphism, it too is chain homotopic to zero.

$$\begin{array}{ccccccccc}
0 & \longrightarrow & F(X_\alpha) & \longrightarrow & F(U_\alpha) \oplus F(V_\alpha) & \longrightarrow & F(U_\alpha \cap V_\alpha) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(X'_\alpha) & \longrightarrow & F(U'_\alpha) \oplus F(V'_\alpha) & \longrightarrow & F(U'_\alpha \cap V'_\alpha) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(X''_\alpha) & \longrightarrow & F(U''_\alpha) \oplus F(V''_\alpha) & \longrightarrow & F(U''_\alpha \cap V''_\alpha) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(S) & \longrightarrow & F(U) \oplus F(V) & \longrightarrow & F(U \cap V) & \longrightarrow & 0
\end{array}$$

If k is finite, exactness follows by a transfer argument. Any element a in the homology of $F(MV(Q))$ must vanish when we pass to $Q \otimes_k k'$ for any infinite algebraic extension k' of k . Since a must vanish for some finite subextension k'_0 , a has exponent $[k'_0 : k]$. Since $[k'_0 : k]$ can be chosen to be a power of any prime, we conclude that $a = 0$. \square

Lemma 22.10 corrects [CohTh, 4.23], which omitted the passage from U_0 to U .

COROLLARY 22.11. *Let S' and S'' be semilocal schemes of a smooth quasi-projective scheme X at finite sets of points, and set $S = S' \cup S''$. Then the Mayer-Vietoris sequence $F(MV(Q))$ is exact, where $Q = Q(S, S', S'')$:*

$$0 \rightarrow F(S) \rightarrow F(S') \oplus F(S'') \rightarrow F(S' \cap S'') \rightarrow 0.$$

PROOF. Write S' as the intersection of open sets $U_\alpha \subset S$ and S'' as the intersection of open sets $V_\beta \subset S$. The sequence $F(MV(Q))$ is the direct limit of the sequences $F(MV(Q_{\alpha\beta}))$, where $Q_{\alpha\beta} = Q(S, U_\alpha, V_\beta)$. By 22.10, there are $U_{\alpha\beta} \subset U_\alpha$ such that the sequences $F(MV(Q(S, U_{\alpha\beta}, V_\beta)))$ are exact. Hence the morphisms from $F(MV(Q_{\alpha\beta}))$ to $F(MV(Q))$ are zero on homology. Passing to the direct limit, we see that the homology of $F(MV(Q))$ is zero, i.e., it is exact. \square

Note that the sequence $0 \rightarrow \mathcal{F}(S) \rightarrow \mathcal{F}(S') \oplus \mathcal{F}(S'') \rightarrow \mathcal{F}(S' \cap S'')$ is always exact when \mathcal{F} is a Zariski sheaf on S . This is because it is the direct limit of the exact sequences $0 \rightarrow \mathcal{F}(S) \rightarrow \mathcal{F}(U_\alpha) \oplus \mathcal{F}(V_\beta) \rightarrow \mathcal{F}(U_\alpha \cap V_\beta)$ associated to the family of open covers $\{U_\alpha, V_\beta\}$ of S with $S' \subset U_\alpha$ and $S'' \subset V_\beta$.

LEMMA 22.12. *Let S be the semilocal scheme of a smooth quasi-projective scheme X at a finite set of points. Then $F_{Zar}(S) = F(S)$.*

PROOF. By 11.1, $F(S) = (s_{Zar}F)(S)$. Since $s_{Zar}F$ is a homotopy invariant presheaf with transfers by 22.9, we may replace F by $s_{Zar}F$ and assume that F is separated. We now proceed by induction on the number of the closed points of S .

Let S' be the local scheme at a closed point x of S , and S'' the semilocal scheme at the remaining points. Consider the following commutative diagram.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F(S) & \longrightarrow & F(S') \oplus F(S'') & \longrightarrow & F(S' \cap S'') \\
 & & \downarrow & & \downarrow = & & \downarrow \text{into} \\
 0 & \longrightarrow & F_{Zar}(S) & \longrightarrow & F_{Zar}(S') \oplus F_{Zar}(S'') & \longrightarrow & F_{Zar}(S' \cap S'')
 \end{array}$$

The top row is exact by 22.11, and we have noted that the bottom row is exact because F_{Zar} is a Zariski sheaf. The right vertical map is an injection because F is separated. The middle vertical map is the identity by induction. A diagram chase shows that the left vertical map is an isomorphism, as desired. \square

We need an analogue of lemma 6.16 for the Zariski topology, showing that we can lift finite correspondences to open covers under mild conditions.

LEMMA 22.13. *Let W be a closed subset of $X \times Y$, $x \in X$ a point and $V \subset Y$ an open subset such that $p^{-1}(x) \subset \{x\} \times V$, where $p : W \rightarrow X$ is the projection. Then there is a neighborhood U of x such that $W \times_X U$ is contained in $U \times V$.*

PROOF. The subset $Z = W - W \cap (X \times V)$ is closed, and $x \notin p(Z)$. Because p is a closed map, $p(Z)$ is closed and $U = X - p(Z)$ is an open neighborhood of x . By construction, $W \times_X U$ is contained in $U \times V$. \square

COROLLARY 22.14. *Let $\mathscr{W} \in \text{Cor}(X, Y)$ have support W and let $p : W \rightarrow X$ be the projection. If $x \in X$ and $V \subset Y$ are such that $p^{-1}(x) \subset \{x\} \times V$, then there is a neighborhood U of x and a canonical $\mathscr{W}_U \in \text{Cor}(U, V)$ such that the following diagram commutes.*

$$\begin{array}{ccc}
 U & \xrightarrow{\mathscr{W}_U} & V \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{\mathscr{W}} & Y
 \end{array}$$

PROOF. Writing $\mathscr{W} = \sum n_i[W_i]$, we may apply lemma 22.13 to each W_i . Since W_i is finite over X , $W_i \times_X U$ is finite over U , so $\mathscr{W}_U = \sum n_i[W_i \times_X U]$ is the required finite correspondence. It is canonical because if $U' \subset U$, the composition of $U' \subset U$ with \mathscr{W}_U is $\mathscr{W}_{U'} = \sum n_i[W_i \times_X U']$. \square

THEOREM 22.15. *Let F be a homotopy invariant presheaf with transfers. Then the Zariski sheaf F_{Zar} has a unique structure of presheaf with transfers such that $F \rightarrow F_{Zar}$ is a morphism of presheaves with transfers.*

PROOF. By 22.9 we may assume that F is separated, i.e., that $F(V) \subseteq F_{Zar}(V)$ for every V . We may also assume that X and Y are irreducible without loss of generality. We begin by defining an element $\mathscr{W}^*(f)$ in $F_{Zar}(X)$ for every element $f \in F_{Zar}(Y)$ and every finite correspondence \mathscr{W} from X to Y .

The first step is to fix a point $x \in X$ and construct an element $\mathscr{W}^*(f)_x$ of $F_{Zar}(U_x)$ for an appropriate neighborhood U_x of x . Since $p: W \rightarrow X$ is finite, the image of $p^{-1}(x)$ under the natural map $W \rightarrow Y$ consists of only finitely many points; let S denote the semilocal scheme of Y at these points. Since $F(S) = F_{Zar}(S)$ by 22.12, there is an open $V_x \subset Y$ such that $f_x = f|_{V_x} \in F_{Zar}(V_x)$ lies in the subgroup $F(V_x) \subseteq F_{Zar}(V_x)$. By 22.14, there is a neighborhood U_x of x such that \mathscr{W} restricts to a finite correspondence \mathscr{W}_x from U_x to V_x . Let $\mathscr{W}^*(f)_x$ denote the image of f_x under $\mathscr{W}_x^*: F(V_x) \rightarrow F(U_x) \subseteq F_{Zar}(U_x)$.

Uniqueness of $\mathscr{W}^(f)_x$.* Suppose that $F \rightarrow F_{Zar}$ is a morphism of presheaves with transfers. Given $\mathscr{W} \in Cor(X, Y)$ and $f \in F_{Zar}(Y)$, it suffices to show that $\mathscr{W}^*(f) \in F_{Zar}(X)$ is uniquely defined in some neighborhood of any point x . The construction above shows that the image of $\mathscr{W}^*(f)$ in $F_{Zar}(U_x)$ must equal $\mathscr{W}^*(f)_x$, which is defined using only the sheaf structure on F_{Zar} and the transfer structure on F .

Existence of $\mathscr{W}^(f)_x$.* Fix $\mathscr{W} \in Cor(X, Y)$ and $f \in F_{Zar}(Y)$. In the above construction, we produced a neighborhood U_x of every point $x \in X$, an open set V_x in Y so that $f_x = f|_{V_x}$ belongs to the subgroup $F(V_x)$ of $F_{Zar}(V_x)$, and we considered the image $\mathscr{W}^*(f)_x = \mathscr{W}_x^*(f_x)$ of f_x in $F(U_x) \subseteq F_{Zar}(U_x)$. This construction corresponds to the top row of figure 22.1.




FIGURE 22.1. The transfer map for F_{Zar}

To construct the rest of figure 22.1, pick two points $x, x' \in X$ and set $U_{xx'} = U_x \cap U_{x'}$, $V_{xx'} = V_x \cap V_{x'}$. Since $W \times_X U_x$ lies in $U_x \times V_x$ for all x (by 22.13), it follows that $W \times_X U_{xx'}$ lies in $U_{xx'} \times (V_x \cap V_{x'})$. Hence there is a finite correspondence $\mathscr{W}_{xx'}$ from $U_{xx'}$ lifting both \mathscr{W}_x and $\mathscr{W}_{x'}$ in the sense of 22.14. That is, the middle square commutes in figure 22.1.

A diagram chase on 22.1 shows that the $\mathscr{W}^*(f)_x$ agree on all intersections $U_{xx'} = U_x \cap U_{x'}$. Thus the element $\mathscr{W}^*(f) \in F_{Zar}(X)$ exists by the sheaf axiom.

Fix $x \in X$ and choose $V \subset Y$, $f_V \in F(V)$ and U_x as above. Because F is separated we have $F(V) \subset F_{Zar}(V)$, so the element $f_V \in F(V)$ is well defined. Given a dense $V_0 \subset V$, the map $F(V) \rightarrow F(V_0)$ sends f_V to f_{V_0} , because $F_{Zar}(V) \subset F_{Zar}(V_0)$ by 22.8. Given $U_0 \subset U_x$, the proof of 22.14 shows that the the canonical lift

$\mathscr{W}_{U_0} \in \text{Cor}(U_0, V)$ is the composition of the inclusion $U_0 \subset U$ with the canonical lift $\mathscr{W}_U \in \text{Cor}(U, V)$. Hence $F_{\text{Zar}}(U_x) \rightarrow F_{\text{Zar}}(U_0)$ sends the element $\mathscr{W}^*(f)_x$ to the image of f_{V_0} under $F(V_0) \rightarrow F(U_0) \subset F_{\text{Zar}}(U_0)$.

It is now easy to check using 22.8 that the maps \mathscr{W}^* are additive and give F_{Zar} the structure of a presheaf with transfers. \square

PROOF OF 22.2. We have to prove that $F_{\text{Zar}} = F_{\text{Nis}}$. Let F' and F'' denote the kernel and cokernel presheaves of $F \rightarrow F_{\text{Nis}}$, respectively. By 13.1, they are presheaves with transfers whose associated Nisnevich sheaf is zero. Since sheafification is exact, it suffices to show that $F'_{\text{Zar}} = F''_{\text{Zar}} = 0$. That is, we may assume that $F_{\text{Nis}} = 0$.

By 22.1 and 22.15, F_{Zar} is also a homotopy invariant presheaf with transfers. Since $F_{\text{Nis}} = (F_{\text{Zar}})_{\text{Nis}}$, we may assume that $F = F_{\text{Zar}}$, i.e., that F is a Zariski sheaf. Therefore it suffices to show that $F(S) = 0$ for every local scheme S of a smooth variety X . Let S be the local scheme associated to a point x of X .

By 12.7, it suffices to check that, for any upper distinguished square

$$Q: \begin{array}{ccc} B & \xrightarrow{i} & Y \\ \downarrow f & & \downarrow f \\ A & \xrightarrow{i} & X \end{array}$$

(see definition 12.5), the square $F(Q \times_X S)$ is a pullback. By 21.5, this is equivalent to checking that the complex $F(MV(Q \times_X S))$ is exact. This is evident if $x \in A$, when $A \times_X S = S$ and $B \times_X S = Y \times_X S$, so we may assume that $x \in Z_X$.

Shrinking X around x , we may suppose by 11.17 that X is affine and fits into a standard triple (\bar{X}, X_∞, Z) with $A = X - Z$. Shrinking Y around the finite set $\Sigma = f^{-1}(x)$, we may also suppose by 11.17 that Y is affine, and fits into a standard triple so that Q comes from a covering morphism of standard triples in the sense of 21.1. Hence 21.11 implies that $Q \times_X S \rightarrow Q$ factors through an upper distinguished square Q' in such a way that

$$F(MV(Q)) \rightarrow F(MV(Q')) \rightarrow F(MV(Q \times_X S))$$

is chain homotopic to zero.

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F(X) & \longrightarrow & F(A) \oplus F(Y) & \longrightarrow & F(B) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & F(X') & \longrightarrow & F(A') \oplus F(Y') & \longrightarrow & F(B') & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & F(S) & \longrightarrow & F(A \cap S) \oplus F(Y \times_X S) & \longrightarrow & F(B \times_X S) & \longrightarrow & 0 \end{array}$$

Taking the limit over smaller and smaller neighborhoods X of x , we see that $F(MV(Q \times_X S))$ is exact. But then $F(Q \times_X S)$ is a pullback square, as claimed. \square

LECTURE 23

Contractions

We need one final tool in order to prove theorem 13.8, which says that Nisnevich cohomology preserves homotopy invariance for sheaves with transfers. In this lecture we associate to F a new presheaf F_{-1} (known as the **contraction** of F in the literature). Here is the definition.

Let F be a homotopy invariant presheaf. The presheaf F_{-1} is defined by the formula:

$$F_{-1}(X) = \text{coker} \left(F(X \times \mathbb{A}^1) \rightarrow F(X \times (\mathbb{A}^1 - 0)) \right).$$

For $r > 1$ we define F_{-r} to be $(F_{1-r})_{-1}$. Sometimes we will write $F(X)_{-r}$ for $F_{-r}(X)$.

Since the inclusion $t = 1 : X \hookrightarrow X \times (\mathbb{A}^1 - 0) \subset X \times \mathbb{A}^1$ is split by the projection $X \times \mathbb{A}^1 \rightarrow X$, we have a canonical decomposition $F(X \times (\mathbb{A}^1 - 0)) \cong F(X) \oplus F_{-1}(X)$. Hence, F_{-1} is also homotopy invariant, and if F is a sheaf then so is F_{-1} . Here are some examples of this construction.

EXAMPLE 23.1. If $F = \mathcal{O}^*$ then $F_{-1} = \mathbb{Z}$, because $\mathcal{O}^*(X \times (\mathbb{A}^1 - 0)) = \mathcal{O}^*(X) \times \{t^n\}$ for every integral X . By 4.1, there is a quasi-isomorphism $\mathbb{Z}(1)_{-1} \simeq \mathbb{Z}[-1]$.

More generally, the higher Chow groups $CH^i(-, n)$ are homotopy invariant (see 17.4) and their contractions are given by the formula:

$$(23.1.1) \quad CH^i(X, n)_{-1} \cong CH^{i-1}(X, n-1).$$

This follows from the the Localization Theorem (see 17.4):

$$CH^{i-1}(X, n) \xrightarrow{(t=0)_*} CH^i(X \times \mathbb{A}^1, n) \rightarrow CH^i(X \times (\mathbb{A}^1 - 0), n) \rightarrow CH^{i-1}(X, n-1),$$

which is split as above by the pullback along $t = 1$ (using 19.13).

Theorem 19.1 allows us to rewrite the formula in (23.1.1) as:

$$H^{m,i}(X, \mathbb{Z})_{-1} = \mathbb{H}_{Zar}^m(X, \mathbb{Z}(i))_{-1} \cong \mathbb{H}_{Zar}^{m-1}(X, \mathbb{Z}(i-1)) = H^{m-1, i-1}(X, \mathbb{Z}).$$

This yields the formula $\mathbb{Z}(i)_{-1} \simeq \mathbb{Z}(i-1)[-1]$ in the derived category, and in **DM**.

EXAMPLE 23.2. We will see in the next lecture (in 24.1 and 24.8) that if F is a homotopy invariant Zariski sheaf with transfers, then $H^n(-, F)$ is homotopy invariant and $H_{Zar}^n(-, F)_{-1} \cong H_{Zar}^n(-, F_{-1})$.

EXAMPLE 23.3. Suppose that $1/n \in k$, and let M be a locally constant n -torsion sheaf, such as μ_n . The argument of 23.1 applied to étale cohomology,

shows that

$$H_{\acute{e}t}^m(X, M \otimes \mu_n)_{-1} \cong H_{\acute{e}t}^{m-1}(X, M).$$

EXERCISE 23.4. Let \mathcal{U} be the standard covering of $X \times (\mathbb{A}^n - 0)$ by $U_1 = X \times (\mathbb{A}^1 - 0) \times \mathbb{A}^{n-1}, \dots, U_n = X \times \mathbb{A}^{n-1} \times (\mathbb{A}^1 - 0)$. If F is homotopy invariant and $n \geq 2$, show that $\check{H}^0(\mathcal{U}, F) \cong F(X)$, $\check{H}^{n-1}(\mathcal{U}, F) \cong F_{-n}(X)$, and that $\check{H}^r(\mathcal{U}, F) = 0$ for all other r .

Now suppose that F is a Zariski sheaf, and that its cohomology groups are also homotopy invariant. Show that, for all m and $n > 0$, the cohomology with supports satisfies:

$$H_{X \times \{0\}}^m(X \times \mathbb{A}^n, F) \cong H^{m-n}(X, F)_{-n}.$$

Hint: Use the Čech spectral sequence $\check{H}^p(\mathcal{U}, H^q F) \Rightarrow H^{p+q}(X \times (\mathbb{A}^n - 0), F)$.

PROPOSITION 23.5. *Let F be a homotopy invariant presheaf with transfers. Then $(F_{Nis})_{-1} \cong (F_{-1})_{Nis}$.*

PROOF. By 13.1 and 22.3, F_{Nis} is a homotopy invariant sheaf with transfers. By inspection, the natural map $(F_{-1})_{Nis} \rightarrow (F_{Nis})_{-1}$ is a morphism of presheaves with transfers. By 11.2 (applied to the kernel and cokernel), it suffices to show that $F_{-1}(S) = (F_{Nis})_{-1}(S)$ when $S = \text{Spec } E$ for a field E . The left hand side is $F(\mathbb{A}_E^1 - 0)/F(\mathbb{A}_E^1)$ by definition, while the right side equals $F_{Nis}(\mathbb{A}_E^1 - 0)/F_{Nis}(\mathbb{A}_E^1)$. These are equal by 22.4 and 22.2. \square

In the rest of this lecture, we will compare F_{-1} to various sheaves $F_{(Y,Z)}$, which we now define.

DEFINITION 23.6. Given a closed embedding $i: Z \hookrightarrow Y$, and a presheaf F , we define a Nisnevich sheaf $F_{(Y,Z)}$ on Z as follows. Let $K = K_{(Y,Z)}$ denote the presheaf cokernel of $F \rightarrow j_* j^* F$, where $j: V \hookrightarrow Y$ is the complement of Z . That is, $K(U)$ is the cokernel of $F(U) \rightarrow F(U \times_Y V)$ for all U . We set $F_{(Y,Z)} = (i^* K)_{Nis}$.

Since sheafification is exact, there is a canonical exact sequence of sheaves

$$(23.6.1) \quad F_{Nis} \rightarrow (j_* j^* F)_{Nis} \rightarrow i_* F_{(Y,Z)} \rightarrow 0.$$

EXAMPLE 23.7. If $Z = \{z\}$ is a closed point on Y , then the value at Z of $F_{(Y,Z)}$ is the cohomology with supports, $H_Z^1(Y, F_{Nis})$. Indeed, if S is the Hensel local scheme of Y at Z then $F_{(Y,Z)}(Z)$ is the cokernel of $F_{Nis}(S) \rightarrow F_{Nis}(S - Z, F)$, i.e., $H_Z^1(S, F_{Nis})$. But this equals $H_Z^1(Y, F_{Nis})$ by excision [Har77, Ex.III.2.3]. Similarly, we have $H^n(-, F)_{(Y,Z)} \cong H_Z^{n+1}(Y, F)$ for $n > 0$. This follows from excision and the exact sequence

$$H^{n-1}(S, F) \rightarrow H^{n-1}(U, F) \rightarrow H_Z^n(S, F) \rightarrow 0.$$

EXAMPLE 23.8. Fix a Nisnevich sheaf F and consider the presheaf $H^n(-, F)$. We claim that if $n > 0$ then

$$H^n(-, F)_{(Y,Z)} = i^* R^n j_*(F).$$

Indeed, in (23.6.1) we have $H^n(-, F)_{Nis} = 0$, and $R^n j_*(F)$ is the sheaf on Y associated to the presheaf $j_* j^* H^n(-, F) = j_* H^n(-, F|_V)$. Hence $i_* H^n(-, F)_{(Y, Z)} \cong (j_* j^* H^n(-, F))_{Nis} = R^n j_*(F)$. Now apply i^* and observe that $i^* i_*$ is the identity.

EXAMPLE 23.9. Let $i : S \hookrightarrow S \times \mathbb{A}^1$ be the embedding $i(s) = (s, 0)$, with complement $S \times (\mathbb{A}^1 - 0)$. By definition, $F_{-1}(U) = K(U \times \mathbb{A}^1)$ where the cokernel presheaf K is defined in 23.6. The adjunction yields a natural map from $K(U \times \mathbb{A}^1)$ to $i_* i^* K(U \times \mathbb{A}^1) = i^* K(U)$. That is, we have a natural morphism of sheaves on S :

$$(F_{-1})_{Nis} \rightarrow F_{(S \times \mathbb{A}^1, S \times 0)}.$$

PROPOSITION 23.10. *Let F be a homotopy invariant presheaf with transfers. Then $(F_{-1})_{Nis}|_S \cong F_{(S \times \mathbb{A}^1, S \times 0)}$ for all smooth S .*

PROOF. We need to compare F_{-1} and $j_* j^* F / F$ in a sufficiently small neighborhood of any point s of any smooth affine S . We will use the standard triple $T = (\mathbb{P}_S^1 \rightarrow S, S \times \infty, S \times 0)$, which is split over $S \times \mathbb{A}^1$ by 11.12. For each affine neighborhood U of $S \times 0$ in $S \times \mathbb{A}^1$, set $T_U = (\mathbb{P}_S^1, \mathbb{P}_S^1 - U, S \times 0)$.

We claim that by shrinking S we can make T_U into a standard triple. At issue is whether or not $(\mathbb{P}_S^1 - U) \cup (S \times 0)$ lies in an affine open subscheme of \mathbb{P}_S^1 . Since the fiber U_s over s is open in \mathbb{P}_s^1 , there is an affine open $V \subset \mathbb{P}_k^1$ so that $s \times_k V$ contains both 0 and the finite set $\mathbb{P}_s^1 - U_s$. Hence the complements of U and $S \times V$ in \mathbb{P}_S^1 intersect in a closed subset, disjoint from the fiber \mathbb{P}_s^1 . Since \mathbb{P}_S^1 is proper over S , we may shrink S about s (keeping S affine) to assume that the complements are disjoint. Hence the affine $S \times V$ contains the complement $\mathbb{P}_S^1 - U$ as well as $S \times 0$, as claimed.

Now the identity on \mathbb{P}_S^1 is a finite morphism of standard triples $T_U \rightarrow T$ in the sense of 21.1 by 21.2. Setting $U_0 = U - (S \times 0)$, the square Q coming from this is:

$$\begin{array}{ccc} U_0 & \longrightarrow & U \\ \downarrow & & \downarrow \\ S \times (\mathbb{A}^1 - 0) & \xrightarrow{j} & S \times \mathbb{A}^1. \end{array}$$

By the standard triples theorem 21.6 applied to $Q' = Q$, the complex $F(MV(Q))$ is split exact:

$$0 \rightarrow F(S \times \mathbb{A}^1) \rightarrow F(S \times (\mathbb{A}^1 - 0)) \oplus F(U) \rightarrow F(U_0) \rightarrow 0.$$

Since F is homotopy invariant, this implies that $F(U) \rightarrow F(U_0)$ is injective and that $F_{-1}(S) \cong F(U_0)/F(U)$. Since $j : S \times (\mathbb{A}^1 - 0) \hookrightarrow S \times \mathbb{A}^1$ has $j_* j^* F(U) = F(U_0)$, the right side is $j_* j^* F / F(U)$. Passing to the limit over U and S , we get the statement. \square

LEMMA 23.11. *Let $f : Y \rightarrow X$ be an étale morphism and Z a closed subscheme of X such that $f^{-1}(Z) \rightarrow Z$ is an isomorphism. Then for every presheaf F :*

$$F_{(X, Z)} \xrightarrow{\cong} F_{(Y, f^{-1}(Z))}.$$

PROOF. Since this is to be an isomorphism of Nisnevich sheaves, we may assume that X is Hensel local, and that Z is not empty. Then Y is Hensel semilocal; the assumption that $f^{-1}(Z) \cong Z$ implies that Y is local and in fact $Y \cong X$. In this case the two sides are the same, namely $F(X - Z)/F(X) \cong F(Y - Z)/F(X)$. \square

Lemma 23.11 uses the Nisnevich topology in a critical way. For the Zariski topology, the corresponding result requires F to be a homotopy invariant presheaf with transfers, and may be proven along the same lines as 23.10; see [CohTh, 4.13].

THEOREM 23.12. *Let $i : Z \rightarrow X$ be a closed embedding of smooth schemes of codimension 1, and F a homotopy invariant presheaf with transfers. Then there exists a covering $X = \cup U_\alpha$ and isomorphisms on each $U_\alpha \cap Z$:*

$$F_{(U_\alpha, U_\alpha \cap Z)} \cong (F_{-1})_{\text{Nis}}.$$

That is, for each α there is an exact sequence of Nisnevich sheaves on U_α :

$$0 \rightarrow F_\alpha \rightarrow j_{\alpha*} j_\alpha^* F_\alpha \rightarrow i_*(F_{-1})_{\text{Nis}} \rightarrow 0.$$

Here $F_\alpha = (F|_{U_\alpha})_{\text{Nis}}$ and j_α denotes the inclusion $U_\alpha \cap (X - Z) \hookrightarrow U_\alpha$.

Moreover, for every smooth T we also have isomorphisms on $(U_\alpha \cap Z) \times T$:

$$F_{(U_\alpha \times T, (U_\alpha \cap Z) \times T)} \cong (F_{-1})_{\text{Nis}}.$$

PROOF. We have to show that every smooth pair (X, Z) of codimension one is locally like $(S \times \mathbb{A}^1, S \times 0)$. If $\dim(Z) = d$ then, by shrinking X about any point (and writing X instead of U), we may find an étale map $f : X \rightarrow \mathbb{A}^{d+1}$ such that $Z \cong f^{-1}(\mathbb{A}^d)$.

$$\begin{array}{ccc} Z & \xrightarrow{i} & X \\ \downarrow f & & \downarrow f \\ \mathbb{A}^d & \hookrightarrow & \mathbb{A}^{d+1} \cong \mathbb{A}^d \times \mathbb{A}^1 \end{array}$$

By construction, $Z \times \mathbb{A}^1$ is étale over $\mathbb{A}^d \times \mathbb{A}^1$. Form the pullback $X' = X \times_{\mathbb{A}^{d+1}} Z \times \mathbb{A}^1$ and note that both $X' \rightarrow X$ and $X' \rightarrow Z \times \mathbb{A}^1$ are étale with $Z' = Z \times_{\mathbb{A}^d} Z$ lying above Z and $Z \times 0$, respectively. Since $Z' \rightarrow Z$ is étale and has a canonical section Δ , we can write $Z' = \Delta(Z) \amalg W$. Setting $X'' = X - W$, both $X'' \rightarrow X$ and $X'' \rightarrow Z \times \mathbb{A}^1$ are étale, with $\Delta(Z)$ the inverse image of Z and $Z \times 0$, respectively. Applying lemma 23.11 twice and then 23.10, we obtain the required isomorphisms of Nisnevich sheaves on Z :

$$F_{(X, Z)} \xleftarrow{\cong} F_{(X'', \Delta(Z))} \xrightarrow{\cong} F_{(Z \times \mathbb{A}^1, Z \times 0)} \cong (F_{-1})_{\text{Nis}}.$$

To see that the sequence of sheaves is exact, we only need to observe that F_α injects into $j_* j^* F_\alpha$ by lemma 22.8, since $F_\alpha = (F|_{U_\alpha})_{\text{Zar}}$ by 22.2.

In order to prove the final assertion, it suffices to replace Z, X and \mathbb{A}^d with $Z \times T, X \times T$ and $\mathbb{A}^d \times T$ in the above argument. \square

PORISM 23.13. *The same proof shows that if $Z \rightarrow X$ is a closed embedding of smooth schemes of codimension r , then locally $F_{(X,Z)} \cong F_{(Z \times \mathbb{A}^r, Z \times 0)}$.*

EXAMPLE 23.14. Let M be a locally constant n -torsion étale sheaf and consider $F(X) = H^1(X, M \otimes \mu_n)$. By 23.3, $(F_{-1})_{\text{Nis}} \cong M$. By [Mil80, p. 243], we also have $F_{(X,Z)} \cong M$. In this case, the isomorphisms $F_{(X,Z)} \cong (F_{-1})_{\text{Nis}}$ of 23.12 hold for any cover of X .

Homotopy invariance of cohomology

We finally have all the tools to prove 13.8 which we restate here for the convenience of the reader.

THEOREM 24.1. *Let k be a perfect field and F a homotopy invariant presheaf with transfers. Then $H_{Nis}^n(-, F_{Nis})$ is a homotopy invariant presheaf (with transfers) for every n .*

PROOF. It suffices to prove that the $H_{Nis}^n(-, F_{Nis})$ are homotopy invariant, since we already know that they are presheaves with transfers from 13.4. We shall proceed by induction on n . The case $n = 0$ was completed in theorem 22.3, so we know that F_{Nis} is homotopy invariant. Hence, we may assume that $F = F_{Nis}$.

Consider $X \times \mathbb{A}^1 \xrightarrow{\pi} X$. Since $\pi_*F(U) = F(U \times \mathbb{A}^1) \cong F(U)$, we have $\pi_*F = F$. By induction we know that $R^q\pi_*F = 0$ for $0 < q < n$. By theorem 24.2 below, $R^n\pi_*F = 0$ as well. Hence the Leray spectral sequence

$$H_{Nis}^p(X, R^q\pi_*F) \Rightarrow H_{Nis}^{p+q}(X \times \mathbb{A}^1, F)$$

collapses enough to yield $H_{Nis}^n(X, F) \cong H_{Nis}^n(X \times \mathbb{A}^1, F)$. That is, the presheaf $H_{Nis}^n(-, F)$ is homotopy invariant. \square

We have thus reduced the proof of 24.1 to the following theorem. Recall from [EGA4, 17.5] that the Hensel local scheme $\text{Spec}(R)$ of a smooth variety at some point is **formally smooth**, i.e., geometrically regular.

THEOREM 24.2. *Let k be a perfect field, and F a homotopy invariant Nisnevich sheaf with transfers such that $R^q\pi_*F = 0$ for $0 < q < n$. If S is a formally smooth Hensel local scheme over k , then $H_{Nis}^n(S \times \mathbb{A}^1, F) = 0$.*

The requirement that k be perfect comes from the following fact (see [EGA0, 19.6.4]): if k is perfect, every regular local k -algebra is formally smooth over k .

PROOF. We will proceed by induction on $d = \dim(S)$. If $d = 0$ then $S = \text{Spec}(K)$ for some field K ; in this case, $H_{Nis}^n(S \times \mathbb{A}^1, F) = H_{Nis}^n(\mathbb{A}_K^1, F) = 0$ by 22.7. Here we have used exercise 2.10 to regard F as a homotopy invariant presheaf with transfers over K .

If $\dim(S) > 0$, and U is any proper open subscheme, then $\dim U < d$ (S is local), so $R^q\pi_*F|_U = 0$ for $0 < q \leq n$, by induction on d . Thus the canonical map $\pi|_U^* : H_{Nis}^n(U, F) \rightarrow H_{Nis}^n(U \times \mathbb{A}^1, F)$ is an isomorphism, and its inverse is induced

by the restriction $s|_U$ of the zero section $s : S \rightarrow S \times \mathbb{A}^1$ to U . From the diagram

$$\begin{array}{ccc} H_{Nis}^n(S \times \mathbb{A}^1, F) & \xrightarrow{j^*} & H_{Nis}^n(U \times \mathbb{A}^1, F) \\ s^* \downarrow & & s|_U^* \downarrow \cong \\ 0 = H_{Nis}^n(S, F) & \longrightarrow & H_{Nis}^n(U, F) \end{array}$$

we see that the top map j^* is zero for all such U .

Now $S = \text{Spec}(R)$ for a regular local ring (R, \mathfrak{m}) ; choose $r \in \mathfrak{m} - \mathfrak{m}^2$ and set $Z = \text{Spec}(R/r)$, $U = S - Z$. Because Z is regular and k is perfect, Z is formally smooth over k . For this choice, the map j^* is an injection by proposition 24.3 below. Hence the source $H_{Nis}^n(S \times \mathbb{A}^1, F)$ of j^* must be zero. \square

PROPOSITION 24.3. *Let k be a perfect field and S the Hensel local scheme of a smooth scheme X at some point. Let U be the complement of a smooth divisor Z on S . Under the inductive assumption that $R^q \pi_* F = 0$ for all $0 < q < n$, the following map is a monomorphism:*

$$H_{Nis}^n(S \times \mathbb{A}^1, F) \rightarrow H_{Nis}^n(U \times \mathbb{A}^1, F).$$

PROOF. Let i and j denote the inclusions of $Z \times \mathbb{A}^1$ and $U \times \mathbb{A}^1$ into $S \times \mathbb{A}^1$ respectively. Regarding F as a sheaf on $S \times \mathbb{A}^1$, the map in question factors as:

$$H_{Nis}^n(S \times \mathbb{A}^1, F) \xrightarrow{\tau} H_{Nis}^n(S \times \mathbb{A}^1, j_* j^* F) \xrightarrow{\eta} H_{Nis}^n(U \times \mathbb{A}^1, j^* F).$$

We first show that the right-hand map η is injective. This will follow from 24.4 below, once we have shown that $R^q j_* F = 0$ for $0 < q < n$. The inductive assumption implies that $H^q(F)$ is a homotopy invariant presheaf with transfers. Since $q > 0$ we have $H^q(F)_{Nis} = 0$. Now see from 23.5 that $(H^q(F)_{-1})_{Nis} \cong (H^q(F)_{Nis})_{-1} = 0$. By 23.8 and 23.12 (with $T = \mathbb{A}^1$) we have

$$R^q j_* F \cong i_* H^q(F)_{(S \times \mathbb{A}^1, Z \times \mathbb{A}^1)} \cong i_* (H^q(F)_{-1})_{Nis} = 0.$$

We now prove that the left-hand map τ is injective as well. Since F is a homotopy invariant presheaf with transfers, F injects into $j_* j^* F$ by lemma 22.8. By 23.6, there is a short exact sequence of Nisnevich sheaves on $S \times \mathbb{A}^1$:

$$0 \rightarrow F \rightarrow j_* j^* F \rightarrow i_* F_{(S \times \mathbb{A}^1, Z \times \mathbb{A}^1)} \rightarrow 0.$$

Since S is local, theorem 23.12 (with $T = \mathbb{A}^1$) implies that $F_{(S \times \mathbb{A}^1, Z \times \mathbb{A}^1)} \cong F_{-1}$ on $Z \times \mathbb{A}^1$. Consider the associated long exact sequence in cohomology.

$$\begin{array}{c} H^{n-1}(S \times \mathbb{A}^1, j_* j^* F) \rightarrow H^{n-1}(Z \times \mathbb{A}^1, F_{-1}) \\ \xrightarrow{\partial} H^n(S \times \mathbb{A}^1, F) \rightarrow H^n(S \times \mathbb{A}^1, j_* j^* F) \rightarrow H^n(Z \times \mathbb{A}^1, F_{-1}). \end{array}$$

It suffices to show that the map $H^{n-1}(S \times \mathbb{A}^1, j_* j^* F) \rightarrow H^{n-1}(Z \times \mathbb{A}^1, F_{-1})$ is onto. If $n > 1$, this follows from the homotopy invariance of F_{-1} and the fact that Z is Hensel local:

$$H^{n-1}(Z \times \mathbb{A}^1, F_{-1}) \cong H^{n-1}(Z, F_{-1}) = 0.$$

If $n = 1$, we argue as follows. Since F and F_{-1} are homotopy invariant, the two left horizontal maps are isomorphisms in the commutative diagram:

$$\begin{array}{ccccc}
 F(U) & \xrightarrow{\cong} & F(U \times \mathbb{A}^1) & \xleftarrow{=} & H^0(S \times \mathbb{A}^1, j_* j^* F) \\
 \downarrow \text{23.12 onto} & & \downarrow & & \downarrow \\
 F_{-1}(Z) & \xrightarrow{\cong} & F_{-1}(Z \times \mathbb{A}^1) & \xleftarrow{=} & H^0(Z \times \mathbb{A}^1, F_{-1}).
 \end{array}$$

The left vertical map is onto by 23.12, because S is local. It follows that the right vertical map is onto, as desired. \square

LEMMA 24.4. *Let G be any sheaf on $U \times \mathbb{A}^1$ such that $R^q j_* G = 0$ for $0 < q < n$. Then the canonical map $H^n(X \times \mathbb{A}^1, j_* G) \rightarrow H^n(U \times \mathbb{A}^1, G)$ is an injection.*

PROOF. Consider the Leray spectral sequence

$$H^p(X \times \mathbb{A}^1, R^q j_* G) \implies H^{p+q}(U \times \mathbb{A}^1, G).$$

Using the assumption on the vanishing of the $R^q j_* G$, it is easy to see that there is a short exact sequence:

$$0 \rightarrow H^n(X \times \mathbb{A}^1, j_* G) \rightarrow H^n(U \times \mathbb{A}^1, G) \rightarrow H^0(X \times \mathbb{A}^1, R^n j_* G). \quad \square$$

We have now completed the proof of homotopy invariance of the cohomology sheaves, which was promised in lecture 13 (as theorem 13.8).

For the rest of this lecture, we fix a homotopy invariant Zariski sheaf with transfers F over a perfect field k . Because we have proven theorem 13.8, we may use proposition 13.9, which says that $H_{Zar}^*(X, F) \cong H_{Nis}^*(X, F)$. We will sometimes suppress the subscript and just write $H^*(X, F)$.

COROLLARY 24.5. *If S is a smooth semilocal scheme over k and F is a homotopy invariant sheaf with transfers, then for all $n > 0$:*

- $H^n(S, F) = 0$;
- $H^n(S \times T, F) = 0$ for every open subset T of \mathbb{A}_k^1 .

PROOF. (Cf. 13.9.) By 24.1, each $H^n(-, F)$ is a homotopy invariant presheaf with transfers. If E is the field of fractions of S , then $H^n(\text{Spec } E, F) = 0$ for $n > 0$ because $\dim E = 0$. By 11.1, this implies that $H^n(S, F) = 0$.

Now $H^n(X) = H^n(X \times T, F)$ is also a homotopy invariant presheaf with transfers by 24.1, and $H^n(S)$ injects into $H^n(\text{Spec } E) = H^n(\text{Spec}(E) \times T, F)$ by 11.1. By 2.10 and 22.7, this group vanishes for $n > 0$. \square

EXAMPLE 24.6. Let (R, \mathfrak{m}) be a discrete valuation ring containing k , with field of fractions E and residue field $K = R/\mathfrak{m}$. Setting $S = \text{Spec } R$ and $Z = \text{Spec } K$, theorem 23.12 yields $F_{(S,Z)} \cong F_{-1}$ and an exact sequence of Nisnevich sheaves on S , $0 \rightarrow F \rightarrow j_* F \rightarrow i_* F_{-1} \rightarrow 0$. Since $H_{Nis}^1(S, F) = 0$ by 24.5, the Nisnevich cohomology sequence yields the exact sequence:

$$0 \rightarrow F(\text{Spec } R) \rightarrow F(\text{Spec } E) \rightarrow F_{-1}(\text{Spec } K) \rightarrow 0.$$

More generally, if R is a semilocal principal ideal domain with maximal ideals \mathfrak{m}_i , the same argument (using 24.5) yields an exact sequence:

$$0 \rightarrow F(\text{Spec } R) \rightarrow F(\text{Spec } E) \rightarrow \bigoplus_i F_{-1}(\text{Spec } R/\mathfrak{m}_i) \rightarrow 0.$$

EXERCISE 24.7. If X is a smooth curve over k , show that $F_{-1}(x) \cong H_x^1(X, F)$ for every closed point $x \in X$. Conclude that there is an exact sequence

$$0 \rightarrow F(X) \rightarrow F(\text{Spec } k(X)) \rightarrow \bigoplus_{x \in X} F_{-1}(x) \rightarrow H_{\text{Zar}}^1(X, F) \rightarrow 0.$$

PROPOSITION 24.8. *Let k be a perfect field and F a homotopy invariant Zariski sheaf with transfers. Then $H^n(-, F)_{-1} \cong H^n(-, F_{-1})$ for all smooth X . That is, there is a natural isomorphism:*

$$H_{\text{Zar}}^n(X \times (\mathbb{A}^1 - 0), F) \cong H_{\text{Zar}}^n(X, F) \oplus H_{\text{Zar}}^n(X, F_{-1}).$$

PROOF. Write T for $\mathbb{A}^1 - 0$ and consider the projection $\pi : X \times T \rightarrow X$. Let S be the local scheme at a point x of X . The stalk of $R^q \pi_* F$ at x is $H^q(S \times T, F)$, which vanishes for $q > 0$ by 24.5. Therefore the Leray spectral sequence degenerates to yield $H^n(X \times T, F) \cong H^n(X, \pi_* F)$. But $\pi_* F \cong F \oplus F_{-1}$ by the definition of F_{-1} . \square

EXAMPLE 24.9. Let F be a homotopy invariant Zariski sheaf with transfers. Combining proposition 24.8 with 24.1 and 23.4, we get the formula:

$$H_{Z \times \{0\}}^n(Z \times \mathbb{A}^r, F) \cong H^{n-r}(Z, F_{-r}).$$

If $Z = \text{Spec}(K)$ for a field K , this shows that $H_{\{0\}}^n(\mathbb{A}_K^r, F)$ vanishes for $n \neq r$, while the value of $H_{\{0\}}^r(\mathbb{A}_K^r, F)$ at $\text{Spec}(K)$ is $F_{-r}(\text{Spec}(K))$.

LEMMA 24.10. *Let S be a d -dimensional regular local scheme over a perfect field k . If F is a homotopy invariant sheaf with transfers and Z is the closed point of S , then $H_Z^n(S, F)$ vanishes for $n \neq d$, while $H_Z^d(S, F) \cong F_{-d}(Z)$.*

PROOF. Since the case $d = 0$ is trivial, and $d = 1$ is given in example 24.6, we may assume that $d > 1$. Write U for $S - Z$. Since $F(S)$ injects into $F(U)$ by 11.1, $H_Z^0(S, F) = 0$. For $n > 0$, we may use $H^{n-1}(-, F)$, which is a homotopy invariant presheaf with transfers by 24.1. By 23.11 and two applications of 23.7, we have

$$H_Z^n(S, F) \cong H^{n-1}(-, F)_{(S, Z)} \cong H^{n-1}(-, F)_{(Z \times \mathbb{A}^d, Z \times 0)} \cong H_{Z \times 0}^n(Z \times \mathbb{A}^d, F).$$

By 24.9, this group vanishes for $n \neq d$, and equals $F_{-d}(Z)$ if $n = d$. \square

If z is a point of X with closure Z , and A is an abelian group, let $(i_z)_*(A)$ denote the constant sheaf A on Z , extended to a sheaf on X .

THEOREM 24.11. *Let X be smooth over k , and F a homotopy invariant Zariski sheaf with transfers. Then there is a canonical exact sequence of Zariski sheaves on X :*

$$0 \rightarrow F \rightarrow \coprod_{\text{codim } z=0} (i_z)_*(F) \rightarrow \coprod_{\text{codim } z=1} (i_z)_*(F_{-1}) \rightarrow \cdots \rightarrow \coprod_{\text{codim } z=r} (i_z)_*(F_{-r}) \rightarrow \cdots$$

PROOF. It suffices to assume that X is local with generic point x_0 and closed point x_d , and construct the exact sequence

$$0 \rightarrow F(S) \rightarrow F(x_0) \rightarrow \coprod_{\text{codim } z=1} (F_{-1}(z)) \rightarrow \cdots \rightarrow \coprod_{\text{codim } z=r} (F_{-d}(z)) \rightarrow \cdots \rightarrow F(x_d) \rightarrow 0.$$

When $\dim(X) = 1$ this is 24.6, so we may assume that $d = \dim(X) > 1$. For any $r \leq d$, let $H^n(X^r, F)$ denote the direct limit of the groups $H^n(X - T, F)$ with $\text{codim}(T) > r$. For any Zariski sheaf F , and $r > 0$, the direct limit (over T and all Z of codimension r) of the long exact cohomology sequences $H_Z^*(X - T, F) \rightarrow H^*(X - T, F) \rightarrow H^*(X - Z - T, F)$ yields an exact sequence

$$0 \rightarrow \coprod_{\substack{\text{codim } z \\ =r}} H_z^0(X_z, F) \rightarrow F(X^r) \rightarrow F(X^{r-1}) \rightarrow \coprod_{\substack{\text{codim } z \\ =r}} H_z^1(X_z, F) \rightarrow H^1(X^r, F) \rightarrow \cdots.$$

Each X_z is an r -dimensional local scheme. Hence the groups $H_z^n(X_z, F)$ vanish except for $n = r$ by 24.10, and $H_z^r(X_z, F) \cong F_{-r}(z)$. For $r > 0$ this yields:

$$F(X) \cong F(X^{d-1}) \cong \cdots \cong F(X^r) \cong \cdots \cong F(X^1);$$

$$0 = H^r(X, F) \cong H^r(X^{d-1}, F) \cong \cdots \cong H^r(X^{r+1}, F);$$

and (since X^0 is a point)

$$0 = H^r(X^0, F) \cong H^r(X^1, F) \cong \cdots \cong H^r(X^{r-1}, F).$$

Using these, we get exact sequences:

$$0 \rightarrow F(X) \rightarrow F(x_0) \rightarrow \coprod_{\text{codim } z=1} H_z^1(X_z, F) \rightarrow H^1(X^1, F) \rightarrow 0;$$

and (for $0 < r \leq d$)

$$0 \rightarrow H^{r-1}(X^{r-1}, F) \rightarrow \coprod_{\text{codim } z=r} H_z^r(X_z, F) \rightarrow H^r(X^r, F) \rightarrow 0.$$

Splicing these together (and using 24.10) yields the required exact sequence. \square

REMARK 24.12. Since the sheaves $(i_z)_*(F_{-r})$ are flasque, theorem 24.11 gives a flasque resolution of the sheaf F . Taking global sections yields a chain complex which computes the cohomology groups $H^n(X, F)$. This shows that the coniveau spectral sequence

$$E_1^{p,q} = \bigoplus_{\text{codim } x=p} H_z^{p+q}(X, F) \implies H^{p+q}(X, F)$$

degenerates, with $E_2^{p,0} = H^p(X, F)$ and $E_2^{p,q} = 0$ for $q \neq 0$.

Bibliography

- [ABS87] R. MacPherson A. Beilinson and V. Schechtman, *Notes on motivic cohomology*, Duke Math. J. **54** (1987), 679–710.
- [Ada74] J. F. Adams, *Stable homotopy and generalised homology*, University of Chicago Press, Chicago, Ill., 1974, Chicago Lectures in Mathematics. MR 53 #6534
- [BG73] K. S. Brown and S. M. Gersten, *Algebraic K-theory as generalized sheaf cohomology*, Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Springer, Berlin, 1973, pp. 266–292. Lecture Notes in Math., Vol. 341.
- [BL94] S. Bloch and S. Lichtenbaum, *A spectral sequence for motivic cohomology*, Preprint, 1994.
- [Blo86] S. Bloch, *Algebraic cycles and higher K-theory*, Adv. in Math. **61** (1986), no. 3, 267–304.
- [Blo94] ———, *The moving lemma for higher Chow groups*, J. Algebraic Geom. **3** (1994), no. 3, 537–568.
- [BT73] H. Bass and J. Tate, *The Milnor ring of a global field*, Algebraic K-theory, II: “Classical” algebraic K-theory and connections with arithmetic (Proc. Conf., Seattle, Wash., Battelle Memorial Inst., 1972) (Berlin), Springer, 1973, pp. 349–446. Lecture Notes in Math., Vol. 342.
- [EGA1sv] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique I*, Springer-Verlag, New York, 1971.
- [EGA0] ———, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas.*, Inst. Hautes Études Sci. Publ. Math. (1964), no. 20, Reprinted in [EGA1sv].
- [EGA1] ———, *Éléments de géométrie algébrique. I. Le langage des schémas*, Inst. Hautes Études Sci. Publ. Math. (1960), no. 4, 228, Reprinted in [EGA1sv].
- [EGA3] ———, *Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents.*, Inst. Hautes Études Sci. Publ. Math. (1961,1963), no. 11,17.
- [EGA4] ———, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas.*, Inst. Hautes Études Sci. Publ. Math. (1964-1967), no. 20,24,28,32.
- [DMOS82] P. Deligne, J. Milne, A. Ogus, and K.-Y. Shih, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics, vol. 900, Springer-Verlag, Berlin, 1982, Philosophical Studies Series in Philosophy, 20.
- [FS02] E. M. Friedlander and A. Suslin, *The spectral sequence relating algebraic K-theory to motivic cohomology*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 6, 773–875.
- [Ful75] W. Fulton, *Rational equivalence on singular varieties*, Inst. Hautes Études Sci. Publ. Math. (1975), no. 45, 147–167.
- [Ful84] ———, *Intersection theory*, Springer-Verlag, Berlin, 1984.
- [GL00] T. Geisser and M. Levine, *The K-theory of fields in characteristic p*, Invent. Math. **139** (2000), no. 3, 459–493.
- [GM88] S. Gel'fand and Yu. Manin, *Metody gomologicheskoi algebrы. Tom I*, “Nauka”, Moscow, 1988, Vvedenie v teoriyu kogomologii i proizvodnye kategorii.
- [GM03] ———, *Methods of homological algebra*, second ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003, Translation of [GM88].

- [Gro57] A. Grothendieck, *Sur quelques points d'algèbre homologique*, Tôhoku Math. J. (2) **9** (1957), 119–221.
- [Har77] R. Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.
- [Jou73] J.-P. Jouanolou, *Une suite exacte de Mayer-Vietoris en K -théorie algébrique*, Algebraic K -theory, I: Higher K -theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Springer, Berlin, 1973, pp. 293–316. Lecture Notes in Math., Vol. 341.
- [KS86] K. Kato and S. Saito, *Global class field theory of arithmetic schemes*, Applications of algebraic K -theory to algebraic geometry and number theory, Part I, II (Boulder, Colo., 1983), Amer. Math. Soc., Providence, RI, 1986, pp. 255–331.
- [Lev94] M. Levine, *Bloch's higher Chow groups revisited*, K -theory (Strasbourg, 1992), 1994, pp. 235–320.
- [Lev98] ———, *Mixed motives*, American Mathematical Society, Providence, RI, 1998.
- [Lic84] S. Lichtenbaum, *Values of zeta-functions at nonnegative integers*, Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Springer, Berlin, 1984, pp. 127–138.
- [Lic94] ———, *Motivic complexes*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 303–313.
- [Mac71] S. Mac Lane, *Categories for the working mathematician*, Springer-Verlag, New York, 1971.
- [Man68] Ju. Manin, *Correspondences, motifs and monoidal transformations*, Mat. Sb. (N.S.) **77** (119) (1968), 475–507.
- [May01] J. P. May, *The additivity of traces in triangulated categories*, Adv. Math. **163** (2001), no. 1, 34–73.
- [Mil70] J. Milnor, *Algebraic K -theory and quadratic forms*, Invent. Math. **9** (1969/1970), 318–344.
- [Mil80] J. Milne, *Étale cohomology*, Princeton University Press, Princeton, N.J., 1980.
- [Nis89] Ye. A. Nisnevich, *The completely decomposed topology on schemes and associated descent spectral sequences in algebraic K -theory*, Algebraic K -theory: connections with geometry and topology (Lake Louise, AB, 1987), Kluwer Acad. Publ., Dordrecht, 1989, pp. 241–342.
- [NS89] Yu. Nesterenko and A. Suslin, *Homology of the general linear group over a local ring, and Milnor's K -theory*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 1, 121–146.
- [Rec71] D. Rector, *K -theory of a space with coefficients in a (discrete) ring*, Bull. Amer. Math. Soc. **77** (1971), 571–575.
- [RG71] M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de "platification" d'un module*, Invent. Math. **13** (1971), 1–89.
- [Ric89] J. Rickard, *Derived categories and stable equivalence*, J. Pure Appl. Algebra **61** (1989), no. 3, 303–317.
- [Ser65] J.-P. Serre, *Algèbre locale. Multiplicités*, Springer-Verlag, Berlin, 1965.
- [SGA1] *Revêtements étales et groupe fondamental*, Springer-Verlag, Berlin, 1971, Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de M. Raynaud, Lecture Notes in Mathematics, Vol. 224.
- [SGA4] *Théorie des topos et cohomologie étale des schémas*, Springer-Verlag, Berlin, 1972–1973, Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J.-L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 269, 270, 305.
- [Sus82] A. Suslin, *Mennicke symbols and their applications in the K -theory of fields*, Proceedings of a Conference held at Oberwolfach, June 16–20, 1980 (R. K. Dennis, ed.), Springer-Verlag, Berlin, 1982, pp. 334–356.
- [Sus83] ———, *On the K -theory of algebraically closed fields*, Invent. Math. **73** (1983), no. 2, 241–245.

- [SV96] A. Suslin and V. Voevodsky, *Singular homology of abstract algebraic varieties*, Invent. Math. **123** (1996), no. 1, 61–94.
- [SV00] ———, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, The Arithmetic and Geometry of Algebraic Cycles (B. Gordon, J. Lewis, S. Müller-Stach, S. Saito, N. Yui, ed.), Nato ASI series C, vol. 548, Kluwer, 2000, pp. 117–189.
- [Swa62] R. G. Swan, *Vector bundles and projective modules*, Trans. Amer. Math. Soc. **105** (1962), 264–277.
- [Swa72] ———, *Some relations between higher K-functors*, J. Algebra **21** (1972), 113–136.
- [Ver96] J.-L. Verdier, *Des catégories dérivées des catégories abéliennes*, Astérisque (1996), no. 239, xii+253 pp. (1997), With a preface by Luc Illusie, Edited and with a note by Georges Maltsiniotis.
- [Voe02] V. Voevodsky, *Cancellation theorem*, Preprint. <http://www.math.uiuc.edu/K-theory/541>, 2002.
- [VSF00] V. Voevodsky, A. Suslin and E. M. Friedlander, *Cycles, transfers, and motivic homology theories*, Annals of Mathematics Studies, vol. 143, Princeton University Press, 2000.
- [Wei89] C. Weibel, *Homotopy algebraic K-theory*, Algebraic K-theory and algebraic number theory (Honolulu, HI, 1987), Amer. Math. Soc., Providence, RI, 1989, pp. 461–488.
- [Wei94] ———, *An introduction to homological algebra*, Cambridge University Press, Cambridge, 1994.
- [Whi78] G. Whitehead, *Elements of homotopy theory*, Springer-Verlag, New York, 1978.

References in [VSF00].

- [RelCh] A. Suslin and V. Voevodsky, *Relative Cycles and Chow Sheaves*, in [VSF00], pp. 10–86.
- [CohTh] V. Voevodsky, *Cohomological Theory of Presheaves with Transfers*, in [VSF00], pp. 87–137.
- [BivCy] E. M. Friedlander and V. Voevodsky, *Bivariant Cycle Cohomology*, in [VSF00], pp. 138–187.
- [TriCa] V. Voevodsky, *Triangulated Categories of Motives Over a Field*, in [VSF00], pp. 188–238.
- [HighCh] A. Suslin, *Higher Chow Groups and Étale Cohomology*, in [VSF00], pp. 239–254.

Glossary

- $\otimes^{\mathbb{L}}$ total tensor product, 56
- \otimes^{tr} tensor product of presheaves with transfers, 57
- \otimes_L^{tr} tensor product of complexes of presheaves with transfers, 57
- $\otimes_{\acute{e}t}^{tr}$ tensor product of étale sheaves with transfers, 58
- $\otimes_{L, Nis}^{tr}$ tensor product on $\mathbf{D}^- Sh_{Nis}(Cor_k, R)$, 109
- $\otimes_{L, \acute{e}t}^{tr}$ tensor product of complexes of étale sheaves with transfers, 58
- $\otimes_{\mathcal{L}}$ tensor product on \mathcal{L} , 73
- $a_{\acute{e}t}(F)$ étale sheafification of F , 42
- $a_{Nis}(F)$ Nisnevich sheafification of F , 90
- \mathcal{A}^{\oplus} the closure of \mathcal{A} under infinite direct sums, 55
- $A(q)$ the complex of presheaves with transfers $\mathbb{Z}(q) \otimes A$, 21
- $A_{r,i}(Y, X)$ bivariant cycle cohomology group, 130
- $c(X/S, 0)$ universally integral relative cycles of X finite and surjective over S , 10
- $C_0(X/S)$ same as $c(X/S, 0)$, 47
- C_*F the complex obtained from the simplicial presheaf $F(- \times \Delta^\bullet)$, 17
- $C_*^{DK}F$ normalized complex associated to C_*F , 17
- $\mathcal{C}[T^{-1}]$ category obtained from \mathcal{C} by inverting $\otimes T$, 65
- $CH^i(X)$ Chow group of codimension i cycles of X , 14
- $\mathbf{Ch}^-(\mathcal{A})$ category of bounded above cochain complexes in \mathcal{A} , 56
- $CH^i(X, m)$ Bloch's higher Chow group, 135
- $Chow$ ($Chow^{\text{eff}}$) the category of (effective) Chow motives, 167
- Cor_k category of finite correspondences, 4
- $Cor(X, Y)$ group of finite correspondences from X to Y , 3
- Cor_S category of finite correspondences over a Noetherian scheme S , 7
- $Cor_S(X, Y)$ the group $c(X \times_S Y/X, 0)$, 11
- $Cycl(X/S, r)$ free abelian group of the relative cycles W on X over S such that each component has dimension r over S , 9
- \mathbf{D}^- or $\mathbf{D}_{\acute{e}t}^-$ or $\mathbf{D}^-(Sh_{\acute{e}t}(Cor_k, R))$ derived category of étale sheaves of R -modules with transfers, 67
- \mathbf{D}^- or $\mathbf{D}^-(Sh_{Nis}(Cor_k, R))$ derived category of Nisnevich sheaves with transfers, 109
- \mathbf{D}_{Nis}^- or $\mathbf{D}_{Nis}^-(Sh_{Nis}(Sm/k))$ derived category of Nisnevich sheaves, 92
- Δ^\bullet cosimplicial scheme with $\Delta^n \cong \mathbb{A}^n$, 16
- $\mathbf{D}^-(G, \mathbb{Z}/m)$ derived category of discrete \mathbb{Z}/n -modules over G , 74
- $\mathbf{DM}^-(k, R)$ category of motives, 110

- $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}$ or $\mathbf{DM}_{\acute{e}t}^{\text{eff},-}(k, R)$ category of effective étale motives, 67
 $\mathbf{DM}_{\acute{e}t}^-$ or $\mathbf{DM}_{\acute{e}t}^-(k, R)$ category of étale motives, 68
 $\mathbf{DM}_{\text{gm}}^{\text{eff}}(k, R)$ category of effective geometric motives, 109
 $\mathbf{DM}_{\text{gm}}(k, R)$ category of geometric motives, 110
 $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}$ or $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}(k, R)$ category of effective motives, 109
 $\mathcal{E}_{\mathbb{A}^1}$ the thick subcategory of \mathbf{D}^- such that $\mathbf{DM}^- = \mathbf{D}^- / \mathcal{E}_{\mathbb{A}^1}$, 67
 Et/k category of smooth schemes over k of dimension zero, 38
 F_{-r} contraction of the presheaf F , 191
 F_{cdh} cdh sheafification of F , 95
 $F_{\acute{e}t}$ same as $a_{\acute{e}t}(F)$, 42
 F_{Nis} same as $a_{\text{Nis}}(F)$, 90
 $F_{(Y,Z)}$ Nisnevich sheaf on Z associated to a closed embedding of Z into Y , 192
 Γ_f the graph of f , 3
 \mathbb{G}_m the pointed scheme $(\mathbb{A}^1 - 0, 1)$, 15
 $\mathbb{G}_{\bar{X}, Y}$ units of \bar{X} equal to 1 on Y , 49
 $H^{p,q}(X, A)$ motivic cohomology group, 22
 $H_{n,i}^{BM}(X, R)$ (Borel-Moore) motivic homology with compact supports, 130
 $H_c^{n,i}(X, R)$ motivic cohomology with compact supports, 130
 $H_L^{p,q}(X, A)$ étale (or Lichtenbaum) motivic cohomology, 75
 $H_0^{\text{sing}}(X/k)$ the group $H_0 C_* \mathbb{Z}_{\text{tr}}(X)(\text{Spec } k)$, 18
 $H_*^{\text{sing}}(X/S)$ algebraic singular homology of X over S , 47
 $H_*^{\text{sing}}(X, R)$ algebraic singular homology, 78
 h_X additive functor $\text{Hom}_{\mathcal{A}}(-, X)$, 55
 $\mathbb{H}_{\text{cdh}}^n(X, L)$ cdh hypercohomology of a complex of cdh sheaves, 115
 $\mathbb{H}_{\acute{e}t}^i(X, K)$ étale hypercohomology of a complex of sheaves, 45
 $\mathbb{H}_{\text{Nis}}^n(X, K)$ Nisnevich hypercohomology of a complex of sheaves, 100
 $\mathbb{H}_{\text{Zar}}^p(X, L)$ Zariski hypercohomology of a complex of sheaves, 22
 $\underline{\text{Hom}}(M, N)$ internal Hom in \mathbf{DM}_{gm} , 170
 $\underline{\text{Hom}}(F, G)$ Hom presheaf, 56
 $\mathbf{K}^-(\mathcal{A})$ chain homotopy category of complexes in \mathcal{A} , 57
 $K_*^M(k)$ the Milnor K -theory of a field k , 29
 K_n K -theory group, 14
 \mathbb{L} the Lefschetz motive $\mathbb{Z}(1)[2]$, 167
 \mathcal{L} or $\mathcal{L}_{\acute{e}t}$ \mathbb{A}^1 -local objects in $\mathbf{D}^-(\text{Sh}_{\acute{e}t}(\text{Cor}_k, R))$, 71
 \mathcal{L} of \mathcal{L}_{Nis} \mathbb{A}^1 -local objects in $\mathbf{D}^-(\text{Sh}_{\text{Nis}}(\text{Cor}_k, R))$, 111
 \mathcal{L}_{Δ} line bundle on $U \times_S \bar{X}$ corresponding to the diagonal map, 85
 (\mathcal{L}, t) line bundle \mathcal{L} with a trivialization t , 49
 $\mathcal{M}^*(\mathbb{P}^1; 0, \infty)$ sheaf sending X to the rational functions on $X \times \mathbb{P}^1$ which are regular in a neighborhood of $X \times \{0, \infty\}$ and equal to 1 on $X \times \{0, \infty\}$, 25
 M^* the dual of a motive M , 168
 $M(q)$ Tate twist $M \otimes_{L_{\text{Nis}}}^{\text{tr}} R(q)$, 110
 $M^c(X)$ motive with compact support of X , 128

- μ_n sheaf of n^{th} roots of unity, 14
 $MV(Q)$ Mayer-Vietoris sequence of a square Q , 177
 $M(X)$ the motive of X in $\mathbf{DM}_{Nis}^{\text{eff},-}$, \mathbf{DM}^- or \mathbf{DM}_{gm} , 109
 \mathcal{O} sheaf of global functions, 13
 \mathcal{O}^* sheaf of global units, 13
 $\mathcal{O}_{X,x}^h$ Hensel local ring of X at x , 44
 $\text{Pic}(X)$ the Picard group of X , 25
 $\text{Pic}(\bar{X}, Y)$ relative Picard group, 49
 $\text{PreSh}(\text{Cor}_k)$ category of additive presheaves with transfers, 13
 $\mathbf{PST}(k)$ same as $\text{PreSh}(\text{Cor}_k)$, 13
 $Q(X, Y, A)$ cartesian square of schemes, 90
 $\underline{RHom}(M, L)$ internal Hom in $\mathbf{DM}_{Nis}^{\text{eff},-}$, 113
 Sch/k category of schemes of finite type over k , 94
 $\text{Sh}_{\acute{e}t}(\text{Cor}_k)$ category of étale sheaves with transfers, 37
 $\text{Sh}_{\acute{e}t}(\text{Sm}/k)$ category of étale sheaves on smooth schemes, 37
 $\text{Sh}_{\acute{e}t}^{\text{lc}}$ category of locally constant sheaves in $\text{Sh}_{\acute{e}t}(\text{Sm}/k)$, 38
 $\text{Sh}_{Nis}(\text{Cor}_k)$ category of Nisnevich sheaves with transfers, 99
 $\text{Sh}_{Nis}(\text{Sm}/k)$ category of Nisnevich sheaves on smooth schemes, 90
 S_l Henselization at $\{0\}$ in \mathbb{A}^l , 52
 Sm/k category of smooth separated schemes, 3
 $W_{\mathbb{A}}$ multiplicative system of \mathbb{A}^1 -weak equivalences, 67
 (\bar{X}, X_{∞}, Z) standard triple, 84
 $Z_1 \cdot Z_2$ intersection product of cycles, 143
 $\mathbb{Z}(\mathcal{A})$ category of additive presheaves on \mathcal{A} , 55
 $z_{\text{equi}}(T, r)$ sheaf of equidimensional cycles of relative dimension r , 125
 $z_{\text{equi}}^i(X, m)$ same as $z_{\text{equi}}(X, \dim X - i)(\Delta^m)$, 149
 $z^i(X, m)$ Bloch's cycle group, 135
 $z^i(Y, m)_{\not\mathcal{W}}$ cycles in $z^i(Y, m)$ meeting \mathcal{W} properly, 137
 $\mathbb{Z}(q)$ the motivic complex $C_*\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge q})[-q]$, 21
 $\mathbb{Z}^{SF}(i)$ Suslin-Friedlander motivic complex, 126
 $\mathbb{Z}_{tr}(\mathbb{G}_m)$ the presheaf with transfers $\mathbb{Z}_{tr}(\mathbb{A}^1 - 0)/\mathbb{Z}$, 15
 $\mathbb{Z}_{tr}(\mathbb{G}_m^{\wedge q})$ smash product, 16
 $\mathbb{Z}_{tr}(\check{U})$ Čech complex associated to a cover \check{U} of X , 39
 $\mathbb{Z}_{tr}(X)$ representable presheaf with transfers associated to X , 15
 $\mathbb{Z}_{tr}(X, x)$ cokernel of the map $x_* : \mathbb{Z} \rightarrow \mathbb{Z}_{tr}(X)$, 15
 $\mathbb{Z}(X)$ Nisnevich sheafification of $\mathbb{Z}[\text{Hom}_{\text{Sm}/k}(-, X)]$, 92

Index

Page numbers in **boldface** refer to definitions.

- \mathbb{A}^1 -homotopy, **19**, 19–20
 - \mathbb{A}^1 -homotopic, **19**, 19–20, 47, 69, **69**, 71, 85, 86, 113, **113**, 121–123, 141, 142, 177–179
 - \mathbb{A}^1 -homotopy equivalence, **20**, 20, 69, 120, 178
 - strict, **72**, 72–74, 100
- \mathbb{A}^1 -local complexes, *see* Nisnevich \mathbb{A}^1 -local
- \mathbb{A}^1 -local objects, *see* étale \mathbb{A}^1 -local, Nisnevich \mathbb{A}^1 -local
- \mathbb{A}^1 -weak equivalences, *see* étale \mathbb{A}^1 -weak equivalences, Nisnevich \mathbb{A}^1 -weak equivalences
- abelian category, 13, 15, 39, 42–44, 55, 90, 99
- abstract blow-up, *see* blow-up
- additive category, 3–5, 19, 37, 55–57, 63, 64, 67
- adjoint functors, 6, 20, 38, 42, 51, 56, 61, 70, 71, 99, 112, 113, 169, 193
- algebraic singular homology $H_i^{sing}(X/S)$, **48**, 47–54, **78**, 78–80, 179
 - and Chow groups, 136
 - and higher Chow groups, 136
 - and motivic homology, 114
 - and transfers, 48
- balanced functor, 56
- Bass-Tate lemma, 33
- Bertini’s Theorem, 87
- bivariant cycle cohomology group, 130
- Bloch’s cycle complex $z^i(X, *)$, **135**, 135–142, 149–154
 - and equidimensional cycle complex, 149
 - subcomplex $z^i(X, *)_{\neq}$, **137**, 136–139, 163–165
 - and equidimensional cycles, 160
 - is isomorphic when X affine, 137
- Bloch’s cycle presheaf $z^i(- \times T, *)$, **160**, 160–163
- Bloch, Spencer, 135, 162, 163
- blow-up
 - abstract, **94**, 94–97, 106, 109, 111, 115
 - decomposition of motives, 123
 - sequence in cohomology, 104
 - sequence in motivic cohomology, 115
 - triangle, 106, 111
- Borel-Moore motivic homology, *see* motivic homology with compact supports
- Brauer group, 44
- Brown-Gersten, 162
- calculus of fractions, 67, 71
- Cancellation Theorem, 110, 131, 135, 164, 167–169
- canonical flasque resolution, *see* flasque resolution
- Cartier divisor, 50, 86, 180
- cdh cohomology, 97
- cdh hypercohomology, 130
 - and hyperext, 115
 - and motivic cohomology, 115
- cdh resolution, 106
- cdh sheaf, 129
- cdh sheaves with transfers, 105–106
- cdh topology, 45, **95**, 94–97, 115
- Čech cohomology, 183, 184, 192
- Čech resolution, 39, 41, 59, 99, 103, 119, 121
- Chevalley’s theorem, 84
- Chow group, 14, 135, 136, 167
 - and algebraic singular homology, 136
 - and motivic cohomology, 159
 - and $z_{equi}(T, r)$, 126
 - higher, *see* higher Chow group
 - is a presheaf with transfers, 14
- Chow motives, 167–169

- cohomological dimension
 - cdh, 95, 115
 - étale, 45, 71–73, 117
 - Nisnevich, 89, 100, 101, 112
 - Zariski, 101
- cohomology, *see* motivic cohomology, étale cohomology, Nisnevich cohomology, cdh cohomology
- compact support, *see* motive with compact support
- compactification, *see* good compactification
- contraction, **191**, 198
 - of higher Chow groups, 191
 - of homotopy invariant Zariski sheaf, 200
 - of locally constant étale sheaf, 191
 - of motivic cohomology, 191
 - of motivic complex, 191
 - of relative sheaf, 193, 194
 - of Zariski cohomology, 191
- corrections, 69, 186
- correspondence
 - \mathbb{A}^1 -homotopic, *see* \mathbb{A}^1 -homotopy composition, **4**
 - elementary, **3**, 3–6, 121
 - finite, **3**, 3–6
 - lifts to a vector bundle torsor, 139
- covering morphism, *see* standard triple
- cycles
 - equidimensional, *see* equidimensional cycles
 - in good position, 138, 139, **144**, 143–147
 - intersection, 4, 135, 143
 - properly intersecting, 5, 135–138, **143**, 143–147, 150
 - pullback, *see* pullback
 - push-forward, *see* push-forward
- deformation retract, **93**
- Deligne, Pierre, 28
- direct limit, 6, 15, 83
- Dold-Kan correspondence, 17
- dual motive, *see* geometric motive
- dual vector space, 78
- Duality Theorem, 131, 164, 167, 168
- effective étale motives, *see* étale motives
- effective geometric motives, *see* geometric motives
- effective motives, **109**, 109–118
 - and Chow motives, 167
 - and Nisnevich \mathbb{A}^1 -local, 113
 - list of properties, 110
 - \mathbb{Q} coefficients, 116–118
 - are effective étale motives, 118
- Eilenberg-Zilber theorem, 24, 76
- elementary correspondence, *see* correspondences
- elementary matrices, 122
- enough injectives, 13, 43, 44, 79, 99
- enough projectives, 13, 56
- equidimensional cycle complex $z_{equi}^i(X, *)$, **149**, 149–154
 - and Bloch's cycle complex, 149
- equidimensional cycle presheaf $z_{equi}(T, r)$, **125**, 125–131, 135, 136, 159
 - and Chow group, 126
 - and motives with compact support, 128
 - is a presheaf with transfers, 126
 - is an étale sheaf, 125
 - is $\mathbb{Z}_{rr}(T)$ if T proper, 125
- equidimensional cycles, 9, 11, 143, 145
 - locally, 137
- equidimensional scheme, 135, 149–150, 164
- étale \mathbb{A}^1 -local, **71**, 71–80, 117–118
 - and étale effective motives, 74
 - is Nisnevich \mathbb{A}^1 -local rationally, 117
- étale \mathbb{A}^1 -weak equivalence, **67**, 67–80
- étale cohomology, 27, 45
 - and Ext, 45
 - preserves transfers, 44
- étale hypercohomology
 - and hyperext, 45
- étale motives, **68**
 - and étale effective motives, 68
 - effective, **67**, 67–80
- étale motivic cohomology, 67, **75**, 75–80, 117, 118
 - and étale cohomology, 75
 - is motivic cohomology rationally, 117
 - is representable, 78, 118
- étale resolution, 61, 103
- étale sheaf, **37**
 - locally constant, **38**, 38–39, 51, 58, 60, 61, 72–74, 79, 191, 195
 - with transfers, **37**, 37–45
- étale sheafification
 - preserves transfers, 42
- étale topology, 40, 160
 - and Tate twist, *see* Tate twist
- Ext group, 78
- fat point, **8**
- ffp cohomology, 28
- field
 - extension, *see* field extension

- perfect, 100–103, 105, 112, 113, 117–119, 124, 126, 128, 131, 159, 167, 183, 197–200
- separable closure, 39, 51
- separably closed, 51, 52, 78–80
- field extension, 6–9, 23, 26, 30–33, 87, 88, 102, 186
 - Galois, 6, 10, 39, 60
 - inseparable, 10, 33
 - inseparable degree, 39
 - normal, 39
 - purely inseparable, 10
 - separable, 6, 10, 15, 20, 23, 33, 38, 89
 - separable degree, 39
- finite correspondence, *see* correspondences
- flasque resolution, 61, 162, 201
 - canonical, 44, 45, 92, 99, 100
- flat presheaf, *see* tensor product
- flat site, 160
- formally smooth, 197
- Friedlander, Eric, 130, 131
- Friedlander-Voevodsky Duality, *see* Duality Theorem
- Gabber, Offer, 84
- Galois group, 6, 10, 38, 39, 67, 74
- Galois module, 38, 39, 58, 60, 67
- Generic Equidimensionality Theorem, 151, 153, 155
- geometric motives, **110**, 167–171
 - are rigid, 170
 - double dual isomorphism, 169
 - dual, **168**
 - dual and motive with compact supports, 169
 - dual and tensor product, 170
 - effective, **109**, 167–171
- geometric point, 51
- geometrically regular, 197
- Gersten resolution, 135, 200
- global functions
 - is a presheaf with transfers, 13
 - is zero in motives, 71
- global units, 184, 191
 - is a presheaf with transfers, 13
 - is $\mathbb{Z}(1)[1]$, 25
- good compactification, 47–54, 84, 85, 176
- Gysin
 - map, 123
 - map for compact supports, 130
 - triangle, 111, 124
- h-topology, 95
- Hensel local ring or scheme, 40, 51, **90**, 101, 102, 117, 128, 192, 194, 197, 198
 - strictly, 40, 44
- Henselization, 52
- higher Chow groups, 125, 130, 131, **135**, 135–142, 149, 150, 159–165, 191
 - and algebraic singular homology, 136
 - and Chow groups, 136
 - and vector bundle torsors, 139
 - are functors on affine schemes, 139
 - are $H_{i,n}^{BM}$, 159, 164
 - are hypercohomology, 162
 - are motivic cohomology, 159, 163
 - are presheaves with transfers, 142
 - are representable, 165
 - definition of transfer, 140
 - Homotopy Invariance, 136, 139, 149, 161, 165, 191
 - Localization Theorem, 136, 139, 150, 161, 191
- Hom presheaf, 56, 61
- homogeneous form, 156
- homotopic
 - \mathbb{A}^1 –, *see* \mathbb{A}^1 -homotopy
 - weekly, *see* weakly homotopic
- homotopy commutative, **177**, 177–181
- homotopy invariance, **17**, 17–20, 83–88
 - higher Chow groups, *see* higher Chow groups
 - Nisnevich \mathbb{A}^1 -local and, 112
 - Nisnevich cohomology preserves, 100, 197
 - Nisnevich sheaf, 100
 - is the Zariski sheaf, 183
 - Nisnevich sheaf and contractions, 192
 - Nisnevich sheafification preserves, 183
 - presheaf $H_n C_* F$, 18
 - Zariski sheaf, 199
 - Zariski sheafification and, 183, 187
- homotopy invariant presheaf with transfers, *see* homotopy invariance
- hyperext, *see* cohomology
- injective resolution, 79
- internal Hom, 113, 168
 - of geometric motives, 170
- intersection cycle, *see* cycles
- intersection multiplicity, 143, 144
 - and the Tor formula, 143
- Jouanolou’s device, 139, 140
- K -theory, 14, 48
 - Milnor, *see* Milnor K -theory

- Kummer Theory, 50, 51
- leading form, **156**
- Lefschetz motive, 167
- Levine, Marc, 137, 138, 163
- Lichtenbaum motivic cohomology, *see* étale motivic cohomology
- line bundle, 14, 50, 85–87, 123, 176, 178, 179
trivialization, 49, 50, 84–86, 123, 176, 178–181
- localization, 64, 65, 67, 68, 109, 110
- localization sequence, 127, 136
- Localization Theorem, *see* higher Chow groups
- locally constant, *see* étale sheaf
- Mayer-Vietoris property, **162**
- Mayer-Vietoris sequence $MV(Q)$, **177**, 183–186, 189, 193
- Mayer-Vietoris triangle, 103, 110, 123, 129
- Milnor K -theory, 29–34
is $H^{n,n}$, 29
- motive $M(X)$, *see also* effective motives, geometric motives, motives, 20, **109**
Chow, *see* Chow motive
decomposition of projective space, 121
dual, *see* geometric motive
geometric, *see* geometric motive
is homotopy invariant, 20
isomorphic to $C_*\mathbb{Z}_{tr}(X)$, 110
list of properties, 110
projective bundle, 111, 123
vector bundle, 111
- motive with compact support $M^c(X)$, **128**, 128–131, 168–169
Duality Theorem, 131
embedding triangle, 129
is geometric, 129
is $M(X)$ for X proper, 129
- motives $\mathbf{DM}^-(k, R)$, 110, 168
- motivic cohomology, **22**
and cdh hypercohomology, 115
and Chow group, 159
and Nisnevich hypercohomology, 101
colimits, 23
independent of k , 23
is a presheaf with transfers, 14
is étale motivic cohomology rationally, 117
is higher Chow groups, 159
is representable, 101, 110, 114
pairing, 24, 122
product is graded-commutative, 122
singular X , **114**
with compact supports, **130**
- motivic complex, **21**, 21–24
and the projective space, 119
is $\mathbb{Z}^{SF}(i)$, 126
product, 24, 76, 122
- motivic homology, **114**
with compact supports, **130**, 159, 164
is higher Chow groups, 164
- multiplicative system, 64, 67, 109
- Nisnevich \mathbb{A}^1 -local, **111**, 111–118
and effective motives, 113
and homotopy invariance, 112
is étale \mathbb{A}^1 -local rationally, 117
- Nisnevich cohomology
and cdh cohomology, 97, 106
and Ext, 92, 99
and Zariski cohomology, 101
is étale cohomology rationally, 116
preserves homotopy invariance, 100, 197
preserves transfers, 99
- Nisnevich cover, 96
- Nisnevich hypercohomology
and hyperext, 100
and Zariski hypercohomology, 101
is étale hypercohomology rationally, 117
- Nisnevich lifting property, 89, 95
- Nisnevich resolution, 40, 92, 94, 99, 106
- Nisnevich sheaf, 40, 89–94, 96, 102
criterion, 90
not étale sheaf, 91
with transfers, 99–106
is étale after $\otimes\mathbb{Q}$, 116
 $\mathbb{Z}(X)$, **92**, 92–94, 103–105, 115
- Nisnevich sheafification
preserves transfers, 99
- Nisnevich topology, 40–42, 44, 45, 68, 83, **89**, 194
and Zariski topology, 102
points are Hensel local, 90
- nodal curve, 97
- norm map, 13, 30–34, 53, 181
Milnor K -theory, 30–34
- normal bundle, 123
- normal surface, 97
- perfect field, *see* field
- Picard group, 25–27, 44, 49, 50, 91, 122, 135
relative, 47, **49**, 49–54, 83–86, 178–181
- platification, 7, 96, 106, 129
- Pontrjagin dual, 61, 80

- presheaf with transfers, **13**, 13–20
 homotopy invariant, *see* homotopy invariance
 $\mathbb{Z}_{tr}(X)$, *see* $\mathbb{Z}_{tr}(X)$
- Projection Formula, 137, 145
- projective bundle, 95, 122, 123
- Projective Bundle Theorem, 123
- projective objects, 15, 16, 55–58, 79
- projective resolution, 55–58, 60, 61
- proper birational cover, **95**
- proper cdh cover, **95**, 94–97
- pseudo pretheory, 48
- pullback of cycles, 5, 126, 131, 137, 145–147, 150, 160, 191
 along a fat point, **8**, 8–11
 along a finite correspondence, **144**
 and relative cycles, 125, 145
 flat, 7, 8, 14, 37, 128, 137, 139, 144, 160, 163, 164
 is defined, **143**
- pure subgroup, 83
- push-forward of cycles, 4, 5, 11, 14, 30, 33, 53, 54, 126, 136, 144, 147
- relative cycle, **9**, 8–11, 125, 145
 universally integral, **7**, **10**, 10–11
- relative Picard group, *see* Picard group
- relative sheaf, 192, 198
- representable presheaf, 55
- resolution of singularities, 45, 78, 97, 105, 106, 109, 111, 115, 116, 127, 129–131, 135, 149, 159, 164, 167
- rigid tensor category, 168–171
- Rigidity Theorem, 51, 73, 79
- roots of unity, 14, 25, 27, 28, 38, 49–51, 58, 61, 72, 75, 76, 78, 99, 192, 195
 vs. $\mathbb{Z}/l(1)$, 27, 99
 $\mu_n^{\otimes q} \cong \mathbb{Z}/n(q)$, 77
- semilocal ring or scheme, 3, 40, 83, 86, 185–188, 199, 200
- separable closure of a field, *see* field
- separably closed field, *see* field
- separated presheaf, 185
- sheaf, *see* Zariski sheaf, étale sheaf, Nisnevich sheaf, cdh sheaf
- simplicial decomposition, 18, 152
- simplicial group, 16, 24, 135, 136, 149, 159, 160
- simplicial homotopy, 18, 153
- simplicial map, 24, 76
- simplicial presheaf, 16, 17, 76, 126, 159, 160
- singular schemes, *see* $\mathbb{Z}_{tr}(X)$, cdh topology
- skeletal homotopy, 152
- skeletal map, **150**, 150–154
- smash product \wedge , 16, 29, 31, 64, 76–78, 119–122
 and tensor product, 58
- smooth curve, 85, 176, 184, 200
- spectral sequence, 60, 73, 79, 94, 102, 103
 Čech, 192
 coniveau, 201
 hypercohomology, 23, 117
 hyperext, 104, 112
 hyperhomology, 59
 Leray, 101, 117, 197, 199, 200
- stable homotopy category, 64
- standard triple, **84**, 84–88, 189, 193
 covering morphism, **175**, 175–185
 split, **85**, 85–88, 176, 178, 179, 182
- strong deformation retract, *see* deformation retract
- Suslin, Andrei, 51, 78, 149, 155, 157
- Suslin-Friedlander motivic complex $\mathbb{Z}^{SF}(i)$, **126**, 126–129, 159–163
 is independent of k , 126
 is isomorphic to Bloch’s cycles sheaf, 161
 is $\mathbb{Z}(i)$, 126
- suspension, 64
- symmetric group, 121–122
- symmetric monoidal category, 6, 55–57, 63–66, 170
- Tate twist, **110**
 étale is invertible, 61, 68
- tensor product
 and smash product, 58
 in Cor_k , 6
 in derived category, 60
 of \mathbb{A}^1 -local objects, 113
 of flat presheaves, 56, 92
 of motives with compact support, 129
 of Nisnevich sheaves, 92
 of presheaves, 56–61, 92
 of presheaves with transfers, 57
 of sheaves, 58–61
 total, *see* total tensor product
- tensor triangulated category, 57, 60, **63**, 63–66, 68, 74, 109, 113
 $\mathcal{E}[T^{-1}]$ is a, 65
 of motives, 110
- thick, **67**, 68, 71, 74, 109, 113
- topology, *see* Zariski topology, étale topology, Nisnevich topology, cdh topology
- total Hom, 79

- trace map, 13, 38
- truncation
 - brutal, 68
 - good, 102, 105
- unibranch scheme
 - geometrically, 9
- universally integral cycles, *see* relative cycles
- upper distinguished square, **90**, 91, 175, 177, 181, 182, 184, 189
 - coming from a covering morphism of standard triples, **175**, 176, 177, 181
- Vanishing Theorems, 22, 159
- vector bundle, 103, 111, 122, 123
 - torsor, 139–142, 160, 164
- Voevodsky, Vladimir, 95, 110, 130, 131

- Walker, Mark, 87
- weakly homotopic, **153**
- Weil divisor, 25, 26, 50, 53, 178, 198
- Weil Reciprocity, 30, 34

- Yoneda lemma, 15, 55, 57, 69, 92, 100, 105, 127, 168

- Zariski covering, 183
- Zariski descent, **161**, 162, 163
- Zariski resolution, 41
- Zariski sheaf, 103, 128
 - with transfers, 116, 183–189, 191, 199–201
- Zariski topology, 40, 42, 44, 83, 100, 187, 194
- Zariski's Main Theorem, 49
- $\mathbb{Z}_{tr}(X)$
 - and flatness, 56
 - is a sheaf, 21, 37
 - is projective, 15
 - singular X , 15, 37, 114, 115

