

# The $(\Pi, \lambda)$ -structures on the C-systems defined by universe categories<sup>1</sup>

Vladimir Voevodsky<sup>2</sup>

## Abstract

We introduce the notion of a  $(\Pi, \lambda)$ -structure on a C-system and construct a bijection, for a given C-system, between the sets of  $(\Pi, \lambda)$ -structures and structures of products of families of types introduced previously by Cartmell and Streicher.

We then define the notion of a  $P$ -structure on a universe in a locally cartesian closed category and construct a  $(\Pi, \lambda)$ -structure on the C-systems  $CC(\mathcal{C}, p)$  from a  $P$ -structure on  $p$ .

In the last section we define homomorphisms of C-systems with  $(\Pi, \lambda)$ -structures and functors of universe categories with  $P$ -structures and show that the construction of the previous section is functorial relative to these definitions.

## Contents

|          |   |           |
|----------|---|-----------|
| <b>1</b> | <b>Introduction</b>   | <b>1</b>  |
| <b>2</b> | <b>Presheaves <math>\mathcal{O}b_i</math> and <math>\widetilde{\mathcal{O}b}_i</math> on the C-systems defined by universe categories</b> | <b>4</b>  |
| 2.1      | Functor $Sig$ and functor isomorphisms $S\mathcal{O}b_i$ and $S\widetilde{\mathcal{O}b}_i$ . . . . .                                      | 4         |
| 2.2      | The functor $D_p$ . . . . .   | 9         |
| 2.3      | Isomorphisms of presheaves $u_1$ and $\tilde{u}_1$ . . . . .  | 12        |
| 2.4      | Functor isomorphisms $SD_p$ . . . . .   | 18        |
| 2.5      | Isomorphisms of presheaves $u_n$ and $\tilde{u}_n$ for $n \geq 2$ . . . . .   | 20        |
| 2.6      | The case of a locally cartesian closed $\mathcal{C}$ . . . . .  | 22        |
| <b>3</b> | <b><math>P</math>-structures on universes and <math>(\Pi, \lambda)</math>-structures</b>  | <b>27</b> |
| 3.1      | Construction of $(\Pi, \lambda)$ -structures on the C-systems $CC(\mathcal{C}, p)$ . . . . .  | 27        |
| <b>4</b> | <b>Functoriality</b>  | <b>30</b> |
| 4.1      | Universe category functors and the $D_p$ construction . . . . .   | 30        |
| 4.2      | Universe category functors and isomorphisms $u_n$ and $\tilde{u}_n$ . . . . .   | 38        |
| 4.3      | Universe category functors and the $I_p$ construction . . . . .   | 38        |

---

<sup>1</sup>2000 *Mathematical Subject Classification*: 03F50, 18C50 03B15, 18D15,

<sup>2</sup>School of Mathematics, Institute for Advanced Study, Princeton NJ, USA. e-mail: vladimir@ias.edu

|          |   |           |
|----------|---|-----------|
| 4.4      | More on universe category functors . . . . .  | 40        |
| 4.5      | Functoriality properties of the $(\Pi, \lambda)$ -structures constructed from $P$ -structures . . . . . | 44        |
| <b>5</b> | <b>Appendices</b>   | <b>46</b> |
| 5.1      | Appendix A. Categories with binary products and binary cartesian closed categories . . . . .            | 46        |
| 5.2      | Appendix B. Slice categories, pullbacks and locally cartesian closed categories . . . . .               | 51        |
| <b>6</b> | <b>Acknowledgements</b>   | <b>56</b> |

# 1 Introduction

The concept of a C-system in its present form was introduced in [?]. The type of the C-systems is constructively equivalent to the type of contextual categories defined by Cartmell in [?] and [?] but the definition of a C-system is slightly different from the Cartmell’s foundational definition.

In this paper we consider what might be the most important structure on C-systems - the structure that corresponds, for the syntactic C-systems, to the operations of dependent product,  $\lambda$ -abstraction and application. A C-system formulation of this structure was introduced by John Cartmell in [?, pp. 3.37 and 3.41] as a part of what he called a strong M.L. structure. It was studied further by Thomas Streicher in [?, p.71] who called a C-system (contextual category) together with such a structure a “contextual category with products of families of types”.

The constructions and proofs of the main part of the paper require knowing many facts about C-systems. These facts are established in Section ???. Many of these facts are new, some have been stated by Cartmell [?] and Streicher [?], but without proper mathematical proofs. Among notable new facts we can mention Lemma ??? that shows that the canonical direct product in a C-system is strictly associative.

In Section ??? we construct on any C-system presheaves  $\mathcal{O}b_n$  and  $\widetilde{\mathcal{O}b}_n$ . These presheaves play a major role in our approach to the C-system formulation of systems of operations that correspond to systems of inference rules. The main result here is Construction ??? for Problem ???. It is likely that constructions for various other variants of this problem involving morphisms between presheaves  $\mathcal{O}b_*$  and  $\widetilde{\mathcal{O}b}_*$  can be given. The full generality of this result should involve as the source fiber products of  $\mathcal{O}b_*$  and  $\widetilde{\mathcal{O}b}_*$  relative to morphisms satisfying certain properties and as the target  $\mathcal{O}b_*$  or  $\widetilde{\mathcal{O}b}_*$ . We limit ourselves to Construction ??? here because it is the only case that will be required later in the paper.

In Section 2.3 we first remind the definition of the product of families of types structure on a C-system. Then, in Definition???, we give the first of the two main definitions of this paper, the definition of a  $(\Pi, \lambda)$ -structure. In the rest of this section we work on constructing a bijection between the sets of structures of products of families of types and  $(\Pi, \lambda)$ -structures

on a given C-system. This is probably the most technical part of the paper which is not surprising considering how different Definitions ?? and ?? are.

This construction uses most of the results of Section ??.

The  $(\Pi, \lambda)$ -structures correspond to the  $(\Pi, \lambda, app, \beta, \eta)$ -system of inference rules. In Remark ?? we outline the definitions of classes of structures that correspond to the similar systems but without the  $\beta$ - or  $\eta$ -rules. Such structures appear as natural variations of the  $(\Pi, \lambda)$ -structures.

In Section 3 we consider the case of C-systems of the form  $CC(\mathcal{C}, p)$  introduced in [?]. They are defined, in a functorial way, by a category  $\mathcal{C}$  with a final object and a morphism  $p : \tilde{U} \rightarrow U$  together with the choice of pullbacks of  $p$  along all morphisms in  $\mathcal{C}$ . A morphism with such choices is called a universe in  $\mathcal{C}$ . As a corollary of general functoriality we also obtain a construction of an isomorphism that connects the C-systems  $CC(\mathcal{C}, p)$  corresponding to different choices of pullbacks and different choices of final objects. It makes it possible to say that  $CC(\mathcal{C}, p)$  is defined by  $\mathcal{C}$  and  $p$ .

We provide several intermediate results about  $CC(\mathcal{C}, p)$  when  $\mathcal{C}$  is a locally cartesian closed category leading to the main result of this paper - Construction 3.4 that produces a  $(\Pi, \lambda)$ -structure on  $CC(\mathcal{C}, p)$  from a simple pullback<sup>3</sup> based on  $p$ . This construction was first announced in [?]. It and the ideas that it is based on are among the most important ingredients of the construction of the univalent model of the Martin-Lof type theory.

In the following sections we study the behavior of our construction with respect to universe category functors and prove that it is functorial with respect to functors equipped with an additional structure that reflects compatibility with the choice of the generating pullback.

One may wonder how the construction of this paper relates to the earlier ideas of Seely [?] and their refinement by Clairambault and Dybjer [?]. This question requires further study.

The methods of this paper are fully constructive.

The paper is written in the formalization-ready style that is in such a way that no long arguments are hidden even when they are required only to substantiate an assertion that may feel obvious to readers who are closely associated with a particular tradition of mathematical thought.

As a result, a number of lemmas, especially in the appendices, may be well know to many readers. Their proofs are nevertheless included to comply with the requirements of the formalization ready style.

On the other hand, not all preliminary lemmas are included or a reference to a complete proof is given. There are some, but very much fewer than is usual in today's papers, exceptions.

The main result of this paper is not a theorem but a construction and so are many of the intermediate results. Because of the importance of constructions for this paper we use a special pair of names Problem-Construction for the specification of the goal of a construction and the description of the particular solution.

---

<sup>3</sup>We say “a pullback” instead of “a pullback square”.

In the case of a Theorem-Proof pair one usually refers (by name or number) to the theorem when using the proof of this theorem. This is acceptable in the case of theorems because the future use of their proofs is such that only the fact that there is a proof but not the particulars of the proof matter.

In the case of a Problem-Construction pair the content of the construction often matters in the future use. Because of this we have to refer to the construction and not to the problem and we assign in this paper numbers both to Problems and to Constructions.

We use below the concept of a universe. In the Zermelo-Fraenkel set theory, the main intended formalization base for this paper, a universe is simply a set  $U$  that is usually assumed to satisfy some properties such as, for example, that it is closed under formation of pairs - if two sets  $A$  and  $B$  are elements of  $U$  then the set representing the pair  $(A, B)$  is an element of  $U$ . We do not provide a precise set of such conditions that we assume. To assume the universes mentioned in the paper to be Grothendieck universes would certainly suffice but in most cases we need a much weaker set of conditions. It is likely that the conditions that we need are weak enough to be able to prove the existence of such universes inside the “canonical” Zermelo-Fraenkel theory without any large cardinal axioms.

In this paper we continue to use the diagrammatic order of writing composition of morphisms, i.e., for  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  the composition of  $f$  and  $g$  is denoted by  $f \circ g$ .

We denote by  $\Phi^\circ$  the functor  $PreShv(C') \rightarrow PreShv(C)$  given by the pre-composition with a functor  $\Phi^{op} : C^{op} \rightarrow (C')^{op}$ , that is,

$$\Phi^\circ(F)(X) = F(\Phi(X))$$

In the literature this functor is denoted both by  $\Phi^*$  and  $\Phi_*$  and we decided to use a new unambiguous notation instead.

Acknowledgements are at the end of the paper.

While abbreviated notations may be helpful for getting a general impression from a brief scroll through the paper, long notations become indispensable when one seeks true understanding.

In view of Lemma 3.5, Construction ?? can be used not only to construct the product of families of types structures on C-systems, but also to prove that such structures do not exist. This applies also to structures corresponding to other systems of inference rules in type theory. For example, a similar technique may be used not only to construct a model of a particular kind of higher inductive types, but also to show that for a given universe  $p$  no such model on  $CC(\mathcal{C}, p)$  exists.

## 2 Presheaves $\mathcal{O}b_i$ and $\widetilde{\mathcal{O}b}_i$ on the C-systems defined by universe categories

### 2.1 Functor $Sig$ and functor isomorphisms $S\mathcal{O}b_i$ and $S\widetilde{\mathcal{O}b}_i$

In this section we consider three constructions that apply to any C-system  $CC$ . The functor  $Sig : PreShv(CC) \rightarrow PreShv(CC)$  and two families of isomorphisms parameterized by  $i \in \mathbf{N}$ :

$$S\mathcal{O}b_i : Sig(\mathcal{O}b_i) \rightarrow \mathcal{O}b_{i+1}$$

and

$$S\widetilde{\mathcal{O}b}_i : Sig(\widetilde{\mathcal{O}b}_i) \rightarrow \widetilde{\mathcal{O}b}_{i+1}$$

Let  $\mathcal{G}$  be a presheaf on  $CC$ . For  $\Gamma \in CC$  we set

$$[2016.08.30.eq7] Sig(\mathcal{G})(\Gamma) = \coprod_{T \in \mathcal{O}b_1(\Gamma)} \mathcal{G}(T) \quad (2.1)$$

and for  $f : \Gamma' \rightarrow \Gamma$

$$[2016.08.30.eq8] Sig(\mathcal{G})(f)(T, g) = (f^*(T), \mathcal{G}(q(f, T))(T)) \quad (2.2)$$

**Lemma 2.1** [2016.08.28.11] *The presheaf data  $Sig$  is a presheaf, that is, one has:*

1. for  $\Gamma \in CC$ ,

$$Sig(\mathcal{G})(Id_\Gamma) = Id_{Sig(\mathcal{G})(\Gamma)}$$

2. for  $f' : \Gamma'' \rightarrow \Gamma'$ ,  $f : \Gamma' \rightarrow \Gamma$ ,

$$Sig(\mathcal{G})(f' \circ f) = Sig(\mathcal{G})(f) \circ Sig(\mathcal{G})(f')$$

**Proof:** For the identity we have

$$Sig(\mathcal{G})(Id_\Gamma)(T, g) = (Id_\Gamma^*(T), \mathcal{G}(q(Id_\Gamma, T))(g)) = (T, g)$$

where the second equality is by axioms of the C-system structure. For the composition we have

$$\begin{aligned} Sig(\mathcal{G})(f')(Sig(\mathcal{G})(f)(T, g)) &= Sig(\mathcal{G})(f')(f^*(T), \mathcal{G}(q(f, T))(g)) = \\ ((f')^*(f^*(T)), \mathcal{G}(q(f', f^*(T)))(\mathcal{G}(q(f, T))(g))) &= ((f')^*(f^*(T)), \mathcal{G}(q(f', f^*(T)) \circ q(f, T))(g)) = \\ ((f' \circ f)^*(T), \mathcal{G}(q(f' \circ f, T))(g)) &= Sig(\mathcal{G})(f' \circ f)(T, g) \end{aligned}$$

where the first two equalities are by definition of  $Sig(\mathcal{G})$ , the third by the composition property of  $\mathcal{G}$ , the fourth by the axioms of the C-system structure and the fifth again by the definition of  $Sig(\mathcal{G})$ . This completes the proof of Lemma 2.1.

One defines  $Sig$  on morphisms of presheaves  $r : \mathcal{G} \rightarrow \mathcal{G}'$  by the family of morphisms

$$[\mathbf{2016.08.30.eq9}] Sig(r)_\Gamma(T, g) = (T, r_T(g)) \quad (2.3)$$

For  $r : \mathcal{G} \rightarrow \mathcal{G}'$  and  $f : \Gamma' \rightarrow \Gamma$ , we have

$$Sig(\mathcal{G})(f) \circ Sig(r)_{\Gamma'} = Sig(r)_\Gamma \circ Sig(\mathcal{G}')(f)$$

that is, the family of functions  $Sig(r)_\Gamma$  parametrized by  $\Gamma \in CC$  is a morphism of presheaves.

For  $\mathcal{G} \in PreShv(CC)$  we have

$$[\mathbf{2016.12.14.eq1}] Sig(Id_{\mathcal{G}})_\Gamma(T, g) = (T, (Id_{\mathcal{G}})_T(g)) = (T, g) \quad (2.4)$$

and for  $r : \mathcal{G} \rightarrow \mathcal{G}'$ ,  $r' : \mathcal{G}' \rightarrow \mathcal{G}''$  we have

$$[\mathbf{2016.12.14.eq2}] Sig(r \circ r')_\Gamma(T, g) = (T, (r \circ r')_T(g)) = (T, r'_T(r_T(g))) = Sig(r')_\Gamma(Sig(r)_\Gamma(T, g)) \quad (2.5)$$

These two equalities show that the functor data given by  $Sig$  on presheaves and  $Sig$  on morphisms of presheaves is a functor that we also denote by

$$Sig : PreShv(CC) \rightarrow PreShv(CC)$$

**Remark 2.2**  $[\mathbf{2016.12.14.rem1}]$  The construction of  $Sig$  works in more general setting than presheaves.

Indeed, for any family of sets  $G(\Gamma)$  parametrized by  $\Gamma \in CC$  the formula (2.1) defines a new family of sets  $Sig(G)(\Gamma)$  also parametrized by  $\Gamma \in CC$ . For any two families  $G, G'$  and a family of functions  $r_\Gamma : G(\Gamma) \rightarrow G'(\Gamma)$  the formula (2.3) defines a family of functions  $Sig(r)_\Gamma : Sig(G)(\Gamma) \rightarrow Sig(G')(\Gamma)$ . The properties (2.4) and (2.5) hold in this more general setting.

We can also define  $Sig(G)$  for any presheaf data, that is, for any pair consisting of a family  $G(\Gamma)$  of sets parametrized by  $\Gamma \in CC$  and a family of functions  $G(f) : G(\Gamma) \rightarrow G(\Gamma')$  parametrized by  $f : \Gamma' \rightarrow \Gamma$  in  $Mor(CC)$ . For this we can again use formulas (2.1) and (2.2).

If  $r_\Gamma : G(\Gamma) \rightarrow G'(\Gamma)$  is a morphism of functor data, that is functions  $r_*$  commute with functions  $G(*)$ , then  $Sig(r)$  is a morphism of functor data as well.

The presheaves  $\mathcal{O}b_n$  on  $CC$  were defined in  $[\text{?}, \text{Sec. 3}]$ . On objects they are given by

$$[\mathbf{2016.11.15.eq5}] \mathcal{O}b_n(\Gamma) = \{T \in Ob(CC) \mid l(T) = l(\Gamma) + n, ft^n(T) = \Gamma\} \quad (2.6)$$

and on morphisms  $f : \Gamma' \rightarrow \Gamma$  by  $T \mapsto f^*(T)$ .

**Problem 2.3**  $[\mathbf{2016.08.30.prob1}]$  For  $i \geq 0$  to construct an isomorphism of presheaves

$$S\mathcal{O}b_i : Sig(\mathcal{O}b_i) \rightarrow \mathcal{O}b_{i+1}$$

In constructing a solution of this problem and other problems where one needs to build an isomorphism of presheaves we will use the following lemma that is often used without an explicit reference.

**Lemma 2.4** [2016.11.14.11] *Let  $\Phi, \Phi' : \mathcal{C} \rightarrow \mathcal{D}$  be functors and  $\phi : \Phi \rightarrow \Phi'$  a natural transformation. Then  $\phi$  is an isomorphism of functors if and only if for all objects  $X$  of  $\mathcal{C}$  the morphism  $\phi_X : \Phi(X) \rightarrow \Phi'(X)$  is an isomorphism in  $\mathcal{D}$ .*

*The inverse isomorphism is formed by the family of morphisms  $\phi_X^{-1} = (\phi_X)^{-1}$ .*

**Proof:** One should first verify that the family  $Id_X$  forms the identity isomorphism of functors. This is immediate from the definitions.

If  $\phi$  is an isomorphism and  $\phi^{-1}$  is its inverse, then the functions  $\phi_X^{-1}$  form inverses to the functions  $\phi_X$ . This proves the “only if part”.

If all morphisms  $\phi_X$  are isomorphisms then the family  $(\phi_X)^{-1}$  forms a morphism of presheaves  $\phi^{-1} : \Phi' \rightarrow \Phi$ . Indeed, for  $f : X \rightarrow Y$  one has

$$\phi_X^{-1} \circ \Phi(f) = \Phi'(f) \circ \phi_Y^{-1}$$

This equality follows by taking its composition with  $\phi_X$  on the left and  $\phi_Y$  on the right. That  $\phi^{-1}$  is both the left and the right inverse to  $\phi$  is immediate from its definition. This proves the “if” part.

Next will also need the following lemma.

**Lemma 2.5** [2016.09.01.11] *Let  $\Gamma \in CC$ . Then one has:*

1. *if  $T \in \mathcal{O}b_1(\Gamma)$  and  $X \in \mathcal{O}b_i(T)$  then  $X \in \mathcal{O}b_{i+1}(\Gamma)$ ,*
2. *if  $X \in \mathcal{O}b_{i+1}(\Gamma)$  then  $ft^i(X) \in \mathcal{O}b_1(\Gamma)$  and  $X \in \mathcal{O}b_i(ft^i(X))$ .*

**Proof:** The first assertion follows from the equalities  $l(X) = l(T) + i = l(\Gamma) + 1 + i$  and  $ft^{i+1}(X) = ft(ft^i(X)) = ft(T) = \Gamma$ .

To prove the second assertion let  $X \in \mathcal{O}b_{i+1}(\Gamma)$ . Since  $l(X) \geq i$  we have  $l(ft^i(X)) = l(X) - i = l(\Gamma) + (i + 1) - i = l(\Gamma) + 1$ . The equality  $ft^1(ft^i(X)) = ft^{i+1}(X) = \Gamma$  is obvious and we conclude that  $ft^i(X) \in \mathcal{O}b_1(\Gamma)$ . Next, again because  $l(X) \geq i$ , we have  $l(X) = l(ft^i(X)) + i$  and since  $ft^i(X) = ft^i(X)$  we have that  $X \in \mathcal{O}b_i(ft^i(X))$ .

**Construction 2.6** [2016.08.30.constr1] Let  $\Gamma \in CC$ . Then  $Sig(\mathcal{O}b_i)(\Gamma)$  is the set of pairs  $(T, X)$  where  $T \in \mathcal{O}b_1(\Gamma)$  and  $X \in \mathcal{O}b_i(T)$ . By Lemma 2.5(1), the formula

$$[2016.09.01.eq4] S\mathcal{O}b_{i,\Gamma}(T, X) = X \tag{2.7}$$

defines a function  $Sig(\mathcal{O}b_i)(\Gamma) \rightarrow \mathcal{O}b_{i+1}(\Gamma)$ .

Conversely, by Lemma 2.5(2), the formula

$$[\mathbf{2016.09.01.eq5}] S\mathcal{O}b_{i,\Gamma}^{-1}(X) = (ft^i(X), X) \quad (2.8)$$

defines a function  $\mathcal{O}b_{i+1}(\Gamma) \rightarrow \text{Sig}(\mathcal{O}b_i)(\Gamma)$ .

If  $\Phi = S\mathcal{O}b_{i,\Gamma}$  and  $\Psi = S\mathcal{O}b_{i,\Gamma}^{-1}$  then

$$\Phi(\Psi(X)) = \Phi((ft^i(X), X)) = X$$

and

$$\Psi(\Phi(T, X)) = \Psi(X) = (ft^i(X), X) = (T, X)$$

where the last equality follows from the equality  $T = ft^i(X)$ . We conclude that  $S\mathcal{O}b_{i,\Gamma}$  and  $S\mathcal{O}b_{i,\Gamma}^{-1}$  are mutually inverse bijections.

In view of Lemma 2.4, it remains to verify that the family of bijections  $S\mathcal{O}b_{i,\Gamma}$  parametrized by  $\Gamma \in CC$  is a morphism of presheaves, that is, that for any  $f : \Gamma' \rightarrow \Gamma$  and  $(T, X) \in \text{Sig}(\mathcal{O}b_i)(\Gamma)$  we have

$$[\mathbf{2016.08.30.eq10}] \mathcal{O}b_{i+1}(f)(S\mathcal{O}b_{i,\Gamma}((T, X))) = S\mathcal{O}b_{i,\Gamma'}(\text{Sig}(\mathcal{O}b_i)(f)((T, X))) \quad (2.9)$$

Computing we get

$$\mathcal{O}b_{i+1}(f)(S\mathcal{O}b_{i,\Gamma}((T, X))) = f^*(X)$$

$$S\mathcal{O}b_{i,\Gamma'}(\text{Sig}(\mathcal{O}b_i)(f)((T, X))) = S\mathcal{O}b_{i,\Gamma'}(f^*(T), q(f, T)^*(X)) = q(f, T)^*(X)$$

and (2.9) follows from [?, Lemma 2.7]. This completes Construction 2.6.

As a corollary of Construction 2.6 and Lemma 2.4 we obtain the fact that the family of functions (2.8) parametrized by  $\Gamma \in CC$  is an isomorphism of presheaves that is inverse to  $S\mathcal{O}b_i$ .

We proceed now to the construction of isomorphisms  $S\widetilde{\mathcal{O}b}_i$ . Recall that for a morphism  $p : Y \rightarrow X$  we set

$$\text{sec}(p) = \{s \in \text{Mor}(X, Y) \mid s \circ p = \text{Id}_X\}$$

Elements of  $\text{sec}(p)$  are called sections of  $p$ .

The presheaves  $\widetilde{\mathcal{O}b}_n$  where defined in [?, Sec. 3]. On objects they are given by

$$[\mathbf{2016.11.15.eq6}] \widetilde{\mathcal{O}b}_n(\Gamma) = \{o \in \text{Mor}(CC) \mid \text{codom}(o) \in \mathcal{O}b_n(\Gamma), o \in \text{sec}(p_{\text{codom}(o)}), \text{codom}(o) > \Gamma\} \quad (2.10)$$

and on morphisms  $f : \Gamma' \rightarrow \Gamma$  by  $o \mapsto f^*(o)$ , where  $f^*(o)$  is defined in [?, Lemma 2.13].

For an element  $o \in \mathcal{O}b_n(\Gamma)$  we let  $\partial_\Gamma(o)$ , or simply  $\partial(o)$ , denote the object  $\text{codom}(o)$ .

Recall from [?, Sec. 3], that  $\widetilde{\mathcal{O}b}(CC)$  is the set of elements  $o \in \text{Mor}(CC)$  such that  $o \in \text{sec}(p_{\text{codom}(o)})$  and  $l(\text{codom}(o)) > 0$ . For such elements we also denote  $\text{codom}(o)$  by  $\partial(o)$ .

It follows easily from (2.10) that for  $\Gamma \in \text{Ob}(CC)$  and  $n > 0$  one has  $o \in \widetilde{\mathcal{O}b}_n(\Gamma)$  if and only if  $o \in \widetilde{\mathcal{O}b}(CC)$  and  $\partial(o) \in \mathcal{O}b_n(\Gamma)$ . It also follows from (2.10) that  $\mathcal{O}b_0(\Gamma) = \emptyset$ .



**Problem 2.7** [2016.08.30.prob2] For  $i \geq 1$  to construct an isomorphism of presheaves

$$S\widetilde{\mathcal{O}b}_i : \text{Sig}(\widetilde{\mathcal{O}b}_i) \rightarrow \widetilde{\mathcal{O}b}_{i+1}$$

**Lemma 2.8** [2016.11.18.11] Let  $\Gamma \in CC$ . Then one has:

1. if  $T \in \mathcal{O}b_1(\Gamma)$  and  $o \in \widetilde{\mathcal{O}b}_i(T)$  then  $o \in \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$ ,
2. if  $o \in \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$  then  $ft^i(\partial(o)) \in \mathcal{O}b_1(\Gamma)$  and  $o \in \widetilde{\mathcal{O}b}_i(ft^i(\partial(o)))$ .

**Proof:** If  $o \in \widetilde{\mathcal{O}b}_i(T)$  we have  $i > 0$  and therefore  $o \in \widetilde{\mathcal{O}b}(CC)$  and  $\partial(o) \in \mathcal{O}b_i(T)$ . By Lemma 2.5(1) we have  $\partial(o) \in \mathcal{O}b_{i+1}(\Gamma)$ . Therefore  $o \in \widetilde{\mathcal{O}b}_{i+1}(T)$ . This proves the first assertion.

If  $o \in \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$  then  $o \in \widetilde{\mathcal{O}b}(CC)$  and  $\partial(o) \in \mathcal{O}b_{i+1}(\Gamma)$ . By Lemma 2.5(2) we have  $ft^i(\partial(o)) \in \mathcal{O}b_1(\Gamma)$  and  $\partial(o) \in \mathcal{O}b_i(ft^i(\partial(o)))$ . Therefore  $o \in \widetilde{\mathcal{O}b}_i(ft^i(\partial(o)))$ .

We can now provide a construction for Problem 2.7.

**Construction 2.9** [2016.09.01.constr2] For  $\Gamma \in CC$  we have

$$\text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma) = \{(T, o) \mid T \in \mathcal{O}b_1(\Gamma), o \in \widetilde{\mathcal{O}b}_i(T)\}$$

For  $(T, o) \in \text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma)$  we have  $o \in \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$  by Lemma 2.8(1) and therefore the formula

$$[2016.09.01.eq6] S\widetilde{\mathcal{O}b}_{i,\Gamma}(T, o) = o \tag{2.11}$$

defines a function  $\text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma) \rightarrow \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$ .

If  $o \in \widetilde{\mathcal{O}b}_{i+1}(\Gamma)$  then by Lemma 2.8(2),  $ft^i(\partial(o)) \in \mathcal{O}b_1(\Gamma)$  and  $o \in \widetilde{\mathcal{O}b}_i(ft^i(\partial(o)))$ . Therefore the formula

$$[2016.09.01.eq7] S\widetilde{\mathcal{O}b}_{i,\Gamma}^{-1}(o) = (ft^i(\partial(o)), o) \tag{2.12}$$

defines a function  $\widetilde{\mathcal{O}b}_{i+1}(\Gamma) \rightarrow \text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma)$ .

One verifies in the same way as in Construction 2.6 that  $S\widetilde{\mathcal{O}b}_{i,\Gamma}$  and  $S\widetilde{\mathcal{O}b}_{i,\Gamma}^{-1}$  are mutually inverse bijections.

In view of Lemma 2.4 it remains to verify that the family of functions  $S\widetilde{\mathcal{O}b}_{i,\Gamma}$  parametrized by  $\Gamma \in CC$  is a morphism of functors, that is, that for  $f : \Gamma' \rightarrow \Gamma$  and  $(T, o) \in S\mathcal{O}b_{i,\Gamma}$  one has

$$[2016.09.01.eq2b] \widetilde{\mathcal{O}b}_{i+1}(f)(S\widetilde{\mathcal{O}b}_{i,\Gamma}(T, o)) = S\widetilde{\mathcal{O}b}_{i,\Gamma'}(\text{Sig}(\widetilde{\mathcal{O}b}_i)(f)(T, o)) \tag{2.13}$$

Computing we get

$$\widetilde{\mathcal{O}b}_{i+1}(f)(S\widetilde{\mathcal{O}b}_{i,\Gamma}(T, o)) = \widetilde{\mathcal{O}b}_{i+1}(f)(o) = f^*(o)$$

$$S\widetilde{\mathcal{O}b}_{i,\Gamma'}(\text{Sig}(\widetilde{\mathcal{O}b}_i)(f)(T, o)) = S\widetilde{\mathcal{O}b}_{i,\Gamma'}(f^*(T), q(f, T)^*(o)) = q(f, T)^*(o)$$

and we conclude that (2.13) holds by [?, Lemma 2.15].

This completes Construction 2.9.

As a corollary of Construction 2.9 and Lemma 2.4 we obtain the fact that the family of functions (2.12) parametrized by  $\Gamma \in CC$  is an isomorphism of presheaves that is inverse to  $S\widetilde{\mathcal{O}b}_i$ .

**Lemma 2.10** [2016.12.04.11] *For any  $i \geq 1$  the square of morphisms of presheaves*

$$\begin{array}{ccc}
\text{Sig}(\widetilde{\mathcal{O}b}_i) & \xrightarrow{S\widetilde{\mathcal{O}b}_i} & \widetilde{\mathcal{O}b}_{i+1} \\
\text{[2016.12.04.eq1]} \downarrow \text{Sig}(\partial) & & \downarrow \partial \\
\text{Sig}(\mathcal{O}b_i) & \xrightarrow{S\mathcal{O}b_i} & \text{Sig}(\mathcal{O}b_i)
\end{array} \tag{2.14}$$

*commutes.*

**Proof:** Let  $\Gamma \in CC$ . By definition we have

$$\text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma) = \{(T, o) \mid T \in \mathcal{O}b_1(\Gamma), o \in \widetilde{\mathcal{O}b}_i(T)\}$$

Let  $(T, o) \in \text{Sig}(\widetilde{\mathcal{O}b}_i)(\Gamma)$ . Then, again by definitions,

$$\partial_\Gamma(S\widetilde{\mathcal{O}b}_{i,\Gamma}(T, o)) = \partial_\Gamma(o)$$

and

$$S\mathcal{O}b_{i,\Gamma}(\text{Sig}(\partial)_\Gamma(T, o)) = S\mathcal{O}b_{i,\Gamma}(T, \partial_\Gamma(o)) = \partial_\Gamma(o)$$

The lemma is proved.

**Remark 2.11** [2016.11.18.rem1] Define  $\text{Sig}^i$  by induction on  $i$ , setting  $\text{Sig}^0 = Id_{\text{PreShv}(CC)}$  and  $\text{Sig}^{i+1} = \text{Sig}^i \circ \text{Sig}$ . Then, also by induction on  $i$ , we can construct isomorphisms

$$S\mathcal{O}b_j^i : \text{Sig}^i(\mathcal{O}b_j) \rightarrow \mathcal{O}b_{i+j}$$

where  $S\mathcal{O}b_j^0 = Id_{\mathcal{O}b_j}$  and  $S\mathcal{O}b_j^{i+1}$  is the composition

$$\text{Sig}^{i+1}(\mathcal{O}b_j) = \text{Sig}(\text{Sig}^i(\mathcal{O}b_j)) \xrightarrow{\text{Sig}(S\mathcal{O}b_j^i)} \text{Sig}(\mathcal{O}b_{i+j}) \xrightarrow{S\mathcal{O}b_{i+j}} \mathcal{O}b_{i+j+1}$$

In exactly the same way we construct isomorphisms

$$S\widetilde{\mathcal{O}b}_j^i : \text{Sig}^i(\widetilde{\mathcal{O}b}_j) \rightarrow \widetilde{\mathcal{O}b}_{i+j}$$

## 2.2 The functor $D_p$

In this section we work in the context of a category  $\mathcal{C}$  with a universe  $p$ . The goal of the section is to construct, for any such pair, a functor

$$D_p : \text{PreShv}(\mathcal{C}) \rightarrow \text{PreShv}(\mathcal{C})$$

The definition of a universe in a category was given in [?, Definition 2.1]. We repeat it here for the convenience of the reader.

**Definition 2.12** [2009.11.1.def1] Let  $\mathcal{C}$  be a category. A universe structure on a morphism  $p : \tilde{U} \rightarrow U$  in  $\mathcal{C}$  is a mapping that assigns to any morphism  $f : X \rightarrow U$  in  $\mathcal{C}$  a pullback of the form

$$\begin{array}{ccc} (X, f) & \xrightarrow{Q(F)} & \tilde{U} \\ \text{[2016.12.02.eq8]} p_{X,F} \downarrow & & \downarrow p \\ X & \xrightarrow{F} & U \end{array} \quad (2.15)$$

A universe in  $\mathcal{C}$  is a morphism together with a universe structure on it.

We usually refer to a universe by the name of the corresponding morphism without mentioning the choices of pullbacks explicitly. To shorten the notation we will write  $p_F$  instead of  $p_{X,F}$ .

For  $f : W \rightarrow X$  and  $g : W \rightarrow \tilde{U}$  such that  $f \circ F = g \circ p$  we will denote by  $f *_F g$  the unique morphism  $W \rightarrow (X; F)$  such that

$$\begin{aligned} \text{[2016.11.10.eq1]} (f *_F g) \circ p_F &= f \\ (f *_F g) \circ Q(F) &= g \end{aligned} \quad (2.16)$$

For  $X' \xrightarrow{f} X \xrightarrow{F} U$  we let  $Q(f, F)$  denote the morphism

$$\text{[2016.12.02.eq4]} Q(f, F) = (p_{f \circ F} \circ f) *_F Q(f \circ F) : (X'; f \circ F) \rightarrow (X; F) \quad (2.17)$$

Observe that one has

$$\text{[2016.08.24.eq4]} Q(f \circ F) = Q(f, F) \circ Q(F) \quad (2.18)$$

$$\text{[2016.08.26.eq2]} Q(Id_X, F) = Id_{(X;F)} \quad (2.19)$$

$$\text{[2016.08.26.eq3]} Q(f' \circ f, F) = Q(f', f \circ F) \circ Q(f, F) \quad (2.20)$$

where the first equality follows directly from the definition, the second from the definition and the uniqueness of the morphisms  $f *_F g$  satisfying (2.2) and the third is proved in [?, Lemma 2.5].

Let us fix a category  $\mathcal{C}$  and a universe  $p$  in it.

For any  $\mathcal{G} \in PreShv(\mathcal{C})$  we define functor data  $D_p(\mathcal{G})$  given on objects by

$$\text{[2016.08.30.eq4]} D_p(\mathcal{G})(X) := \coprod_{F: X \rightarrow U} \mathcal{G}((X; F)) \quad (2.21)$$

and on morphisms by

$$\text{[2016.08.30.eq5]} D_p(\mathcal{G})(f) : (F, \gamma) \mapsto (f \circ F, \mathcal{G}(Q(f, F))(\gamma)) \quad (2.22)$$

**Lemma 2.13** [2016.09.07.11] *The functor data  $D_p(\mathcal{G})$  specified above is a presheaf, i.e., one has*

1. for any  $X \in \mathcal{C}$ ,  $D_p(\mathcal{G})(Id_X) = Id_{D_p(\mathcal{G})(X)}$ ,
2. for any  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$  in  $\mathcal{C}$ ,

$$D_p(\mathcal{G})(f \circ g) = D_p(\mathcal{G})(g) \circ D_p(\mathcal{G})(f)$$

**Proof:** For the first property we have

$$D_p(\mathcal{G})(Id_X)((F, \gamma)) = (Id_X \circ F, \mathcal{G}(Q(Id_X, F))(\gamma)) = (F, \gamma)$$

where the second equality is by (2.19) and the identity morphism axiom from the presheaf  $\mathcal{G}$ .

For the second one we have

$$\begin{aligned} D_p(\mathcal{G})(f \circ g)(F, \gamma) &= (f \circ g \circ F, \mathcal{G}(Q(f \circ g, F))(\gamma)) = \\ &= (f \circ g \circ F, \mathcal{G}(Q(f, g \circ F) \circ Q(g, F))(\gamma)) = (f \circ (g \circ F), \mathcal{G}(Q(f, g \circ F))(\mathcal{G}(Q(g, F))(\gamma))) = \\ &= D_p(\mathcal{G})(f)(D_p(\mathcal{G})(g)(F, \gamma)) = (D_p(\mathcal{G})(g) \circ D_p(\mathcal{G})(f))(F, \gamma) \end{aligned}$$

where the second equality is by (2.20) and the third one by the composition axiom of the presheaf  $\mathcal{G}$ .

One defines  $D_p$  on morphisms of presheaves  $r : \mathcal{G} \rightarrow \mathcal{G}'$  by the family of morphisms

$$[2016.08.30.eq6] D_p(r)_X(F, \gamma) = (F, r_{(X;F)}(\gamma)) \quad (2.23)$$

For  $f : X \rightarrow X'$  and  $r : \mathcal{G} \rightarrow \mathcal{G}'$  we have

$$[2016.11.14.eq2] D_p(\mathcal{G})(f) \circ D_p(r)_X = D_p(r)_{X'} \circ D_p(\mathcal{G}')(f) \quad (2.24)$$

that is, the family of functions  $D_p(r)_X$  parametrized by  $X \in \mathcal{C}$  is a morphism of presheaves.

For  $\mathcal{G} \in PreShv(\mathcal{C})$  we have

$$D_p(Id_{\mathcal{G}})_X = Id_{D_p(\mathcal{G})(X)}$$

and for  $r : \mathcal{G} \rightarrow \mathcal{G}'$  and  $r' : \mathcal{G}' \rightarrow \mathcal{G}''$  we have

$$[2016.12.18.eq4] D_p(r \circ r')_X = D_p(r)_X \circ D_p(r')_X \quad (2.25)$$

These two equalities show that the functor data given by  $D_p$  on presheaves and  $D_p$  on morphisms of presheaves is a functor that we also denote by

$$D_p : PreShv(\mathcal{C}) \rightarrow PreShv(\mathcal{C})$$

Note that for the presheaves of the form  $Yo(A)$ , where  $Yo$  is the Yoneda embedding, we have

$$[2016.11.14.eq4] D_p(Yo(A))(X) = \Pi_{F: X \rightarrow U} Mor_{\mathcal{C}}((X; F), A) \quad (2.26)$$

and for a morphism  $f : X \rightarrow X'$ ,

$$[2016.11.14.eq4a] D_p(Y \circ A)(f)(F_1, F_2) = (f \circ F_1, Q(f, F_1) \circ F_2) \quad (2.27)$$

For a morphism  $a : A' \rightarrow A$  we have

$$[2016.12.02.eq6] D_p(Y \circ a)_X(F_1, F_2) = (F_1, F_2 \circ a) \quad (2.28)$$

**Remark 2.14** [2015.07.29.rem2] Define

$$D_p^n(X, Y) = D_p^n(Y \circ Y)(X)$$

such that in particular one has

$$D_p^0(X, Y) = \text{Mor}_{\mathcal{C}}(X, Y)$$

It is likely that there are composition functions

$$[2016.12.18.eq3] D_p^n(X, Y) \times D_p^m(Y, Z) \rightarrow D_p^{n+m}(X, Z) \quad (2.29)$$

that satisfy the unity and associativity axioms and one obtains, from any universe category  $(\mathcal{C}, p)$ , a new category  $(\mathcal{C}, p)_*$  with the same collection of objects and morphisms between two objects given by

$$\text{Mor}_{(\mathcal{C}, p)_*}(X, Y) = \coprod_{n \geq 0} D_p^n(X, Y)$$

We will use the sets  $D_p^n(X, Y)$  below, but the study of the possible composition functions (2.29) and categories  $(\mathcal{C}, p)_*$  is deferred to a later paper.

**Remark 2.15** [2016.12.14.rem2] The observations of Remark 2.2 apply, with obvious modifications, to the construction  $D_p$  as well.

## 2.3 Isomorphisms of presheaves $u_1$ and $\tilde{u}_1$

[Sec.2.3]

We now consider a universe category, that is, a category  $\mathcal{C}$  with a universe  $p$  and a choice of a final object  $pt$ . We usually denote a universe category as  $(\mathcal{C}, p)$  without mentioning the final object. For any universe category we have constructed in [?, Section 2] a C-system  $CC(\mathcal{C}, p)$ .

The main goal of this section is to provide constructions for Problems 2.17 and 2.20.

Let us first recall the construction of  $CC(\mathcal{C}, p)$ . One defines first, by induction on  $n$ , pairs  $(Ob_n, int_n : Ob_n \rightarrow \mathcal{C})$  where  $Ob_n = Ob_n(\mathcal{C}, p)$  is a set and  $int_n$  is a function from  $Ob_n$  to objects of  $\mathcal{C}$ . The definition is as follows:

1.  $Ob_0$  is the standard one point set *unit* whose element we denote by  $tt$ . The function  $int_0$  maps  $tt$  to the final object  $pt$  of the universe category structure on  $\mathcal{C}$ ,
2.  $Ob_{n+1} = \coprod_{A \in Ob_n} Mor(int(A), U)$  and  $int_{n+1}(A, F) = (int(A); F)$ .

We then define  $Ob(CC(\mathcal{C}, p))$  as  $\coprod_{n \geq 0} Ob_n$  such that elements of  $Ob(CC(\mathcal{C}, p))$  are pairs  $\Gamma = (n, A)$  where  $A \in Ob_n(\mathcal{C}, p)$ . We define the function  $int : Ob(CC(\mathcal{C}, p)) \rightarrow \mathcal{C}$  as the sum of functions  $int_n$ . Where no confusion between  $int$  and  $int_n$  is likely we will omit the index  $n$  at  $int_n$ .

The morphisms in  $CC(\mathcal{C}, p)$  are defined by

$$Mor_{CC(\mathcal{C}, p)} = \coprod_{\Gamma, \Gamma' \in Ob(CC)} Mor_{\mathcal{C}}(int(\Gamma), int(\Gamma'))$$

and the function  $int$  on morphisms maps a triple  $((\Gamma, \Gamma'), a)$  to  $a$ . Note that the subset in  $Mor$  that consists of  $f$  such that  $dom(f) = \Gamma$  and  $codom(f) = \Gamma'$  is not equal to the set  $Mor_{\mathcal{C}}(int(\Gamma), int(\Gamma'))$  but instead to the set of triples of the form  $f = ((\Gamma, \Gamma'), a)$  where  $a \in Mor_{\mathcal{C}}(int(\Gamma), int(\Gamma'))$ . The functor  $int$  maps  $((\Gamma, \Gamma'), a)$  to  $a$ . This map is bijective and therefore the functor is fully faithful but its morphism component is not the identity function.

The length function is defined by  $l((n, A)) = n$ .

One defines  $pt$  as  $pt = (0, tt)$ . It is the only object of length 0.

If  $\Gamma = (n, B)$  where  $n > 0$  then, by construction,  $B = (A, F)$  where  $F : int(A) \rightarrow U$ . The  $ft$  function is defined on such  $\Gamma$  by  $ft(\Gamma) = (n - 1, A)$  and on  $pt$  by  $ft(pt) = pt$ .

**Lemma 2.16** [2016.08.22.11] *For  $\Gamma = (n, A)$  and  $T = (n', B) \in Ob(CC(\mathcal{C}, p))$  one has  $T \in \mathcal{O}b_1(\Gamma)$  if and only if  $n' = n + 1$  and there exists  $F : int(A) \rightarrow U$  such that  $B = (A, F)$ .*

**Proof:** By definition of the length function  $l$ , we have  $l(\Gamma) = n$  and  $l(T) = n'$ . By definition of  $\mathcal{O}b_1$ ,  $T \in \mathcal{O}b_1(\Gamma)$  if and only if  $n' = n + 1$  and  $ft(T) = \Gamma$ .

If  $T = (n + 1, (A, F))$  then  $n' = n + 1$ . In particular,  $l(T) > 0$  and therefore  $ft(T) = (n, A) = \Gamma$ . This proves the "if" part.

Assume that  $T = (n', B) \in \mathcal{O}b_1(\Gamma)$ . Then  $n' = n + 1$ . Since  $n' > 0$ ,  $B$  is a pair of the form  $(A', F)$ . Since  $ft(T) = (n, A') = (n, A)$  we have  $A' = A$ . This proves the "only if" part.

The  $p$ -morphism for  $\Gamma = (n, A)$  where  $n > 0$  and  $A = (B, F)$  is given by  $((\Gamma, ft(\Gamma)), p_F)$  where  $p_F$  are the  $p$ -morphisms of the universe structure.

For  $f : (n, A') \rightarrow (n, A)$  and  $T$  such that  $l(T) = l(\Gamma) + 1$  and  $ft(T) = \Gamma$  one has, by Lemma 2.16,  $T = (n + 1, (A, F))$  and one defines

$$[2016.08.22.eq2] f^*(T) = (n + 1, (A', int(f) \circ F)) \quad (2.30)$$

and

$$[2016.08.22.eq3] q(f, T) = ((f^*(T), T), Q(int(f), F)) \quad (2.31)$$

The C-system axioms are verified in [?].

Let us denote by

$$int^\circ : PreShv(\mathcal{C}) \rightarrow PreShv(CC)$$

is the functor of pre-composition with  $int^{op}$  and by

$$Yo : \mathcal{C} \rightarrow PreShv(\mathcal{C})$$

the Yoneda embedding of  $\mathcal{C}$ .

**Problem 2.17** [2015.04.30.prob1a] *To construct an isomorphism of presheaves*

$$[2016.11.12.eq2]u_1 : \mathcal{O}b_1 \rightarrow int^\circ(Yo(U)) \quad (2.32)$$

such that for  $\Gamma = (n, A)$  and  $T = (n + 1, (A, F))$  one has

$$[2015.04.30.eq3a]u_{1,\Gamma}(T) = F \quad (2.33)$$

**Construction 2.18** [2016.08.22.constr1] By definition of  $int^\circ$  and  $Yo$  and Lemma 2.4, an isomorphism of presheaves of the form (2.32) is a family of functions of the form

$$u_{1,\Gamma} : \mathcal{O}b_1(\Gamma) \rightarrow Mor_{\mathcal{C}}(int(\Gamma), U)$$

parametrized by  $\Gamma \in Ob(CC(\mathcal{C}, p))$  such that for any  $f : \Gamma' \rightarrow \Gamma$  and any  $T \in \mathcal{O}b_1(\Gamma)$  one has

$$[2015.04.30.eq1a]u_{1,\Gamma'}(f^*(T)) = int(f) \circ u_{1,\Gamma}(T) \quad (2.34)$$

and for any  $\Gamma$  the function  $u_{1,\Gamma}$  is a bijection.

By Lemma 2.16, the conditions (2.33) define our family completely and it remains to verify (2.34) and the bijectivity condition.

For  $\Gamma = (n, A)$ ,  $T = (n + 1, (A, F))$ ,  $\Gamma' = (n', A')$  and  $f : \Gamma' \rightarrow \Gamma$  we have, by (2.30),

$$f^*(T) = (n' + 1, (A', int(f) \circ F))$$

Therefore,

$$u_{1,\Gamma'}(f^*(T)) = u_{1,\Gamma'}((n' + 1, (A', int(f) \circ F))) = int(f) \circ F = int(f) \circ u_{1,\Gamma}(T)$$

which proves (2.34).

By Lemma 2.16, for  $\Gamma = (n, A)$ , the formula  $F \mapsto (n + 1, (A, F))$  defines a function

$$Mor_{\mathcal{C}}(int(A), U) \rightarrow \mathcal{O}b_1(\Gamma)$$

By the same lemma and (2.33) this function is inverse to  $u_{1,\Gamma}$ . This proves the bijectivity condition and completes Construction 2.18.

Using again Lemma 2.16 and (2.33) we see that for any  $\Gamma \in Ob(CC(\mathcal{C}, p))$  and  $T \in \mathcal{O}b_1(\Gamma)$ ,

$$[2015.05.02.eq1a] \text{int}(T) = (\text{int}(\Gamma); u_{1,\Gamma}(T)) \quad (2.35)$$

and

$$[2016.08.24.eq3] \text{int}(p_T) = p_{u_{1,\Gamma}(T)} \quad (2.36)$$

For  $f : \Gamma' \rightarrow \Gamma$  and  $T$  as above we have

$$[2016.08.30.eq3] \text{int}(q(f, T)) = Q(\text{int}(f), u_{1,\Gamma}(T)) \quad (2.37)$$

**Lemma 2.19** [2016.08.22.12] For  $\Gamma = (n, A)$  and  $o \in \widetilde{\mathcal{O}b}_1(\Gamma)$  one has

$$[2016.08.22.eq1] \text{codom}(\text{int}(o)) = (\text{int}(\Gamma); u_{1,\Gamma}(\partial(o))) \quad (2.38)$$

**Proof:** We have  $\text{codom}(o) = \partial(o) \in \mathcal{O}b_1(\Gamma)$ . Therefore (2.38) follows from the equality  $\text{codom}(\text{int}(f)) = \text{int}(\text{codom}(f))$  and (2.35).

The second problem whose solution is constructed in this section is as follows.

**Problem 2.20** [2015.04.30.prob1b] To construct an isomorphism of presheaves

$$[2016.11.12.eq3] \tilde{u}_1 : \widetilde{\mathcal{O}b}_1 \rightarrow \text{int}^\circ(Yo(\tilde{U})) \quad (2.39)$$

such that for  $o \in \widetilde{\mathcal{O}b}_1(\Gamma)$  one has

$$[2015.04.30.eq4a] \tilde{u}_{1,\Gamma}(o) = \text{int}(o) \circ Q(u_{1,\Gamma}(\partial(o))) \quad (2.40)$$

where the right hand side is defined by (2.38) and the equality  $\text{dom}(Q(F)) = (\text{dom}(F); F)$ .

To construct a solution for this problem we will need the following two lemmas.

**Lemma 2.21** [2016.08.26.11] For a universe  $p$  in  $\mathcal{C}$  and  $X \in \mathcal{C}$ , the function

$$\amalg_{F \in \text{Mor}(X, U)} \text{sec}(p_F) \rightarrow \text{Mor}(X, \tilde{U})$$

given by the formula  $(F, s) \mapsto s \circ Q(F)$  is a bijection. The inverse bijection is given by the formula  $\tilde{F} \mapsto (\tilde{F} \circ p, Id_X *_{\tilde{F} \circ p} \tilde{F})$  where  $Id_X *_{\tilde{F} \circ p} \tilde{F}$  is defined because  $Id_X \circ \tilde{F} \circ p = \tilde{F} \circ p$ .

**Proof:** Let us denote the first function by  $\Phi$  and second one by  $\Psi$ . We have

$$\Phi(\Psi(\tilde{F})) = \Phi(\tilde{F} \circ p, Id_X *_{\tilde{F} \circ p} \tilde{F}) = (Id_X *_{\tilde{F} \circ p} \tilde{F}) \circ Q(\tilde{F} \circ p) = \tilde{F}$$

where the last equality is by the definition of  $*_{\tilde{F} \circ p}$ , and

$$\Psi(\Phi(F, s)) = \Psi(s \circ Q(F)) = ((s \circ Q(F)) \circ p, Id_X *_{(s \circ Q(F)) \circ p} (s \circ Q(F)))$$



Next we have

$$[\mathbf{2016.11.12.eq1}] s \circ Q(F) \circ p = s \circ p_F \circ F = F \quad (2.41)$$

It remains to compare  $Id_X *_{s \circ Q(F) \circ p} (s \circ Q(F))$  with  $s$ . To do it we need to compare its post-compositions with  $p_F$  and  $Q(F)$  with the same post-compositions for  $s$ .

By (2.41) we may replace  $s \circ Q(F) \circ p$  with  $F$ . We have

$$\begin{aligned} Id_X *_{F} (s \circ Q(F)) \circ p_F &= Id_X = s \circ p_F \\ Id_X *_{F} (s \circ Q(F)) \circ Q(F) &= s \circ Q(F) = s \circ Q(F) \end{aligned}$$

Therefore,  $Id_X *_{F} (s \circ Q(F)) = s$  and

$$\Psi(\Phi(F, s)) = (F, s)$$

The lemma is proved.

**Lemma 2.22** [2016.08.26.14] *Let  $p : Y \rightarrow X$  be a morphism in  $\mathcal{C}$  and  $\Phi : \mathcal{C} \rightarrow \mathcal{C}'$  a functor. Then for  $s \in \text{sec}(p)$  one has  $\Phi(s) \in \text{sec}(\Phi(p))$ .*

*If  $\Phi$  is fully faithful then the resulting function*

$$\Phi_{\text{sec}, p} : \text{sec}(p) \rightarrow \text{sec}(\Phi(p))$$

*is a bijection.*

**Proof:** The first assertion follows immediately from the definition of  $\text{sec}$  and the axioms of a functor.

Assume that  $\Phi$  is fully faithful. To prove that  $\Phi_{\text{sec}, p}$  is a bijection let

$$\Phi_{A,B}^{-1} : \text{Mor}_{\mathcal{C}'}(\Phi(A), \Phi(B)) \rightarrow \text{Mor}_{\mathcal{C}}(A, B)$$

be the inverse to the function  $\Phi_{A,B} : \text{Mor}_{\mathcal{C}}(A, B) \rightarrow \text{Mor}_{\mathcal{C}'}(\Phi(A), \Phi(B))$  that we denoted simply by  $\Phi$ . One verifies easily that for any  $A, B, C \in \mathcal{C}$  the functions  $\Phi_{A,B}^{-1}$ ,  $\Phi_{B,C}^{-1}$  and  $\Phi_{A,C}^{-1}$  commute with the compositions and for any  $A \in \mathcal{C}$  one has  $\Phi_{A,A}^{-1}(Id_{\Phi(A)}) = Id_A$ .

Therefore, for  $s' \in \text{sec}(\Phi_{Y,X}(p))$  we have  $\Phi_{X,Y}^{-1}(s') \in \text{sec}(p)$ . Indeed,

$$\Phi_{X,X}(\Phi_{X,Y}^{-1}(s') \circ p) = \Phi_{X,Y}(\Phi_{X,Y}^{-1}(s')) \circ \Phi_{Y,X}(p) = s' \circ \Phi_{Y,X}(p) = Id_{\Phi(X)}$$

and since  $\Phi_{X,X}^{-1}(Id_{\Phi(X)}) = Id_X$  we obtain that  $\Phi_{X,Y}^{-1}(s') \circ p = Id_X$ . This implies that  $\Phi_{\text{sec}, p}$  and the restriction of  $\Phi_{X,Y}^{-1}$  to  $\text{sec}(\Phi(p))$  form a pair of functions between  $\text{sec}(p)$  and  $\text{sec}(\Phi(p))$  and one sees immediately that they are mutually inverse.

**Construction 2.23** [2016.08.22.constr2] By definition of  $\text{int}^\circ$  and  $Yo$  and Lemma 2.4, an isomorphism of presheaves of the form (2.39) is a family of functions of the form

$$\tilde{u}_{1,\Gamma} : \widetilde{\mathcal{O}b}_1(\Gamma) \rightarrow \text{Mor}_{\mathcal{C}}(\text{int}(\Gamma), \tilde{U})$$

parametrized by  $\Gamma \in Ob(CC(\mathcal{C}, p))$  such that for any  $f : \Gamma' \rightarrow \Gamma$  and any  $o \in \widetilde{\mathcal{O}b}_1(\Gamma)$  one has

$$[\mathbf{2015.04.30.eq1b}] \widetilde{u}_{1,\Gamma'}(f^*(o)) = int(f) \circ \widetilde{u}_{1,\Gamma}(o) \quad (2.42)$$

and for any  $\Gamma$  the function  $\widetilde{u}_{1,\Gamma}$  is a bijection.

The equalities (2.40) define our family completely and it remains to prove (2.42) and the bijectivity condition.

For the proof of (2.42) we have the following, where we write  $u$  instead of  $u_{1,\Gamma}$  and  $u_{1,\Gamma'}$  and  $\widetilde{u}$  instead of  $\widetilde{u}_{1,\Gamma}$  and  $\widetilde{u}_{1,\Gamma'}$ ,

$$\begin{aligned} \widetilde{u}(f^*(o)) &= int(f^*(o)) \circ Q(u(\partial(f^*(o)))) = int(f^*(o)) \circ Q(u(f^*(\partial(o)))) = \\ int(f^*(o)) \circ Q(int(f) \circ u(\partial(o))) &= int(f^*(o)) \circ Q(int(f), u(\partial(o))) \circ Q(u(\partial(o))) = \\ int(f^*(o)) \circ int(q(f, \partial(o))) \circ Q(u(\partial(o))) &= int(f^*(o) \circ q(f, \partial(o))) \circ Q(u(\partial(o))) = \\ int(q(f, \Gamma) \circ o) \circ Q(u(\partial(o))) &= int(f \circ o) \circ Q(u(\partial(o))) = \\ int(f) \circ int(o) \circ Q(u(\partial(o))) &= int(f) \circ \widetilde{u}(o) \end{aligned}$$

where the first equality is by (2.40), second is by definition of  $f^*(o)$ , the third is by (2.34), the fourth is by (2.18), the fifth is by (2.37), the sixth is because  $int$  is a functor, the seventh is by [?, (2.19)], the eighth is by definition of  $q(f, -)$ , the ninth is because  $int$  is a functor and the tenth is again by (2.40). This completes the proof of (2.42).

To prove that the function  $\widetilde{u}_{1,\Gamma}$  is a bijection we will represent it as the composition of functions that we can show to be bijections. The functions are of the form

$$\widetilde{\mathcal{O}b}_1(\Gamma) \rightarrow \Pi_{T \in \mathcal{O}b_1(\Gamma)} \partial^{-1}(T) \rightarrow \Pi_{F: int(\Gamma) \rightarrow U} sec(p_F) \rightarrow Mor(int(\Gamma), \widetilde{U})$$

and are given by the formulas

$$o \mapsto (\partial(o), o) \quad (T, o) \mapsto (u(T), int(o)) \quad (F, s) \mapsto s \circ Q(F)$$

The first function is the function  $X \rightarrow \Pi_{y \in Y} f^{-1}(y)$ , which is defined and is a bijection for any function of sets  $f : X \rightarrow Y$ . The second one is the total function of the function  $u$  and the family of functions  $int_{sec, p_T}$  of Lemma 2.22. Since  $u$  and the functions  $int_{sec, p_T}$  are bijections the total function is a bijection. The third function is the bijection of Lemma 2.21.

Let us show that the composition of these bijections equals  $\widetilde{u}$ . Indeed, for  $o \in \widetilde{\mathcal{O}b}_1(\Gamma)$  we have

$$o \mapsto (\partial(o), o) \mapsto (u(\partial(o)), int(o)) \mapsto int(o) \circ Q(u(\partial(o))) = \widetilde{u}(o)$$

This completes Construction 2.23.

**Remark 2.24** [2016.08.26.rem1] The inverse to  $\widetilde{u}_{1,\Gamma}$  can be expressed by the formula

$$\widetilde{u}_{1,\Gamma}^{-1}(H) = int_{\Gamma, u_{1,\Gamma}^{-1}(H \circ p)}^{-1}(Id_{int(\Gamma)} *_{H \circ p} H)$$

Note that while we can omit explicitly mentioning  $dom(f)$  and  $codom(f)$  when we write  $int(f)$  we must specify them when we write  $int^{-1}(f)$  because  $int$  is bijective only on the subsets of morphisms with fixed domain and codomain. This makes the expression for  $\widetilde{u}_{1,\Gamma}^{-1}$  longer than one would prefer.

The family of functions  $\partial_\Gamma$  forms a morphism of presheaves  $\widetilde{\mathcal{O}b}_n \rightarrow \mathcal{O}b_n$  that we usually denote simply by  $\partial$ .

**Lemma 2.25** [2016.12.02.14] *The square of morphisms of presheaves*

$$\begin{array}{ccc}
\widetilde{\mathcal{O}b}_1 & \xrightarrow{\tilde{u}_1} & \text{int}^\circ(Yo(\widetilde{U})) \\
\text{[2016.08.20.eq1]} \downarrow \partial & & \downarrow \text{int}^\circ(Yo(p)) \\
\mathcal{O}b_1 & \xrightarrow{u_1} & \text{int}^\circ(Yo(U))
\end{array} \tag{2.43}$$

*commutes.*

**Proof:** For  $\Gamma$  and  $o \in \widetilde{\mathcal{O}b}_1(\Gamma)$  we have

$$\begin{aligned}
\text{int}^\circ(Yo(p))_\Gamma(\tilde{u}_{1,\Gamma}(o)) &= (\tilde{u}_{1,\Gamma}(o)) \circ p = \text{int}(o) \circ Q(u_{1,\Gamma}(\partial(o))) \circ p = \\
\text{int}(o) \circ p_{u_{1,\Gamma}(\partial(o))} \circ u_{1,\Gamma}(\partial(o)) &= \text{int}(o \circ p_{\partial(o)}) \circ u_{1,\Gamma}(\partial(o)) = u_{1,\Gamma}(\partial(o))
\end{aligned}$$

where the first equality is by definition of  $\text{int}^\circ$  and  $Yo$ , the second by (2.40), the third by commutativity of (2.15), the fourth by (2.36) and the fifth by the definition  $\widetilde{\mathcal{O}b}_1(\Gamma)$  in (2.10) and the fact that  $\partial(o) = \text{codom}(o)$ . The lemma is proved.

## 2.4 Functor isomorphisms $SD_p$

In this section we continue to consider a universe category  $(\mathcal{C}, p)$ . For any  $(\mathcal{C}, p)$  we will relate the functor  $D_p$  on  $PreShv(\mathcal{C})$  and the functor  $Sig$  on  $PreShv(CC(\mathcal{C}, p))$ .

**Problem 2.26** [2016.08.28.probl] *For a universe category  $(\mathcal{C}, p)$  to construct an isomorphism of functors  $PreShv(\mathcal{C}) \rightarrow PreShv(CC)$  of the form*

$$SD_p : \text{int}^\circ \circ Sig \rightarrow D_p \circ \text{int}^\circ$$

**Construction 2.27** [2016.08.28.constr1] In view of Lemma 2.4, we have to construct, for any  $\mathcal{G} \in PreShv(\mathcal{C})$ , an isomorphism of presheaves on  $CC$  of the form

$$SD_{p,\mathcal{G}} : Sig(\text{int}^{op} \circ \mathcal{G}) \rightarrow \text{int}^{op} \circ D_p(\mathcal{G})$$

and to show that these isomorphisms are natural in  $\mathcal{G}$ , that is, that for a morphism of presheaves  $r : \mathcal{G} \rightarrow \mathcal{G}'$  one has

$$SD_{p,\mathcal{G}} \circ \text{int}^\circ(D_p(r)) = Sig(\text{int}^\circ(r)) \circ SD_{p,\mathcal{G}'}$$

Applying Lemma 2.4 again, we see that we need to construct, for each  $\mathcal{G}$  and  $\Gamma \in CC$ , a bijection  $SD_{p,\mathcal{G},\Gamma}$ , which we will denote  $\phi_{\mathcal{G},\Gamma}$  for the duration of the proof, of the form

$$\phi_{\mathcal{G},\Gamma} : Sig(\text{int}^{op} \circ \mathcal{G})(\Gamma) \rightarrow (\text{int}^{op} \circ D_p(\mathcal{G}))(\Gamma) = D_p(\mathcal{G})(\text{int}(\Gamma))$$

and to show that two conditions hold:

1. for any  $f : \Gamma' \rightarrow \Gamma$  we have

$$[\mathbf{2016.08.30.eq1}] \phi_{\mathcal{G},\Gamma} \circ D_p(\mathcal{G})(\text{int}(f)) = \text{Sig}(\text{int}^{op} \circ \mathcal{G})(f) \circ \phi_{\mathcal{G},\Gamma'} \quad (2.44)$$

that is, the square

$$\begin{array}{ccc} \text{Sig}(\text{int}^{op} \circ \mathcal{G})(\Gamma) & \xrightarrow{\phi_{\mathcal{G},\Gamma}} & D_p(\mathcal{G})(\text{int}(\Gamma)) \\ [\mathbf{2016.11.19.eq1}] \text{int}^{op} \circ \mathcal{G}(f) \downarrow & & \downarrow D_p(\mathcal{G})(\text{int}(f)) \\ \text{Sig}(\text{int}^{op} \circ \mathcal{G})(\Gamma') & \xrightarrow{\phi_{\mathcal{G},\Gamma'}} & D_p(\mathcal{G})(\text{int}(\Gamma')) \end{array} \quad (2.45)$$

commutes.

2. for any  $r : \mathcal{G} \rightarrow \mathcal{G}'$  and  $\Gamma \in CC$  we have

$$[\mathbf{2016.08.30.eq2}] \phi_{\mathcal{G},\Gamma} \circ D_p(r)_{\text{int}(\Gamma)} = \text{Sig}(\text{int}^\circ(r))_\Gamma \circ \phi_{\mathcal{G}',\Gamma} \quad (2.46)$$

that is, the square

$$\begin{array}{ccc} \text{Sig}(\text{int}^{op} \circ \mathcal{G})(\Gamma) & \xrightarrow{\phi_{\mathcal{G},\Gamma}} & D_p(\mathcal{G})(\text{int}(\Gamma)) \\ [\mathbf{2016.11.19.eq2}] \text{int}^\circ(r)_\Gamma \downarrow & & \downarrow D_p(r)_{\text{int}(\Gamma)} \\ \text{Sig}(\text{int}^{op} \circ \mathcal{G}')(\Gamma) & \xrightarrow{\phi_{\mathcal{G}',\Gamma}} & D_p(\mathcal{G}')(\text{int}(\Gamma)) \end{array} \quad (2.47)$$

commutes.

To construct  $\phi_{\mathcal{G},\Gamma}$  we first compute using (2.1)

$$\text{Sig}(\text{int}^{op} \circ \mathcal{G})(\Gamma) = \coprod_{T \in \text{Ob}_1(\Gamma)} \mathcal{G}(\text{int}(T))$$

and using (2.21)

$$D_p(\mathcal{G})(\text{int}(\Gamma)) = \coprod_{F: \text{int}(\Gamma) \rightarrow U} \mathcal{G}((\text{int}(\Gamma); F))$$

and define the function  $\phi_{\mathcal{G},\Gamma}$  by the formula

$$[\mathbf{2016.09.01.eq3}] \phi_{\mathcal{G},\Gamma}((T, g)) = (u_{1,\Gamma}(T), g) \quad (2.48)$$

where the right hand side is defined because of (2.35). The function  $\phi_{\mathcal{G},\Gamma}$  is a bijection as the total function of the bijection  $u_{1,\Gamma}$  and the family of bijections, namely the identity functions.

To prove equality (2.44) we compute using (2.2)

$$\text{Sig}(\text{int} \circ \mathcal{G})(f)(T, g) = (f^*(T), \mathcal{G}(\text{int}(q(f, T)))(\text{int}(T)))$$

and using (2.22)

$$D_p(\mathcal{G})(\text{int}(f))(F, g) = (\text{int}(f) \circ F, \mathcal{G}(Q(\text{int}(f), F))(g))$$

Equality (2.44) follows now from (2.34) and (2.37).

To prove equality (2.46) we compute using (2.3)

$$Sig(int^\circ(r))_\Gamma(T, g) = (T, r_{int(T)}(g))$$

and using (2.23)

$$D_p(r)_{int(\Gamma)}(F, g) = (F, r_{(int(\Gamma); F)}(g))$$

and (2.46) follows from (2.35).

This completes Construction 2.27.

## 2.5 Isomorphisms of presheaves $u_n$ and $\tilde{u}_n$ for $n \geq 2$

In this section we continue to consider a universe category  $(\mathcal{C}, p)$ . For any such  $(\mathcal{C}, p)$  and any  $n \geq 1$ , we construct isomorphisms of presheaves on  $CC(\mathcal{C}, p)$  of the form

$$[2016.11.22.eq1] u_n : \mathcal{O}b_n \rightarrow int^\circ(D_p^{n-1}(Yo(U))) \quad (2.49)$$

and

$$[2016.11.22.eq2] \tilde{u}_n : \widetilde{\mathcal{O}b}_n \rightarrow int^\circ(D_p^{n-1}(Yo(\tilde{U}))) \quad (2.50)$$

where  $D_p^0 = Id_{PreShv(\mathcal{C})}$ , and  $u_1$  and  $\tilde{u}_1$  are the isomorphisms constructed in Section 2.3. We show that

$$[2016.12.02.eq7] \tilde{u}_n \circ int^\circ(D_p^{n-1}(Yo(p))) = \partial \circ \tilde{u}_n \quad (2.51)$$

Let us fix a universe category  $(\mathcal{C}, p)$ .

**Problem 2.28** [2016.11.22.probl] *Let  $n \geq 2$ . To construct an isomorphism of presheaves on  $CC(\mathcal{C}, p)$  of the form (2.49).*

**Construction 2.29** [2016.11.22.constr1] We proceed by induction on  $n$  starting with  $n = 1$ . Observe that  $SD_{p, \mathcal{G}}$  is an isomorphism of the form

$$[2016.11.22.eq3] Sig(int^\circ(\mathcal{G})) \rightarrow int^\circ(D_p(\mathcal{G})) \quad (2.52)$$

The isomorphism  $u_1$  was constructed in Section 2.3. For the successor, define  $u_{n+1}$  as the following composition

$$\begin{array}{ccccc} \mathcal{O}b_{n+1} & \xrightarrow{S\mathcal{O}b_n^{-1}} & Sig(\mathcal{O}b_n) & \xrightarrow{Sig(u_n)} & Sig(int^\circ(D_p^{n-1}(Yo(U)))) & \xrightarrow{SD_{p, D_p^{n-1}(Yo(U))}} \\ & & int^\circ(D_p(D_p^{n-1}(Yo(U)))) & \xlongequal{\quad} & int^\circ(D_p^n(Yo(U))) & \end{array}$$

The isomorphism  $u_{n+1, \Gamma}$  is of the form

$$T \mapsto (ft^n(T), T) \mapsto (ft^n(T), u_{n, ft^n(T)}(T)) \mapsto (u_{1, \Gamma}(ft^n(T)), u_{n, ft^n(T)}(T))$$

where the form of the first map is by (2.8), the second by (2.3) and the third by (2.48). In particular, for  $n = 1$  we get

$$u_{2,\Gamma}(T) = (u_{1,\Gamma}(ft(T)), u_{1,ft(T)}(T))$$

**Problem 2.30** [2016.11.22.prob2] *Let  $n \geq 2$ . To construct an isomorphism of presheaves on  $CC(\mathcal{C}, p)$  of the form (2.50).*

**Construction 2.31** [2016.11.22.constr2] We proceed by induction on  $n$  starting with  $n = 1$ . The isomorphism  $\tilde{u}_1$  was constructed in Section 2.3. For the successor, define  $\tilde{u}_{n+1}$  as the following composition, where we use that  $SD_{p,\mathcal{G}}$  is of the form (2.52),

$$\begin{array}{ccccc} \tilde{\mathcal{O}}b_{n+1} & \xrightarrow{S\tilde{\mathcal{O}}b_n^{-1}} & Sig(\tilde{\mathcal{O}}b_n) & \xrightarrow{Sig(\tilde{u}_n)} & Sig(int^\circ(D_p^{n-1}(Yo(\tilde{U})))) & \xrightarrow{SD_{p,D_p^{n-1}(Yo(\tilde{U}))}} \\ & & & & int^\circ(D_p(D_p^{n-1}(Yo(\tilde{U})))) & \xlongequal{\quad} & int^\circ(D_p^n(Yo(\tilde{U}))) \end{array}$$

The isomorphism  $\tilde{u}_{n+1,\Gamma}$  is of the form

$$o \mapsto (ft^n(\partial(o)), o) \mapsto (ft^n(\partial(o)), \tilde{u}_{n,ft^n(\partial(o))}(o)) \mapsto (u_{1,\Gamma}(ft^n(\partial(o))), \tilde{u}_{n,ft^n(\partial(o))}(o))$$

where the form of the first map is by (2.12), the second by (2.3) and the third by (2.48). In particular, for  $n = 1$  we get

$$\tilde{u}_{2,\Gamma}(o) = (u_{1,\Gamma}(ft(\partial(o))), \tilde{u}_{1,ft(\partial(o))}(o))$$

**Lemma 2.32** [2016.12.02.l3] *For any  $n \geq 1$ , (2.51) holds, that is, the square*

$$\begin{array}{ccc} \tilde{\mathcal{O}}b_n & \xrightarrow{\tilde{u}_n} & int^\circ(D_p^{n-1}(Yo(\tilde{U}))) \\ \partial \downarrow & & \downarrow int^\circ(D_p^{n-1}(Yo(p))) \\ \mathcal{O}b_n & \xrightarrow{u_n} & int^\circ(D_p^{n-1}(Yo(U))) \end{array}$$

*commutes.*

**Proof:** We proceed by induction on  $n$  starting at  $n = 1$ . For  $n = 1$  it is shown in Lemma 2.25.

For the successor of  $n$  we have the diagram

$$\begin{array}{ccccccc} \tilde{\mathcal{O}}b_{n+1} & \xrightarrow{S\tilde{\mathcal{O}}b_n^{-1}} & Sig(\tilde{\mathcal{O}}b_n) & \xrightarrow{Sig(\tilde{u}_n)} & Sig(int^\circ(D_p^n(Yo(\tilde{U})))) & \xrightarrow{SD_p} & int^\circ(D_p(D_p^n(Yo(\tilde{U})))) \\ \partial \downarrow & & Sig(\partial) \downarrow & & Sig(int^\circ(D_p^n(Yo(p)))) \downarrow & & int^\circ(D_p(D_p^n(Yo(p)))) \downarrow \\ \mathcal{O}b_{n+1} & \xrightarrow{S\mathcal{O}b_n^{-1}} & Sig(\mathcal{O}b_n) & \xrightarrow{Sig(u_n)} & Sig(int^\circ(D_p^n(Yo(U)))) & \xrightarrow{SD_p} & int^\circ(D_p(D_p^n(Yo(U)))) \end{array}$$

where the composition of the upper horizontal arrows is  $\tilde{u}_n$  and the composition of the lower horizontal ones is  $u_n$ . To prove the lemma it is sufficient to show that the three squares of the diagram commute.

The commutativity of the left square follows easily from Lemma 2.10. The middle square commutes by the inductive assumption using the fact that  $Sig$  is a functor. The right square commutes because  $SD_p$  is an isomorphism of functors, that is, it is natural in morphisms of presheaves.

## 2.6 The case of a locally cartesian closed $\mathcal{C}$

In this section  $\mathcal{C}$  is a locally cartesian closed category (see Appendix 5.2) with a binary product structure (see Appendix 5.1).

The main construction of this section is Construction 2.35 for Problem ?? that provides, for a universe  $p$  in a category  $\mathcal{C}$  as above, representations for the presheaves  $D_p(Yo(V))$ . As a corollary we provide constructions for Problems 2.36 and ??.

For a morphism  $p : \tilde{U} \rightarrow U$  in  $\mathcal{C}$  and an object  $V$  of  $\mathcal{C}$  let

$$I_p(V) := \underline{Hom}_U((\tilde{U}, p), (U \times V, pr_1))$$

and let

$$pr I_p(V) = p \Delta pr_1 : I_p(V) \rightarrow U$$

be the canonical morphism.

For a morphism  $f : V \rightarrow V'$  let

$$I_p(f) = \underline{Hom}_U((\tilde{U}, p), U \times f)$$

By (5.10),(5.11) and Definition 5.4(3) we have

$$I_p(Id_V) = Id_{I_p(V)}$$

and for  $f' : V' \rightarrow V''$  we have

$$I_p(f \circ f') = I_p(f) \circ I_p(f')$$

which shows that the mappings  $V \mapsto I_p(V)$  and  $f \mapsto I_p(f)$  define a functor from  $\mathcal{C}$  to itself.

The main goal of this section is to construct an isomorphism between the functors

$$V \mapsto D_p(Yo(V))$$

and

$$V \mapsto Yo(I_p(V))$$

from  $\mathcal{C}$  to  $PreShv(\mathcal{C})$ .

Note that  $I_p$  depends on the choice of both the locally cartesian closed and the binary product structures on  $\mathcal{C}$ , but does not depend on the universe structure. On the other hand,

the construction of the functors  $D_p(F)$  requires a universe structure on  $p$  but does not require either the locally cartesian closed or the binary product structure on  $\mathcal{C}$ .

The computations below are required because we have to deal with this fact. In particular, we have to take into the account that for  $F : X \rightarrow U$  the fiber product  $(X, F) \times_U (\tilde{U}, p)$  that we have from the structure of a category with pullbacks on  $\mathcal{C}$  need not be equal to  $(X; F)$  that we have from the universe structure on  $p$ .

Let  $p : \tilde{U} \rightarrow U$  be a universe and  $V$  an object of  $\mathcal{C}$ . We assume that  $\mathcal{C}$  is equipped with a locally cartesian closed and a binary product structures. For  $F : X \rightarrow U$  there is a unique morphism

$$\iota_F : (X; F) \rightarrow (X, F) \times_U (\tilde{U}, p)$$

such that

$$\begin{aligned} \text{[2016.12.02.eq3]} \iota_F \circ pr_1 &= p_F \\ \iota_F \circ pr_2 &= Q(F) \end{aligned} \tag{2.53}$$

which is a particular case of the morphisms  $\iota$  of Lemma 5.3.

The evaluation morphism in the case of  $I_p(V)$  is a morphism in  $\mathcal{C}/U$  of the form

$$evI_p : (I_p(V), prI_p(V)) \times_U (U \times V, pr_1) \rightarrow (U \times V, pr_1)$$

Define a morphism

$$st_p(V) : (I_p(V); prI_p(V)) \rightarrow V$$

as the composition:

$$\text{[2016.12.02.eq2]} st_p(V) := \iota_{prI_p(V)} \circ evI_p(V) \circ pr_2 \tag{2.54}$$

We will need to use some properties of these morphisms.

**Lemma 2.33** *[2015.04.14.l2a] Let  $f : V \rightarrow V'$  be a morphism, then one has*

$$Q(I_p(f), prI_p(V')) \circ st_p(V') = st_p(V) \circ f$$

**Proof:** Let  $pr = prI_p(V)$ ,  $pr' = prI_p(V')$ ,  $\iota = \iota_{pr}$ ,  $\iota' = \iota_{pr'}$ ,  $ev = evI_p(V)$  and  $ev' = evI_p(V')$ . Then we have to verify that the outer square of the following diagram commutes:

$$\begin{array}{ccccccc} (I_p(V); pr) & \xrightarrow{\iota} & (I_p(V), pr) \times_U (\tilde{U}, p) & \xrightarrow{ev} & U \times V & \xrightarrow{pr_2} & V \\ Q(I_p(f), pr') \downarrow & & I_p(f) \times Id_{\tilde{U}} \downarrow & & Id_U \times f \downarrow & & \downarrow f \\ (I_p(V'); pr') & \xrightarrow{\iota'} & (I_p(V'), pr') \times_U (\tilde{U}, p) & \xrightarrow{ev'} & U \times V' & \xrightarrow{pr_2} & V' \end{array}$$

The commutativity of the left square is a particular case of Lemma 5.3. The commutativity of the right square is an immediate corollary of the definition of  $Id_U \times f$ . The commutativity of the middle square is a particular case of (5.7).



**Problem 2.34** [2015.03.29.probl1] Let  $\mathcal{C}$  be a locally cartesian closed category with a binary product structure and  $p$  a universe in  $\mathcal{C}$ . To construct, for all  $V \in \mathcal{C}$ , isomorphisms of presheaves

$$\eta_V : D_p(Yo(V)) \rightarrow Yo(I_p(V))$$

that are natural in  $V$ , i.e., such that for all  $f : V \rightarrow V'$  the square

$$\begin{array}{ccc} D_p(Yo(V)) & \xrightarrow{D_p(Yo(f))} & D_p(Yo(V')) \\ \eta_V \downarrow & & \downarrow \eta_{V'} \\ Yo(I_p(V)) & \xrightarrow{Yo(I_p(f))} & Yo(I_p(V')) \end{array}$$

commutes.

**Construction 2.35** [2015.03.29.constr1] We need to construct bijections

$$\eta_{V,X} : D_p(Yo(V))(X) \rightarrow Mor_{\mathcal{C}}(X, I_p(V))$$

such that for all  $f : V \rightarrow V'$ ,  $X \in \mathcal{C}$  and  $d \in D_p(Yo(V))(X)$  one has

$$[2016.09.11.eq1] \eta_{V',X}(d) \circ I_p(f) = \eta_{V',X}(D_p(Yo(f))_X(d)) \quad (2.55)$$

and for any  $f : X' \rightarrow X$  and  $d \in D_p(Yo(V))(X)$  one has

$$[2016.09.11.eq2] f \circ \eta_{V,X}(d) = \eta_{V,X'}(D_p(Yo(V))(f)(d)) \quad (2.56)$$

We will construct bijections

$$\eta_{V,X}^! : Mor(X, I_p(V)) \rightarrow D_p(Yo(V))(X)$$

such that for all  $g : X \rightarrow I_p(V)$  one has:

1. for all  $f : V \rightarrow V'$  one has

$$[2016.09.11.eq3] D_p(Yo(f))_X(\eta^!(g)) = \eta^!(g \circ I_p(f)) \quad (2.57)$$

2. for all  $f : X' \rightarrow X$  one has

$$[2016.09.11.eq4] D_p(Yo(V))(f)(\eta^!(g)) = \eta^!(f \circ g) \quad (2.58)$$

and then define  $\eta_{V,X}$  as the inverse to  $\eta_{V,X}^!$ . One proves easily that (2.55) implies (2.57) and (2.56) implies (2.58).

By (2.26) we have

$$D_p(Yo(V))(X) = \coprod_{F: X \rightarrow U} Mor_{\mathcal{C}}((X; F), V)$$

For  $g : X \rightarrow I_p(V)$  we set

$$[2016.12.02.eq5] \eta_{V,X}^!(g) := (g \circ pr I_p(V), Q(g, pr I_p(V)) \circ st_p(V)) \quad (2.59)$$

as can be seen on the diagram

$$\begin{array}{ccccc}
& & & & V \\
& & & & \uparrow \text{st}_p(V) \\
(X; g \circ \text{pr}I_p(V)) & \xrightarrow{Q(g, \text{pr}I_p(V))} & (I_p(V); \text{pr}I_p(V)) & \xrightarrow{Q(\text{pr}I_p(V))} & \tilde{U} \\
\downarrow & & \downarrow & & \downarrow p \\
X & \xrightarrow{g} & I_p(V) & \xrightarrow{\text{pr}I_p(V)} & U
\end{array}$$

To see that this is a bijection observe first that it equals to the composition

$$Mor(X, I_p(V)) \rightarrow \amalg_{F: X \rightarrow U} Mor_U((X, F), (I_p(V), \text{pr}I_p(V))) \rightarrow \amalg_{F: X \rightarrow U} Mor((X; F), V)$$

where the first function is given by the formula  $g \mapsto (g \circ \text{pr}I_p(V), g)$  and the second is the sum over all  $F : X \rightarrow U$  of functions  $g \mapsto Q(g, \text{pr}I_p(V)) \circ \text{st}_p(V)$ .

The first function is a function of the form  $A \rightarrow \amalg_{b \in B} h^{-1}(b)$ , which is defined and is a bijection for any function of sets  $h : A \rightarrow B$ . It remains to show that the second one is a bijection for every  $F$ .

By definition of the  $\underline{Hom}_U$  structure we know that for each  $F$  the function

$$adj : Mor_U((X, F), (I_p(V), \text{pr}I_p(V))) \rightarrow Mor_U((X, F) \times_U (\tilde{U}, p), (U \times V, \text{pr}_1))$$

given by  $g \mapsto (g \times_U Id_{(\tilde{U}, p)}) \circ evI_p(V)$  is a bijection.

By definition of the binary product, the function of post-composition with  $\text{pr}_2$ ,

$$Mor_U((X, F) \times_U (\tilde{U}, p), (U \times V, \text{pr}_1)) \rightarrow Mor((X, F) \times_U (\tilde{U}, p), V)$$

is a bijection. By Lemma 5.2,  $\iota_F$  is an isomorphism and therefore the pre-composition with it is a bijection. Now we have two functions

$$Mor_U((X, F), (I_p(V), \text{pr}I_p(V))) \rightarrow Mor((X; F), V)$$

given by  $g \mapsto \iota_F \circ (g \times_U Id_{\tilde{U}}) \circ evI_p(V) \circ \text{pr}_2$  and  $g \mapsto Q(g, \text{pr}I_p(V)) \circ \text{st}_p(V)$  of which the first one is the bijection. It remains to show that these functions are equal. In view of (2.54) it is sufficient to show that

$$\iota_F \circ (g \times_U Id_{\tilde{U}}) = Q(g, \text{pr}I_p(V)) \circ \iota_{\text{pr}I_p(V)}$$

To do it we have to show that the compositions of the left and right hand sides with  $\text{pr}_1$  (to  $I_p(V)$ ) and  $\text{pr}_2$  (to  $\tilde{U}$ ) are equal.

For  $\text{pr}_1$  we have

$$\begin{aligned}
\iota_F \circ (g \times_U Id_{\tilde{U}}) \circ \text{pr}_1 &= \iota_F \circ \text{pr}_1 \circ g = p_F \circ g \\
Q(g, \text{pr}I_p(V)) \circ \iota_{\text{pr}I_p(V)} \circ \text{pr}_1 &= Q(g, \text{pr}I_p(V)) \circ p_{\text{pr}I_p(V)} = p_{g \circ \text{pr}I_p(V)} \circ g = p_F \circ g
\end{aligned}$$

where we used the defining equations (2.6) of  $\iota$ , the definition (2.17) of  $Q(-, -)$  and the fact that  $g$  is a morphism over  $U$ .

For  $pr_2$  we have

$$\iota_F \circ (g \times_U Id_{\tilde{U}}) \circ pr_2 = \iota_F \circ pr_2 \circ Id_{\tilde{U}} = \iota_F \circ pr_2 = Q(F)$$

$$Q(g, pr I_p(V)) \circ \iota_{pr I_p(V)} \circ pr_2 = Q(g, pr I_p(V)) \circ Q(pr I_p(V)) = Q(g \circ pr I_p(V)) = Q(F)$$

where we used the defining equations (2.6) of  $\iota$ , (2.18) and the fact that  $g$  is a morphism over  $U$ .

We now have to check the behavior of  $\eta^!$  with respect to morphisms in  $V$  (equality (2.57)) and  $X$  (equality (2.58)).

Let  $pr = pr I_p(V)$  and  $pr' = pr I_p(V')$ . Let  $g : X \rightarrow I_p(V)$  be as above. For  $f : V' \rightarrow V$  we have

$$D_p(Yo(f))_X(\eta^!(g)) = D_p(Yo(f))_X(g \circ pr, Q(g, pr) \circ st_p(V)) = (g \circ pr, Q(g, pr) \circ st_p(V) \circ f)$$

where the first equality is by (2.59) and the second by (2.28) and

$$\eta^!(g \circ I_p(f)) = (g \circ I_p(f) \circ pr', Q(g \circ I_p(f), pr') \circ st_p(V'))$$

where the equality is by (2.59). We have  $pr = I_p(f) \circ pr'$  because  $I_p(f)$  is a morphism over  $U$ . It remains to check that

$$Q(g, pr) \circ st_p(V) \circ f = Q(g \circ I_p(f), pr') \circ st_p(V')$$

By [?, Lemma 2.5] we have

$$Q(g \circ I_p(f), pr') = Q(g, pr) \circ Q(I_p(f), pr')$$

and the remaining equality

$$Q(g, pr) \circ st_p(V) \circ f = Q(g, pr) \circ Q(I_p(f), pr') \circ st_p(V')$$

follows from Lemma 2.33.

Consider now  $f : X' \rightarrow X$ . Then

$$\begin{aligned} D_p(Yo(V))(f)(\eta^!(g)) &= D_p(Yo(V))(f)(g \circ pr, Q(g, pr) \circ st_p(V)) = \\ &= (f \circ g \circ pr, Q(f, g \circ pr) \circ Q(g, pr) \circ st_p(V)) \end{aligned}$$

and

$$\eta^!(f \circ g) = (f \circ g \circ pr, Q(f \circ g, pr) \circ st_p(V))$$

where we used (2.59) and (2.27) and the required equality follows from [?, Lemma 2.5].

**Problem 2.36** [2016.12.02.probl] *For a locally cartesian closed category  $\mathcal{C}$  with a binary product structure and a universe  $p$  in  $\mathcal{C}$  to construct, for all  $n \geq 0$  and  $V \in \mathcal{C}$ , isomorphisms of presheaves*

$$\eta_{n,V} : D_p^n(Yo(V)) \rightarrow Yo(I_p^n(V))$$

that are natural in  $V$ .

**Construction 2.37** [2016.12.02.constr1] Proceed by induction on  $n$  starting with  $n = 0$ . By our convention,  $D_p^0 = Id_{PreShv(\mathcal{C})}$  and  $I_p^0 = Id_{\mathcal{C}}$ . We set  $\eta_{0,V} = Id_{Yo(V)}$ . For the successor we define  $\eta_{n+1,V}$  as the composition

$$\begin{aligned} D_p^{n+1}(Yo(V)) = \\ D_p(D_p^n(Yo(V))) \xrightarrow{D_p(\eta_{n,V})} D_p(Yo(I_p^n(V))) \xrightarrow{\eta_{n,I_p^n(V)}} Yo(I_p(I_p^n(V))) = \\ Yo(I_p^{n+1}(V)) \end{aligned}$$

The naturality in  $V$  is easily proved by induction.

**Problem 2.38** [2015.03.17.prob3] For  $\mathcal{C}$  as above, a universe  $p : \tilde{U} \rightarrow U$  in  $\mathcal{C}$  and  $n \geq 1$  to construct isomorphisms of presheaves

$$\mu_n : \mathcal{O}b_n \rightarrow \text{int}^\circ(Yo(I_p^{n-1}(U)))$$

and

$$\tilde{\mu}_n : \tilde{\mathcal{O}}b_n \rightarrow \text{int}^\circ(Yo(I_p^{n-1}(\tilde{U})))$$

such that the square

$$\begin{array}{ccc} \tilde{\mathcal{O}}b_n & \xrightarrow{\tilde{\mu}_n} & \text{int}^\circ(Yo(I_p^{n-1}(\tilde{U}))) \\ \text{[2016.12.04.eq2]} \downarrow \partial & & \downarrow \text{int}^\circ(Yo(I_p^{n-1}(p))) \\ \mathcal{O}b_n & \xrightarrow{\mu_n} & \text{int}^\circ(Yo(I_p^{n-1}(U))) \end{array} \quad (2.60)$$

commutes.

**Construction 2.39** [2015.03.17.constr2] Compose isomorphism  $u_n$  of Construction 2.29 (resp. isomorphism  $\tilde{u}_n$  of Construction 2.31) with the isomorphism  $\text{int}^\circ(\eta_{n-1,U})$  (resp.  $\text{int}^\circ(\eta_{n-1,\tilde{U}})$ ) of Construction 2.37.

To prove the commutativity of (2.60) consider the diagram

$$\begin{array}{ccc} \tilde{\mathcal{O}}b_n & \xrightarrow{\tilde{u}_n} & \text{int}^\circ(D_p^{n-1}(Yo(\tilde{U}))) & \xrightarrow{\text{int}^\circ(\eta_{n-1,\tilde{U}})} & \text{int}^\circ(Yo(I_p^{n-1}(\tilde{U}))) \\ \partial \downarrow & & \text{int}^\circ(D_p^{n-1}(Yo(p))) \downarrow & & \downarrow \text{int}^\circ(Yo(I_p^{n-1}(p))) \\ \mathcal{O}b_n & \xrightarrow{u_n} & \text{int}^\circ(D_p^{n-1}(Yo(U))) & \xrightarrow{\text{int}^\circ(\eta_{n-1,U})} & \text{int}^\circ(Yo(I_p^{n-1}(U))) \end{array}$$

The composition of the upper arrows is  $\tilde{\mu}_n$  and the composition of the lower ones is  $\mu_n$ . It remains to show that the two squares commute. The left square commutes by Lemma 2.32. The right square commutes because  $\text{int}^\circ$  is a functor and  $\eta_{n-1,V}$  is natural in  $V$ .

**Remark 2.40** [2016.04.23.rem1] In [?] generalized polynomial functors are defined as functors isomorphic to functors of the form  $I_p$ .

### 3 $P$ -structures on universes and $(\Pi, \lambda)$ -structures

#### 3.1 Construction of $(\Pi, \lambda)$ -structures on the C-systems $CC(\mathcal{C}, p)$

We will show now how to construct  $(\Pi, \lambda)$ -structures on C-systems of the form  $CC(\mathcal{C}, p)$  for locally cartesian closed universe categories  $(\mathcal{C}, p)$  with a binary product structure.

That construction for Problem 3.6, without the part that concerns the bijection, exists was originally stated in [?, Proposition 2] with a sketch of a proof given in the 2009 version of [?].

Let us recall the following definition from [?]:

**Definition 3.1** [2015.03.09.def1] *Let  $CC$  be a C-system. A pre- $(\Pi, \lambda)$ -structure on  $CC$  is a pair of morphisms of presheaves*

$$\begin{aligned}\Pi &: \mathcal{O}b_2 \rightarrow \mathcal{O}b_1 \\ \lambda &: \widetilde{\mathcal{O}}b_2 \rightarrow \widetilde{\mathcal{O}}b_1\end{aligned}$$

such that the square

$$\begin{array}{ccc} \widetilde{\mathcal{O}}b_2 & \xrightarrow{\lambda} & \widetilde{\mathcal{O}}b_1 \\ \text{[2015.03.09.eq1]}_{\partial} \downarrow & & \downarrow_{\partial} \\ \mathcal{O}b_2 & \xrightarrow{\Pi} & \mathcal{O}b_1 \end{array} \quad (3.1)$$

commutes.

A pre- $(\Pi, \lambda)$ -structure is called a  $(\Pi, \lambda)$ -structure if the square (3.1) is a pullback.

**Definition 3.2** [2015.03.29.def1] *Let  $\mathcal{C}$  be a locally cartesian closed category with a binary product structure and  $p : \widetilde{U} \rightarrow U$  a universe in  $\mathcal{C}$ . A  $P$ -structure on  $p$  is a pair of morphisms*

$$\begin{aligned}\widetilde{P} &: I_p(\widetilde{U}) \rightarrow \widetilde{U} \\ P &: I_p(U) \rightarrow U\end{aligned}$$

such that the square

$$\begin{array}{ccc} I_p(\widetilde{U}) & \xrightarrow{\widetilde{P}} & \widetilde{U} \\ \text{[2009.prod.square]}_{\mathfrak{h}(p)} \downarrow & & \downarrow_p \\ I_p(U) & \xrightarrow{P} & U \end{array} \quad (3.2)$$

is a pullback.

**Problem 3.3** [2015.03.17.prob0] *Let  $(\mathcal{C}, p)$  be a locally cartesian closed universe category with a binary product structure. Let  $(P, \widetilde{P})$  be a  $P$ -structure on  $p$ . To construct a  $(\Pi, \lambda)$ -structure on  $CC(\mathcal{C}, p)$ .*

**Construction 3.4** [2015.03.17.constr3] Consider the diagram:

$$\begin{array}{ccccccc}
\widetilde{\mathcal{O}}b_2 & \xrightarrow{\widetilde{\mu}_2} & \text{int}^\circ(Yo(I_p(\widetilde{U}))) & \xrightarrow{\text{int}^\circ(Yo(\widetilde{P}))} & \text{int}^\circ(Yo(\widetilde{U})) & \xrightarrow{\mu_1^{-1}} & \widetilde{\mathcal{O}}b_1 \\
\text{[2016.12.09.eq1]} \downarrow \partial & & \downarrow \text{int}^\circ(Yo(I_p(p))) & & \downarrow \text{int}^\circ(Yo(p)) & & \downarrow \partial \\
\mathcal{O}b_2 & \xrightarrow{\mu_2} & \text{int}^\circ(Yo(I_p(U))) & \xrightarrow{\text{int}^\circ(Yo(P))} & \text{int}^\circ(Yo(U)) & \xrightarrow{\mu_1^{-1}} & \mathcal{O}b_1
\end{array} \quad (3.3)$$

Since the squares (2.60) commute, the square (3.2) commutes and both  $Yo$  and  $\text{int}^\circ$  are functors, the external square of this diagram commutes and therefore defines a pre- $(\Pi, \lambda)$  structure. We will show that it is a  $(\Pi, \lambda)$ -structure, that is, that this square is a pullback.

Horizontal composition of pullbacks is a pullback. The left hand side square is a pullback because it is a commutative square with two parallel sides being isomorphisms. The right hand side square is a pullback for the same reason.

It remains to show that the middle square is pullback. This square is obtained by applying first the functor  $Yo$  and then the functor  $\text{int}^\circ$  to the pullback square (3.2).

Our claim follows now from two facts:

1. the Yoneda functor  $Yo : \mathcal{C} \rightarrow \text{PreShv}(\mathcal{C})$  takes pullbacks to pullbacks,
2. for any functor  $F : \mathcal{C}' \rightarrow \mathcal{C}$ , the functor

$$F^\circ : \text{PreShv}(\mathcal{C}) \rightarrow \text{PreShv}(\mathcal{C}')$$

of pre-composition with  $F^{op}$ , takes pullbacks to pullbacks.

We assume that these two facts are known.

We conclude that for a  $P$ -structure  $(P, \widetilde{P})$  the pair of morphisms

$$\begin{array}{l}
\text{[2016.12.09.eq3]} \quad \Pi = \widetilde{\mu}_2 \circ \text{int}^\circ(Yo(\widetilde{P})) \circ \widetilde{\mu}_1^{-1} \\
\lambda = \mu_2 \circ \text{int}^\circ(Yo(P)) \circ \mu_1^{-1}
\end{array} \quad (3.4)$$

is a  $(\Pi, \lambda)$ -structure on  $CC(\mathcal{C}, p)$ .

There is an important class of cases when the function from  $P$ -structures on  $p$  to  $(\Pi, \lambda)$ -structures on  $CC(\mathcal{C}, p)$  is a bijection.

**Lemma 3.5** [2016.09.09.11] *Let  $(\mathcal{C}, p)$  be a universe category such that the functor*

$$Yo \circ \text{int}^\circ : \mathcal{C} \rightarrow \text{PreShv}(CC(\mathcal{C}, p))$$

*is fully faithful. Then the function from the  $P$ -structures on  $p$  to the  $(\Pi, \lambda)$ -structures on  $CC(\mathcal{C}, p)$  defined by Construction 3.4 is a bijection.*

**Proof:** Let  $\tilde{\phi}$  be the inverse to  $(Yo \circ \text{int}^\circ)_{I_p(\tilde{U}), \tilde{U}}$  and  $\phi$  be the inverse to  $(Yo \circ \text{int}^\circ)_{I_p(U), U}$ . Given a  $(\Pi, \lambda)$ -structure  $(\Pi, \lambda)$  let

$$\begin{aligned} \text{[2016.09.09.eq1]} \quad \tilde{P} &= \tilde{\phi}(\tilde{\mu}_2^{-1} \circ \Pi \circ \tilde{\mu}_1) \\ P &= \phi(\mu_2^{-1} \circ \lambda \circ \mu_1) \end{aligned} \quad (3.5)$$

Then  $\tilde{P} : I_p(\tilde{U}) \rightarrow \tilde{U}$  and  $P : I_p(U) \rightarrow U$ . Let  $S$  be the square that  $\tilde{P}$  and  $P$  form with  $I_p(p)$  and  $p$ . The square verifies that  $(Yo \circ \text{int}^\circ)(S)$  is of the form

$$\begin{array}{ccc} \text{int}^\circ(Yo(I_p(\tilde{U}))) & \xrightarrow{\tilde{\mu}_2^{-1} \circ \Pi \circ \tilde{\mu}_1} & \text{int}^\circ(Yo(\tilde{U})) \\ \text{int}^\circ(Yo(I_p(p))) \downarrow & & \downarrow \text{int}^\circ(Yo(p)) \\ \text{int}^\circ(Yo(I_p(U))) & \xrightarrow{\mu_2^{-1} \circ \lambda \circ \mu_1} & \text{int}^\circ(Yo(U)) \end{array}$$

Since the left and right squares of (3.3) commute, the square  $(Yo \circ \text{int}^\circ)(S)$  is isomorphic to the original square formed by  $\Pi$  and  $\lambda$  and as a square isomorphic to a pullback is a pullback.

The functor  $Yo \circ \text{int}^\circ$  is assumed to be fully faithful and if the image of a square under a fully faithful functor is a pullback then the square itself is a pullback. We conclude that formulas (3.5) define a function from  $(\Pi, \lambda)$ -structures to  $P$ -structures.

It remains to verify that this function is inverse on both sides to the function of Construction 3.4, that is, that formulas (3.4) and (3.5) define mutually inverse functions on pairs of morphisms. This is straightforward by rewriting. The lemma is proved.

**Problem 3.6** [2016.12.09.prob2] *Let  $(\mathcal{C}, p)$  be a universe category.*

*To construct a function from the set of  $P$ -structures on  $p$  to the set of structures of products of families of types on  $CC(\mathcal{C}, p)$ .*

*To show that if the functor  $Yo \circ \text{int}^\circ$  is fully faithful than this function is a bijection.*

**Construction 3.7** [2016.12.09.constr2] The required function is the composition of the function of Construction 3.4 with the construction for [?, Problem 4.5] described in that paper.

## 4 Functoriality

### 4.1 Universe category functors and the $D_p$ construction

Let  $(\mathcal{C}, p, pt)$  and  $(\mathcal{C}', p', pt')$  be two universe categories. Recall from [?] the following definition.

**Definition 4.1** [2016.12.09.def1] *A universe category functor from  $(\mathcal{C}, p, pt)$  to  $(\mathcal{C}', p', pt')$  is a triple  $\Phi = (\Phi, \phi, \tilde{\phi})$  where  $\Phi$  is a functor  $\mathcal{C} \rightarrow \mathcal{C}'$  and  $\phi : \Phi(U) \rightarrow U'$ ,  $\tilde{\phi} : \Phi(\tilde{U}) \rightarrow \tilde{U}'$  are two morphisms such that one has:*

1.  $\Phi$  takes the pt to a final object,
2.  $\Phi$  takes the canonical pullbacks based on  $p$  to pullbacks,
3. the square

$$\begin{array}{ccc}
\Phi(\tilde{U}) & \xrightarrow{\tilde{\phi}} & \tilde{U}' \\
\text{[2015.03.21.sq1]}_{(p)} \downarrow & & \downarrow_{p'} \\
\Phi(U) & \xrightarrow{\phi} & U'
\end{array} \tag{4.1}$$

is a pullback.

**Problem 4.2** [2016.12.14.prob1] Let  $\Phi = (\Phi, \phi, \tilde{\phi})$  be a universe category functor  $(\mathcal{C}, p) \rightarrow (\mathcal{C}', p')$ . To construct a functor morphism

$$\text{[2016.12.14.eq3]} \Phi D : \Phi^\circ \circ D_p \rightarrow D_{p'} \circ \Phi^\circ \tag{4.2}$$

**Construction 4.3** [2016.12.14.constr1] Both the left and the right hand side of (4.2) are functors of the form

$$PreShv(\mathcal{C}') \rightarrow PreShv(\mathcal{C})$$

Therefore, we need, for any presheaf  $\mathcal{G}'$  on  $\mathcal{C}'$  and any  $X \in \mathcal{C}$ , to construct a function

$$\text{[2016.12.14.eq4]} \Phi D_{\mathcal{G}', X} : D_p(\Phi^\circ(\mathcal{G}'))(X) \rightarrow \Phi^\circ(D_{p'}(\mathcal{G}'))(X) \tag{4.3}$$

and to prove that

1. the family  $\Phi D_{\mathcal{G}', -}$  is a morphism of presheaves, that is, for any  $a : X \rightarrow Y$  in  $\mathcal{C}$ , the square

$$\begin{array}{ccc}
D_p(\Phi^\circ(\mathcal{G}'))(Y) & \xrightarrow{\Phi D_{\mathcal{G}', Y}} & \Phi^\circ(D_{p'}(\mathcal{G}'))(Y) \\
\text{[2016.12.16.eq2]}_{\Phi^\circ(\mathcal{G}')(a)} \downarrow & & \downarrow_{\Phi^\circ(D_{p'}(\mathcal{G}'))(a)} \\
D_p(\Phi^\circ(\mathcal{G}'))(X) & \xrightarrow{\Phi D_{\mathcal{G}', X}} & \Phi^\circ(D_{p'}(\mathcal{G}'))(X)
\end{array} \tag{4.4}$$

commutes,

2.  $\Phi D$  is a natural transformation of functors to presheaves, that is, for any  $f' : \mathcal{F}' \rightarrow \mathcal{G}'$  and any  $X \in \mathcal{C}$  the square

$$\begin{array}{ccc}
D_p(\Phi^\circ(\mathcal{F}'))(X) & \xrightarrow{\Phi D_{\mathcal{F}', X}} & \Phi^\circ(D_{p'}(\mathcal{F}'))(X) \\
\text{[2016.12.16.eq3]}_{\Phi^\circ(f')_X} \downarrow & & \downarrow_{\Phi^\circ(D_{p'}(f'))_X} \\
D_p(\Phi^\circ(\mathcal{G}'))(X) & \xrightarrow{\Phi D_{\mathcal{G}', X}} & \Phi^\circ(D_{p'}(\mathcal{G}'))(X)
\end{array} \tag{4.5}$$

commutes.



Computing the left and right hand side of (4.3) we see that  $\Phi D_{\mathcal{G}', X}$  should be a function of the form

$$\coprod_{F: X \rightarrow U} \mathcal{G}'(\Phi((X; F))) \rightarrow \coprod_{F': \Phi(X) \rightarrow U'} \mathcal{G}'((\Phi(X); F'))$$

Let  $F : X \rightarrow U$ . Consider  $(\Phi(X); \Phi(F) \circ \phi)$ . Since (4.1) is a pullback there is a unique morphism  $q$  such that  $q \circ \tilde{\phi} = Q(\Phi(F) \circ \phi)$  and  $q \circ \Phi(p) = p_{\Phi(X), \Phi(F) \circ \phi} \circ \Phi(F)$ . Then the external square in the diagram

$$\begin{array}{ccccc} (\Phi(X); \Phi(F) \circ \phi) & \xrightarrow{q} & \Phi(\tilde{U}) & \xrightarrow{\tilde{\phi}} & \tilde{U}' \\ \downarrow p_{\Phi(X), \Phi(F) \circ \phi} & & \Phi(p) \downarrow & & \downarrow p' \\ \Phi(X) & \xrightarrow{\Phi(F)} & \Phi(U) & \xrightarrow{\phi} & U' \end{array}$$

is a pullback and since the right hand side square is a pullback, the left hand side square is a pullback as well. Together with the fact that  $\Phi$  takes pullback squares based on  $p$  to pullback squares this implies that we obtain two pullbacks based on  $\Phi(F)$  and  $\Phi(p)$ .

By Lemma 5.8 and Lemma 5.2 we have a unique morphism, which is an isomorphism,

$$\iota : (\Phi(X); \Phi(F) \circ \phi) \rightarrow \Phi(X; F)$$

such that

$$[\mathbf{2015.04.08.eq1}] \iota \circ \Phi(p_{X, F}) = p_{\Phi(X), \Phi(F) \circ \phi} \tag{4.6}$$

$$[\mathbf{2015.04.08.eq2}] \iota \circ \Phi(Q(F)) \circ \tilde{\phi} = Q(\Phi(F) \circ \phi) \tag{4.7}$$

and we define:

$$[\mathbf{2016.12.16.eq4}] \Phi D_{\mathcal{G}', X}(F, \gamma') = (\Phi(F) \circ \phi, \mathcal{G}'(\iota)(\gamma')) \tag{4.8}$$

To prove that (5.13) commutes let

$$(F : Y \rightarrow U, \gamma' \in \mathcal{G}'(\Phi((Y; F)))) \in D_p(\Phi^\circ(\mathcal{G}'))(Y)$$

Then one path in the square gives us

$$\begin{aligned} & (\Phi^\circ(D_{p'}(\mathcal{G}'))(a))(\Phi D_{\mathcal{F}', X}((F, \gamma'))) = \\ & (\Phi^\circ(D_{p'}(\mathcal{G}'))(a))((\Phi(F) \circ \phi, \mathcal{G}'(\iota)(\gamma'))) = D_{p'}(\mathcal{G}'))(\Phi(a))((\Phi(F) \circ \phi, \mathcal{G}'(\iota)(\gamma'))) = \\ & (\Phi(a) \circ \Phi(F) \circ \phi, \mathcal{G}'(Q(\Phi(a), \Phi(F) \circ \phi))(\mathcal{G}'(\iota)(\gamma'))) = \\ & (\Phi(a \circ F) \circ \phi, \mathcal{G}'(Q(\Phi(a), \Phi(F) \circ \phi) \circ \iota)(\gamma')) \end{aligned}$$

where the first equality is by (4.8), the second by the definition of  $\Phi^\circ$ , the third by (2.22) and the fourth by the composition axiom of  $\Phi$  and  $\mathcal{G}'$ .

The other path gives us

$$\Phi D_{\mathcal{G}', X}(D_p(\Phi^\circ(\mathcal{G}'))(a))((F, \gamma')) =$$

$$\begin{aligned}
\Phi D_{\mathcal{G}', X}((a \circ F, \Phi^\circ(\mathcal{G}')(\mathcal{Q}(a, F))(\gamma'))) &= \Phi D_{\mathcal{G}', X}((a \circ F, \mathcal{G}'(\Phi(\mathcal{Q}(a, F)))(\gamma'))) = \\
&(\Phi(a \circ F) \circ \phi, \mathcal{G}'(\iota)(\mathcal{G}'(\Phi(\mathcal{Q}(a, F)))(\gamma'))) = \\
&(\Phi(a \circ F) \circ \phi, \mathcal{G}'(\iota \circ \Phi(\mathcal{Q}(a, F)))(\gamma'))
\end{aligned}$$

where the first equality is by (2.22), the second by the definition of  $\Phi^\circ$ , the third by (4.8) and the fourth by the composition axiom of  $\mathcal{G}'$ .

It remains to show that

$$[\mathbf{2016.12.16.eq7}] Q(\Phi(a), \Phi(F) \circ \phi) \circ \iota = \iota \circ \Phi(\mathcal{Q}(a, F)) \quad (4.9)$$

We have four pullbacks

$$\begin{array}{ccc}
(\Phi(X); \Phi(a \circ F) \circ \phi) & \xrightarrow{Q(\Phi(a \circ F) \circ \phi)} & \tilde{U}' & & (\Phi(Y); \Phi(F) \circ \phi) & \xrightarrow{Q(\Phi(F) \circ \phi)} & \tilde{U}' \\
p_{\Phi(X), \Phi(a \circ F) \circ \phi} \downarrow & & \downarrow p' & & p_{\Phi(Y), \Phi(F) \circ \phi} \downarrow & & \downarrow p' \\
\Phi(X) & \xrightarrow{\Phi(a \circ F) \circ \phi} & U' & & \Phi(Y) & \xrightarrow{\Phi(F) \circ \phi} & U'
\end{array}$$

and

$$\begin{array}{ccc}
\Phi((X; a \circ F)) & \xrightarrow{\Phi(Q(a \circ F)) \circ \tilde{\phi}} & \tilde{U}' & & \Phi((Y; F)) & \xrightarrow{\Phi(Q(F)) \circ \tilde{\phi}} & \tilde{U}' \\
\Phi(p_{X, a \circ F}) \downarrow & & \downarrow p' & & \Phi(p_{Y, F}) \downarrow & & \downarrow p' \\
\Phi(X) & \xrightarrow{\Phi(a \circ F) \circ \phi} & U' & & \Phi(Y) & \xrightarrow{\Phi(F) \circ \phi} & U'
\end{array}$$

and a morphism  $\Phi(a) : \Phi(X) \rightarrow \Phi(Y)$  such that  $\Phi(a \circ F) \circ \phi = \Phi(a) \circ \Phi(F) \circ \phi$ . Applying to these pullbacks Lemma 5.8 and then applying Lemma 5.3 we obtain a commutative square

$$\begin{array}{ccc}
(\Phi(X); \Phi(a \circ F) \circ \phi) & \xrightarrow{c_1(\Phi(a), Id_{\tilde{U}'})} & (\Phi(Y); \Phi(F) \circ \phi) \\
\downarrow \iota & & \downarrow \iota \\
\Phi((X; a \circ F)) & \xrightarrow{c_2(\Phi(a), Id_{\tilde{U}'})} & \Phi((Y; F))
\end{array}$$

To prove (4.9) it remains to show that

$$[\mathbf{2016.12.16.eq5}] c_1(\Phi(a), Id_{\tilde{U}'}) = Q(\Phi(a), \Phi(F) \circ \phi) \quad (4.10)$$

and

$$[\mathbf{2016.12.16.eq6}] c_2(\Phi(a), Id_{\tilde{U}'}) = \Phi(Q(a, F)) \quad (4.11)$$

In view of the definition of the morphisms  $c_1, c_2$  given in Lemma 5.3 to prove (4.10) we need to show that

$$\begin{aligned}
Q(\Phi(a), \Phi(F) \circ \phi) \circ p_{\Phi(Y), \Phi(F) \circ \phi} &= p_{\Phi(X), \Phi(a \circ F) \circ \phi} \circ \Phi(a) \\
Q(\Phi(a), \Phi(F) \circ \phi) \circ Q(\Phi(F) \circ \phi) &= Q(\Phi(a \circ F) \circ \phi)
\end{aligned}$$

The first equality follows from (2.17). The second equality follows from (2.18). In both cases we need also to use that  $\Phi(a \circ F) = \Phi(a) \circ \Phi(F)$ .

To prove (4.11) we need to show that

$$\Phi(Q(a, F)) \circ \Phi(p_{Y, F}) = \Phi(p_{X, a \circ F}) \circ \Phi(a)$$

$$\Phi(Q(a, F)) \circ \Phi(Q(F)) \circ \tilde{\phi} = \Phi(Q(a \circ F)) \circ \tilde{\phi}$$

The first equality again follows from (2.17) and the composition axiom for  $\Phi$  and the second equality follows from (2.18) and the composition axiom for  $\Phi$ . This completes the proof of commutativity of (5.13).

To prove that (4.5) commutes let

$$(F : X \rightarrow U, \beta' \in \mathcal{F}'(\Phi((X; F)))) \in D_p(\Phi^\circ(\mathcal{F}'))(X)$$

Then one path in the square gives us

$$\begin{aligned} & \Phi^\circ(D_{p'}(f'))_X(\Phi D_{\mathcal{F}', X}((F, \beta'))) = \\ & \Phi^\circ(D_{p'}(f'))_X((\Phi(F) \circ \phi, \mathcal{F}'(\iota)(\beta'))) = D_{p'}(f')_{\Phi(X)}((\Phi(F) \circ \phi, \mathcal{F}'(\iota)(\beta'))) = \\ & (\Phi(F) \circ \phi, f'_{(\Phi(X); \Phi(F) \circ f)}(\mathcal{F}'(\iota)(\beta'))) \end{aligned}$$

where the first equality is by (4.8), the second by the definition of  $\Phi^\circ$  and the third by (2.23).

The other path gives us

$$\begin{aligned} & \Phi D_{\mathcal{G}', X}(D_p(\Phi^\circ(f'))_X((F, \beta'))) = \\ & \Phi D_{\mathcal{G}', X}((F, (\Phi^\circ(f'))_{(X; F)}(\beta'))) = \Phi D_{\mathcal{G}', X}((F, f'_{\Phi((X; F))}(\beta'))) = \\ & (\Phi(F) \circ \phi, \mathcal{G}'(\iota)(f'_{\Phi((X; F))}(\beta'))) \end{aligned}$$

where the first equality is by (2.23), the second by the definition of  $\Phi^\circ$  and the third by (4.8).

It remains to show that

$$f'_{(\Phi(X); \Phi(F) \circ f)}(\mathcal{F}'(\iota)(\beta')) = \mathcal{G}'(\iota)(f'_{\Phi((X; F))}(\beta'))$$

which follows from the axiom of compatibility with morphisms of the natural transformation  $f' : \mathcal{F}' \rightarrow \mathcal{G}'$ . This completes the proof of commutativity of (4.5) and with it Construction 4.3.

**Problem 4.4** [2016.12.18.probl] *Let  $\Phi : (\mathcal{C}, \mathbf{p}) \rightarrow (\mathcal{C}', \mathbf{p}')$  be a universe category functor. Let  $F \in \text{PreShv}(\mathcal{C})$ ,  $F' \in \text{PreShv}(\mathcal{C}')$  and let*

$$m : F \rightarrow \Phi^\circ(F')$$

*be a morphism of presheaves. Let  $n \in \mathbf{N}$ . To construct a morphism of presheaves*

$$D^n(m) : D_p^n(F) \rightarrow \Phi^\circ(D_{p'}^n(F'))$$

**Construction 4.5** [2016.12.18.constr1] We proceed by induction on  $n$ .

For  $n = 0$  we set  $D^0m = m$ .

For the successor of  $n$  we need to construct a morphism

$$D^{n+1}(m) : D_p(D_p^n(F)) \rightarrow \Phi^\circ(D_{p'}(D_{p'}^n(F')))$$

We define it as the composition

$$D_p(D_p^n(F)) \xrightarrow{D_p(D^n(m))} D_p(\Phi^\circ(D_{p'}^n(F'))) \xrightarrow{\Phi D_{p'}^n(F')} \Phi^\circ(D_{p'}(D_{p'}^n(F')))$$

**Lemma 4.6** [2016.12.18.11] *In the assumptions of Problem 4.4 consider a commutative square in  $PreShv(\mathcal{C})$  of the form*

$$\begin{array}{ccc} F_1 & \xrightarrow{m_1} & \Phi^\circ(F'_1) \\ \text{[2016.12.18.eq1]}_p \downarrow & & \downarrow \Phi^\circ(v') \\ F_2 & \xrightarrow{m_2} & \Phi^\circ(F'_2) \end{array} \quad (4.12)$$

Then, for any  $n \in \mathbf{N}$ , the square

$$\begin{array}{ccc} D_p^n(F_1) & \xrightarrow{D^n(m_1)} & \Phi^\circ(D_{p'}^n(F'_1)) \\ \text{[2016.12.18.eq2]}_p^n(v) \downarrow & & \downarrow \Phi^\circ(D_{p'}^n(v')) \\ D_p^n(F_2) & \xrightarrow{D^n(m_2)} & \Phi^\circ(D_{p'}^n(F'_2)) \end{array} \quad (4.13)$$

commutes.

**Proof:** We proceed by induction on  $n$ .

For  $n = 0$  the square (4.13) coincides with the square (4.12).

For the successor of  $n$ , (4.13) is the external square of the diagram

$$\begin{array}{ccccc} D_p(D_p^n(F_1)) & \xrightarrow{D_p(D^n(m_1))} & D_p(\Phi^\circ(D_{p'}^n(F'_1))) & \xrightarrow{\Phi D_{p'}^n(F'_1)} & \Phi^\circ(D_{p'}(D_{p'}^n(F'_1))) \\ D_p(D_p^n(v)) \downarrow & & D_p(\Phi^\circ(D_{p'}^n(v'))) \downarrow & & \downarrow \Phi^\circ(D_{p'}(D_{p'}^n(v'))) \\ D_p(D_p^n(F_2)) & \xrightarrow{D_p(D^n(m_2))} & D_p(\Phi^\circ(D_{p'}^n(F'_2))) & \xrightarrow{\Phi D_{p'}^n(F'_2)} & \Phi^\circ(D_{p'}(D_{p'}^n(F'_2))) \end{array}$$

The left hand side square in this diagram is obtained by applying  $D_p$  to the square (4.13) for  $n$ . It is commutative because  $D_p$  is a functor and in particular satisfies the composition axiom (2.25).

The right hand side square is commutative because  $\Phi D$  is a natural transformation of functors that satisfies the axiom of compatibility with morphisms of presheaves. In our particular case this axiom is applied to the morphism of presheaves  $D_{p'}^n(v')$ .

This completes the proof of the lemma.

The following problem and construction are the only ones in this section where we change our context to simply a functor between two categories.

**Problem 4.7 [2016.12.18.prob3]** *Given a functor  $\Phi : \mathcal{C} \rightarrow \mathcal{C}'$  between two categories to construct, for each  $Y \in \mathcal{C}$ , a morphism of presheaves*

$$yo^{\Phi, Y} : Yo(Y) \rightarrow \Phi^\circ(Yo(\Phi(Y)))$$

and to show that for a morphism  $g : Y \rightarrow Y'$  the square

$$\begin{array}{ccc} Yo(Y) & \xrightarrow{yo^{\Phi, Y}} & \Phi^\circ(Yo(\Phi(Y))) \\ \text{[2016.12.18.eq8]}_{o(g)} \downarrow & & \downarrow \Phi^\circ(Yo(\Phi(g))) \\ Yo(Y') & \xrightarrow{yo^{\Phi, Y'}} & \Phi^\circ(Yo(\Phi(Y'))) \end{array} \quad (4.14)$$

commutes.

**Construction 4.8 [2016.12.18.constr3]** We need to define, for all  $X \in \mathcal{C}$ , functions

$$Yo(Y)(X) = Mor_{\mathcal{C}}(X, Y) \rightarrow Mor_{\mathcal{C}'}(\Phi(X), \Phi(Y)) = \Phi^\circ(Yo(\Phi(Y)))(X)$$

which we define as the restriction of  $\Phi_{Mor}$  to  $Mor_{\mathcal{C}}(X, Y)$ :

$$\text{[2016.12.18.eq7]} yo_X^{\Phi, Y}(f) = \Phi(f) \quad (4.15)$$

Let us show that this family is a morphism of presheaves, i.e., that for any  $a : X' \rightarrow X$  the square

$$\begin{array}{ccc} Yo(Y)(X) & \xrightarrow{yo_X^{\Phi, Y}} & \Phi^\circ(Yo(\Phi(Y)))(X) \\ \text{[2016.12.18.eq5]}_{(Y)(a)} \downarrow & & \downarrow \Phi^\circ(Yo(\Phi(Y)))(a) \\ Yo(Y)(X') & \xrightarrow{yo_{X'}^{\Phi, Y}} & \Phi^\circ(Yo(\Phi(Y)))(X') \end{array} \quad (4.16)$$

commutes. Note that for an element  $f' : \Phi(X) \rightarrow \Phi(Y)$  of  $\Phi^\circ(Yo(\Phi(Y)))(X)$  we have

$$\text{[2016.12.18.eq6]} \Phi^\circ(Yo(\Phi(Y)))(a)(f') = \Phi(a) \circ f' \quad (4.17)$$

Let  $f : X \rightarrow Y$  be an element of  $Yo(Y)(X)$ .

Applying one path in (4.16) to  $f$  we get

$$\Phi^\circ(Yo(\Phi(Y)))(a)(yo_X^{\Phi, Y}(f)) = \Phi^\circ(Yo(\Phi(Y)))(a)(\Phi(f)) = \Phi(a) \circ \Phi(f)$$

where the first equality is by (4.15) and the second is by (4.17).

Applying another path we get

$$yo_{X'}^{\Phi, Y}(Yo(Y)(a)(f)) = \Phi Yo(Y)_{X'}(a \circ f) = \Phi(a \circ f)$$

where the first equality is by definition of  $Yo(Y)$  and the second by (4.15).

We conclude that (4.16) commutes by the composition axiom of  $\Phi$ .

Let  $g : Y \rightarrow Y'$  be a morphism. Note that for an element  $f' : \Phi(X) \rightarrow \Phi(Y)$  of  $\Phi^\circ(Yo(\Phi(Y)))(X)$  we have

$$[\mathbf{2016.12.18.eq9}] \Phi^\circ(Yo(\Phi(g)))(f') = f' \circ \Phi(g) \quad (4.18)$$

Let us show that the square (4.14) commutes. Let  $X \in \mathcal{C}$  and  $f \in Yo(Y)(X)$ .

Applying one path in (4.14) to  $f$  we get

$$\Phi^\circ(Yo(\Phi(g)))(yo^{\Phi, Y}(f)) = \Phi^\circ(Yo(\Phi(g)))(\Phi(f)) = \Phi(f) \circ \Phi(g)$$

where the first equality is by (4.15) and the second by (4.18).

Applying another path we get

$$yo^{\Phi, Y'}(Yo(g)(f)) = yo^{\Phi, Y'}(f \circ g) = \Phi(f \circ g)$$

where the first equality is by the definition of  $Yo(g)$  and the second by (4.15). We conclude that (4.14) commutes by the composition axiom of  $\Phi$ .

This completes the construction.

Recall that for  $X, Y \in \mathcal{C}$  and  $n \geq 0$  we have defined in Remark 2.14 the set  $D_p^n(X, Y)$  as follows:

$$D_p^n(X, Y) = D_p^n(Yo(Y))(X)$$

Since  $D_p^n(Yo(Y))$  is a presheaf we have, for any  $f : X' \rightarrow X$ , the function

$$D_p^n(Yo(Y))(f) : D_p^n(Yo(Y))(X) \rightarrow D_p^n(Yo(Y))(X')$$

that we denote by

$$D_p^n(f, Y) : D_p^n(X, Y) \rightarrow D_p^n(X', Y)$$

Since  $D_p^n$  and  $Yo$  are functors we have, for any  $g : Y \rightarrow Y'$ , a function

$$D_p^n(Yo(g))_X : D_p^n(Yo(Y))(X) \rightarrow D_p^n(Yo(Y'))(X)$$

that we denote by

$$D_p^n(X, g) : D_p^n(X, Y) \rightarrow D_p^n(X, Y')$$

Since the family of functions  $D_p^n(Yo(g))_-$  is a morphism of presheaves the functions  $D_p^n(f, -)$  and  $D_p^n(-, g)$  commute.

**Problem 4.9** [ $\mathbf{2016.12.18.prob2}$ ] *In the assumptions as above, to define, for all  $X, Y \in \mathcal{C}$  and  $n \geq 0$ , functions*

$$\Phi_{X, Y}^n : D_p^n(X, Y) \rightarrow D_{p'}^n(\Phi(X), \Phi(Y))$$

**Construction 4.10** [2016.12.18.constr2] Applying Construction 4.5 to the morphism of presheaves  $yo^{\Phi, Y}$  of Construction 4.8 we obtain morphisms of presheaves

$$D^n(yo^{\Phi, Y}) : D_p^n(Yo(Y)) \rightarrow \Phi^\circ(D_{p'}^n(Yo(\Phi(Y))))$$

Evaluating this morphism on  $X$  we obtain a function

$$[2016.12.20.eq3] D_p^n(X, Y) = D_p^n(Yo(Y))(X) \rightarrow \Phi^\circ(D_{p'}^n(Yo(\Phi(Y))))(X) = D_{p'}^n(\Phi(X), \Phi(Y)) \quad (4.19)$$

**Lemma 4.11** [2016.12.20.11] *In the context of Construction 4.10 one has:*

1. *let  $f : X' \rightarrow X$  be a morphism, then the square*

$$[2016.12.20.eq1] \begin{array}{ccc} D_p^n(X, Y) & \xrightarrow{\Phi_{X, Y}^n} & D_{p'}^n(\Phi(X), \Phi(Y)) \\ \downarrow D_p^n(f, Y) & & \downarrow D_{p'}^n(\Phi(f), \Phi(Y)) \\ D_p^n(X', Y) & \xrightarrow{\Phi_{X', Y}^n} & D_{p'}^n(\Phi(X'), \Phi(Y)) \end{array} \quad (4.20)$$

2. *let  $g : Y \rightarrow Y'$  be a morphism, then the square*

$$[2016.12.20.eq2] \begin{array}{ccc} D_p^n(X, Y) & \xrightarrow{\Phi_{X, Y}^n} & D_{p'}^n(\Phi(X), \Phi(Y)) \\ \downarrow D_p^n(X, g) & & \downarrow D_{p'}^n(\Phi(X), \Phi(g)) \\ D_p^n(X, Y') & \xrightarrow{\Phi_{X, Y'}^n} & D_{p'}^n(\Phi(X), \Phi(Y')) \end{array} \quad (4.21)$$

*commutes.*

**Proof:** Commutativity of (4.20) follows from (4.19) and the fact that  $D^n(yo^{\Phi, Y})$  is a morphism of presheaves.

Commutativity of (4.21) follows from (4.19), the commutativity of (4.14) and Lemma 4.6.

## 4.2 Universe category functors and isomorphisms $u_n$ and $\tilde{u}_n$

## 4.3 Universe category functors and the $I_p$ construction

Note that in the problem below no assumption is made about the compatibility of  $\Phi$  with the locally cartesian closed structures on  $\mathcal{C}$  and  $\mathcal{C}'$ .

**Problem 4.12** [2015.03.21.prob1] *Assume that  $\mathcal{C}$  and  $\mathcal{C}'$  are locally cartesian closed universe categories. For  $\Phi$  as above and  $V \in \mathcal{C}$  to construct a morphism*

$$\chi_\Phi(V) : \Phi(I_p(V)) \rightarrow I_{p'}(\Phi(V))$$

**Construction 4.13** [2015.03.21.constr1] Consider the sequence of functions

$$\begin{array}{ccc} D_p(I_p(V), V) & \xrightarrow{\Phi_{I_p(V), V}^2} & D_{p'}(\Phi(I_p(V)), \Phi(V)) \\ \eta_{V, I_p(V)}^\dagger \uparrow & & \eta_{\Phi(I_p(V)), \Phi(V)} \downarrow \\ \text{Mor}(I_p(V), I_p(V)) & & \text{Mor}(\Phi(I_p(V)), I_{p'}(\Phi(V))) \end{array}$$

where  $\eta$  and its inverse  $\eta^\dagger$  are the bijections of Construction 2.35. Applying it to  $Id_{I_p(V)}$  we obtain

$$\chi_\Phi(V) = \eta_{\Phi(I_p(V)), \Phi(V)}(\Phi_{I_p(V), V}^2(\eta_{V, I_p(V)}^\dagger(Id_{I_p(V)})))$$

Let us show that  $\chi_\Phi$  are natural in  $V$ .

**Lemma 4.14** [2015.04.10.14] For  $\Phi$  as above let  $f : V_1 \rightarrow V_2$  be a morphism. Then the square

$$\begin{array}{ccc} \Phi(I_p(V_1)) & \xrightarrow{\chi_\Phi(V_1)} & I_{p'}(\Phi(V_1)) \\ \Phi(I_p(f)) \downarrow & & \downarrow I_{p'}(\Phi(f)) \\ \Phi(I_p(V_2)) & \xrightarrow{\chi_\Phi(V_2)} & I_{p'}(\Phi(V_2)) \end{array}$$

commutes.

**Proof:** Let  $X_i = I_p(V_i)$  for  $i = 1, 2$ . In the following computations we often omit the indexes that can be recovered from the context. We have:

$$\chi(V_1) \circ I_{p'}(\Phi(V_1)) = \eta(\Phi^2(\eta^\dagger(Id_{X_1}))) \circ I_{p'}(\Phi(f)) = \eta'(D_{p'}(\Phi(X_1), \Phi(f))(\Phi^2(\eta^\dagger(Id_{X_1}))))$$

where the second equality is by (2.56) with respect to  $\Phi(f)$ . Then

$$\begin{aligned} \eta(D_p(X_1, \Phi(f))(\Phi^2(\eta^\dagger(Id_{X_1})))) &= \eta(\Phi^2(D_p(X_1, f)(\eta^\dagger(Id_{X_1})))) = \\ &= \eta(\Phi^2(\eta^\dagger(Id_{X_1} \circ I_p(f)))) = \eta(\Phi^2(\eta^\dagger(I_p(f)))) \end{aligned}$$

where the first equality holds by Lemma ?? and the second by (2.57) with respect to  $f$ .

On the other hand:

$$\begin{aligned} \Phi(I_p(f)) \circ \chi(V_2) &= \Phi(I_p(f)) \circ \eta(\Phi^2(\eta^\dagger(Id_{X_2}))) = \\ &= \eta'(D_{p'}(\Phi(I_p(f)), \Phi(X_2))(\Phi^2(\eta^\dagger(Id_{X_2})))) \end{aligned}$$

where the second equality is by (2.56) with respect to  $\Phi(I_p(f))$ . Then

$$\begin{aligned} \eta(D_{p'}(\Phi(I_p(f)), \Phi(X_2))(\Phi^2(\eta^\dagger(Id_{X_2})))) &= \eta(\Phi^2(D_p(I_p(f), X_2)(\eta^\dagger(Id_{X_2})))) = \\ &= \eta(\Phi^2(\eta^\dagger(I_p(f) \circ Id_{X_2}))) = \eta(\Phi^2(\eta^\dagger(I_p(f)))) \end{aligned}$$

where the first equality holds by Lemma ?? and the second by (2.58) with respect to the  $I_p(f)$ . This completes the proof of Lemma 4.14.



**Lemma 4.15** [2015.05.06.11] For all  $X, V \in \mathcal{C}$  the pentagon

$$\begin{array}{ccc}
D_p(X, V) & \xrightarrow{\eta_{V, X}} & \text{Mor}(X, I_p(V)) \\
\Phi_{V, X}^2 \downarrow & & \downarrow \Phi \\
D_{p'}(\Phi(X), \Phi(V)) & & \text{Mor}(\Phi(X), \Phi(I_p(V))) \\
\eta_{\Phi(V), \Phi(X)} \downarrow & & \downarrow -\circ\chi_{\Phi(V)} \\
\text{Mor}(\Phi(X), I_{p'}(\Phi(V))) & \text{=====} & \text{Mor}(\Phi(X), I_{p'}(\Phi(V)))
\end{array}$$

commutes, that is, for all  $a \in D_p(X, V)$  one has

$$\Phi(\eta(a)) \circ \chi_{\Phi(V)} = \eta(\Phi^2(a))$$

**Proof:** By definition of  $\chi_{\Phi}$  and (2.56) with respect to  $\Phi(\eta(a))$  we have

$$\Phi(\eta(a)) \circ \chi_{\Phi(V)} = \Phi(\eta(a)) \circ \eta(\Phi^2(\eta^!(Id))) = \eta(D_{p'}(\Phi(\eta(a)), \Phi(V))(\Phi^2(\eta^!(Id_{I_p(V)}))))$$

By Lemma ?? we further have:

$$\eta(D_{p'}(\Phi(\eta(a)), \Phi(V))(\Phi^2(\eta^!(Id)))) = \eta(\Phi^2(D_p(\eta(a), V)(\eta^!(Id))))$$

It remains to show that  $D_p(\eta(a), V)(\eta^!(Id)) = a$ . Since  $\eta$  is a bijection we may apply it on both sides and by (2.58) with respect to  $\eta(a)$  we get

$$\eta(D_p(\eta(a), V)(\eta^!(Id))) = \eta(\eta^!(\eta(a) \circ Id)) = \eta(a) \circ Id = \eta(a)$$

## 4.4 More on universe category functors

By [?, Construction 4.7] any universe category functor  $\Phi = (\Phi, \phi, \tilde{\phi})$  from  $(\mathcal{C}, p)$  to  $(\mathcal{C}', p')$  defines a homomorphism of C-systems

$$H : CC(\mathcal{C}, p) \rightarrow CC(\mathcal{C}', p')$$

Let  $\psi_0 : pt' \rightarrow \Phi(pt)$  be the unique morphism. To define  $H$  on objects, one uses the fact that

$$Ob(CC(\mathcal{C}, p)) = \coprod_{n \geq 0} Ob_n(\mathcal{C}, p)$$

and defines  $H(n, A)$  as  $(n, H_n(A))$  where

$$H_n : Ob_n(\mathcal{C}, p) \rightarrow Ob_n(\mathcal{C}', p')$$

To obtain  $H_n$  one defines by induction on  $n$ , pairs  $(H_n, \psi_n)$  where  $H_n$  is as above and  $\psi_n$  is a family of isomorphisms

$$\psi_n(A) : int_n(H_n(A)) \rightarrow \Phi(int_n(A))$$

as follows:

1. for  $n = 0$ ,  $H_0$  is the unique function from a one point set to a one point set and  $\psi_0(A) = \psi_0$ ,
2. for the successor of  $n$  one has

$$[\mathbf{2016.12.10.eq1}] H_{n+1}(A, F) = (H_n(A), \psi_n(A) \circ \Phi(F) \circ \phi) \quad (4.22)$$

and  $\psi_{n+1}(A, F)$  is the unique morphism  $int(H(A, F)) \rightarrow \Phi(int(A, F))$  such that

$$[\mathbf{2016.12.10.eq2}] \psi_{n+1}(A, F) \circ \Phi(Q(F)) \circ \tilde{\phi} = Q(\psi_n(A) \circ \Phi(F) \circ \phi) \quad (4.23)$$

and

$$[\mathbf{2016.12.10.eq3}] \psi_{n+1}(A, F) \circ \Phi(p_F) = p_{\psi_n(A) \circ \Phi(F) \circ \phi} \circ \psi_n(A) \quad (4.24)$$

The function  $H : Ob(CC(\mathcal{C}, p)) \rightarrow Ob(CC(\mathcal{C}', p'))$  is the sum of functions  $H_n$ . For  $\Gamma = (n, A)$  in  $Ob(CC(\mathcal{C}, p))$  we let  $\psi(\Gamma) = \psi_n(A)$  such that  $\psi$  is the sum of families  $\psi_n$ .

The action of  $H$  on morphisms is given, for  $f : \Gamma' \rightarrow \Gamma$ , by

$$[\mathbf{2016.12.10.eq4}] H(f) = \psi(\Gamma') \circ \Phi(int(f)) \circ \psi(\Gamma)^{-1} \quad (4.25)$$

We will often write  $H$  also for the functions  $H_n$  and  $\psi$  for the functions  $\psi_n$ .

**Lemma 4.16** *[2015.03.21.14] Let  $(\Phi, \phi, \tilde{\phi})$  be universe category functor. Then:*

1. for  $T \in Ob_1(\Gamma)$  one has

$$u_{1, H(\Gamma)}(H(T)) = \psi(\Gamma) \circ \Phi(u_{1, \Gamma}(T)) \circ \phi$$

2. for  $o \in \tilde{Ob}_1(\Gamma)$  one has

$$\tilde{u}_{1, H(\Gamma)}(H(o)) = \psi(\Gamma) \circ \Phi(\tilde{u}_{1, \Gamma}(o)) \circ \tilde{\phi}$$

**Proof:** Let  $\Gamma = (n, A)$ .

In the case of  $T \in Ob_1(\Gamma)$ , if  $T = (n + 1, (A, F))$  then

$$u_1(H(T)) = u_1(n + 1, H(A, F)) = u_1(n + 1, (H(A), \psi(\Gamma) \circ \Phi(F) \circ \phi)) = \psi(\Gamma) \circ \Phi(F) \circ \phi$$

where the last equality is by (2.33).

In the case of  $o \in \tilde{Ob}_1(\Gamma)$ , if  $\partial(o) = (n + 1, (A, F))$  then  $\partial(H(o)) = (n + 1, H(A, F))$ . Since  $o : \Gamma \rightarrow \partial(o)$  we have

$$[\mathbf{2016.12.10.eq6}] H(o) = \psi(\Gamma) \circ \Phi(int(o)) \circ \psi(A, F)^{-1} \quad (4.26)$$

and

$$\tilde{u}_1(H(o)) = H(o) \circ Q(u_1(n + 1, H(A, F))) = H(o) \circ Q(\psi(A) \circ \Phi(F) \circ \phi) =$$

$$\begin{aligned}
H(o) \circ \psi(A, F) \circ \Phi(Q(F)) \circ \tilde{\phi} &= \psi(\Gamma) \circ \Phi(\text{int}(o)) \circ \psi(A, F)^{-1} \circ \psi(A, F) \circ \Phi(Q(F)) \circ \tilde{\phi} = \\
&= \psi(\Gamma) \circ \Phi(\text{int}(o)) \circ \Phi(Q(F)) \circ \tilde{\phi} = \psi(\Gamma) \circ \Phi(\text{int}(o) \circ Q(F)) \circ \tilde{\phi} = \\
&= \psi(\Gamma) \circ \Phi(\tilde{u}_1(o)) \circ \tilde{\phi}
\end{aligned}$$

Where the first equality is by (2.40), the second by (4.22) and (2.33), the third by (4.23), the fourth by (4.26) and the seventh again by (2.40).

**Remark 4.17** [2016.12.12.rem1] The assertion of Lemma 4.16(1) can be expressed by saying that the diagram of natural transformations

$$\begin{array}{ccc}
\mathcal{O}b_1 & \xrightarrow{u_1} & \text{int}^\circ(Yo(U)) \\
& & \downarrow \text{int}^\circ(\Phi_{-,U}) \\
& & \text{int}^\circ(\Phi^\circ(Yo(\Phi(U)))) \\
\downarrow H\mathcal{O}b_1 & & \downarrow \text{int}^\circ(\Phi^\circ(Yo(\phi))) \\
& & \text{int}^\circ(\Phi^\circ(Yo(U'))) \\
& & \downarrow \psi^\circ \\
H^\circ(\mathcal{O}b_1) & \xrightarrow{H^\circ(u_1)} & H^\circ(\text{int}^\circ(Yo(U')))
\end{array}$$

commutes. The same, with the replacement of  $U, U'$  and  $\phi$  by  $\tilde{U}, \tilde{U}'$  and  $\tilde{\phi}$  applies to Lemma 4.16(2). To make it precise would, however, require some extra work because we have not proved that the composition with morphisms  $\psi(\Gamma)$  defines a natural transformation of the form  $\Phi^\circ \circ \text{int}^\circ \rightarrow \text{int}^\circ \circ H^\circ$  and we have not proved that for any  $V$  the functions

$$\Phi_{-,V} : \text{Mor}_{\mathcal{C}}(-, V) \rightarrow \text{Mor}_{\mathcal{C}'}(\Phi(-), \Phi(V))$$

form a natural transformation from  $Yo(V)$  to  $\Phi^\circ(Yo(\Phi(V)))$ .

**Lemma 4.18** [2015.03.21.14b] *Let  $(\Phi, \phi, \tilde{\phi})$  be universe category functor. Then:*

1. *for  $T \in \mathcal{O}b_2(\Gamma)$  one has*

$$u_{2,H(\Gamma)}(H(T)) = D_{p'}(\psi(\Gamma), U')(D_{p'}(\text{int}(H(\Gamma)), \phi)(\Phi^2(u_{2,\Gamma}(T))))$$

2. *for  $o \in \tilde{\mathcal{O}}b_2(\Gamma)$  one has*

$$\tilde{u}_{2,H(\Gamma)}(H(o)) = D_{p'}(\psi(\Gamma), \tilde{U}')(D_{p'}(\text{int}(H(\Gamma)), \tilde{\phi})(\Phi^2(\tilde{u}_{2,\Gamma}(o))))$$

**Proof:** Let  $\Gamma = (n, A)$ .

In the case  $T \in \mathcal{O}b_2(\Gamma)$ , if  $T = (n + 2, ((A, F_1), F_2))$  then

$$u_2(H(T)) = u_2(n + 2, H(((A, F_1), F_2))) = u_2(n + 2, (H(A, F_1), \psi(A, F_1) \circ \Phi(F_2) \circ \phi)) =$$

$$u_2(n+2, ((H(A), \psi(A) \circ \Phi(F_1) \circ \phi), \psi(A, F_1) \circ \Phi(F_2) \circ \phi)) = \\ (\psi(A) \circ \Phi(F_1) \circ \phi, \psi(A, F_1) \circ \Phi(F_2) \circ \phi)$$

On the other hand

$$D_{p'}(\psi(A), -)D_{p'}(-, \phi)(\Phi^2(u_2(T))) = D_{p'}(\psi(A), -)D_{p'}(-, \phi)(\Phi^2(u_2(n+2, ((A, F_1), F_2)))) = \\ D_{p'}(\psi(A), -)D_{p'}(-, \phi)(\Phi^2(F_1, F_2)) = D_{p'}(\psi(A), -)D_{p'}(-, \phi)(\Phi(F_1) \circ \phi, \iota \circ \Phi(F_2)) = \\ D_{p'}(\psi(A), -)(\Phi(F_1) \circ \phi, \iota \circ \Phi(F_2) \circ \phi) = (\psi(A) \circ \Phi(F_1) \circ \phi, Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(F_2) \circ \phi)$$

therefore we need to show that

$$[\mathbf{2015.04.12.eq1}]\psi(A, F_1) \circ \Phi(F_2) \circ \phi = Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(F_2) \circ \phi \quad (4.27)$$

which we reduce to  $\psi(A, F_1) = Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota$ . The codomain of both sides is  $\Phi(\text{int}(A, F_1))$ . Using the fact that the external square of the diagram

$$\begin{array}{ccccc} \Phi(\text{int}(A, F_1)) & \xrightarrow{\Phi(Q(F_1))} & \Phi(\tilde{U}) & \xrightarrow{\tilde{\phi}} & \tilde{U}' \\ \Phi(p_{F_1}) \downarrow & & \downarrow \Phi(p) & & \downarrow p' \\ \Phi(\text{int}(A)) & \xrightarrow{\Phi(F_1)} & \Phi(U) & \xrightarrow{\phi} & U' \end{array}$$

is a pullback we see that equality (4.27) would follow from the following two equalities:

$$\psi(A, F_1) \circ \Phi(Q(F_1)) \circ \tilde{\phi} = Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(Q(F_1)) \circ \tilde{\phi}$$

and

$$\psi(A, F_1) \circ \Phi(p_{F_1}) = Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(p_{F_1})$$

For the first equality we have

$$\psi(A, F_1) \circ \Phi(Q(F_1)) \circ \tilde{\phi} = Q(\psi(A) \circ \Phi(F_1) \circ \phi)$$

by definition of  $\psi(\Gamma, F_1)$  and

$$Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(Q(F_1)) \circ \tilde{\phi} = Q(\psi(A), \Phi(F_1) \circ \phi) \circ Q(\Phi(F_1) \circ \phi) = Q(\psi(A) \circ \Phi(F_1) \circ \phi)$$

where the first equality holds by definition of  $\iota$  and second by the definition of  $Q(-, -)$ .

For the second equality we have

$$\psi(A, F_1) \circ \Phi(p_{F_1}) = p_{H(A, F_1)} \circ \psi(A)$$

by definition of  $\psi(A, F_1)$  and

$$Q(\psi(A), \Phi(F_1) \circ \phi) \circ \iota \circ \Phi(p_{F_1}) = Q(\psi(A), \Phi(F_1) \circ \phi) \circ p_{\Phi(F_1) \circ \phi} = p_{H(A, F_1)} \circ \psi_\Gamma$$

by definitions of  $Q$  and  $\iota$ .

The case of  $o \in \widetilde{Ob}_2(\Gamma)$  is strictly parallel to the case of  $T \in Ob_2(\Gamma)$  with  $\Phi(F_2) \circ \phi$  at the end of the formulas replaced by  $\Phi(\tilde{F}_2) \circ \tilde{\phi}$  where instead of  $F_2 : \text{int}(A, F_1) \rightarrow U$  one has  $\tilde{F}_2 : \text{int}(A, F_1) \rightarrow \tilde{U}$  with  $\tilde{F}_2 = \tilde{u}_{1, ft(\partial(o))}(o)$ .

**Remark 4.19** [2016.12.12.rem2]

$$\begin{array}{ccccccc}
\mathcal{O}b_{n+1} & \longrightarrow & \text{Sig}(\mathcal{O}b_n) & \longrightarrow & \text{Sig}(\text{int}^\circ(D_p^n(Yo(U)))) & \longrightarrow & \text{int}^\circ(D_p^{n+1}(Yo(U))) \\
& & & & \downarrow & & \downarrow \\
& & \downarrow & & \text{Sig}(\text{int}^\circ(\Phi^\circ(D_{p'}^n(Yo(U'))))) & \longrightarrow & \text{int}^\circ(D_{p'}(\Phi^\circ(D_{p'}^n(Yo(U'))))) \\
& & & & \downarrow & & \downarrow \\
\downarrow & & \text{Sig}(H^\circ(\mathcal{O}b_n)) & \longrightarrow & \text{Sig}(H^\circ(\text{int}^\circ(D_{p'}^n(Yo(U'))))) & & \text{int}^\circ(\Phi^\circ(D_{p'}^{n+1}(Yo(U')))) \\
& & \downarrow & & \downarrow & & \downarrow \\
H^\circ(\mathcal{O}b_{n+1}) & \longrightarrow & H^\circ(\text{Sig}(\mathcal{O}b_n)) & \longrightarrow & H^\circ(\text{Sig}(\text{int}^\circ(D_{p'}^n(Yo(U'))))) & \longrightarrow & H^\circ(\text{int}^\circ(D_{p'}^{n+1}(Yo(U'))))
\end{array}$$

For  $(\Phi, \phi, \tilde{\phi})$  as above let us denote by

$$\xi_{\Phi} : \Phi(I_p(U)) \rightarrow I_{p'}(U')$$

the composition  $\chi_{\Phi}(U) \circ I_{p'}(\phi)$  and by

$$\tilde{\xi}_{\Phi} : \Phi(I_p(\tilde{U})) \rightarrow I_{p'}(\tilde{U}')$$

the composition  $\chi_{\Phi}(\tilde{U}) \circ I_{p'}(\tilde{\phi})$ .

**Lemma 4.20** [2015.05.06.12] *Let  $(\Phi, \phi, \tilde{\phi})$  be a universe category functor and  $\Gamma \in \text{Ob}(CC(\mathcal{C}, p))$ . Then one has:*

1. for  $T \in \text{Ob}_2(\Gamma)$

$$\mu_2(H(T)) = \psi(\Gamma) \circ \Phi(\mu_2(T)) \circ \xi_{\Phi}$$

2. for  $o \in \tilde{\mathcal{O}}b_2(\Gamma)$

$$\tilde{\mu}_2(H(o)) = \psi(\Gamma) \circ \Phi(\tilde{\mu}_2(o)) \circ \tilde{\xi}_{\Phi}$$

**Proof:** We have

$$\begin{aligned}
\mu_2(H(T)) &= \eta(u_2(H(T))) = \eta(D_{p'}(\psi(\Gamma), -)(D_{p'}(-, \phi)(\Phi^2(u_2(T)))))) = \\
&\quad \psi(\Gamma) \circ \eta(\Phi^2(u_2(T))) \circ I_{p'}(\phi)
\end{aligned}$$

where the first equality holds by the definition of  $\mu_2$  (cf. Construction 2.39), the second equality holds by Lemma 4.16(3) and the third by the naturality of  $\eta$ . Next

$$\eta(\Phi^2(u_2(T))) \circ I_{p'}(\phi) = \Phi(\eta(u_2(T))) \circ \chi_{\Phi}(U) \circ I_{p'}(\phi) = \Phi(\eta(u_2(T))) \circ \xi_{\Phi} = \Phi(\mu_2(T)) \circ \xi_{\Phi}$$

where the first equality holds by Lemma 4.15, the second one by the definition of  $\xi_{\Phi}$  and the third one by the definition of  $\mu_2$ .

The proof of the second part of the lemma is strictly parallel to the proof of the first part.

## 4.5 Functoriality properties of the $(\Pi, \lambda)$ -structures constructed from $P$ -structures

Recall that in [?, pp. 1067-68] we have constructed, for any homomorphism  $H : CC \rightarrow CC'$  of  $C$ -systems, and any  $n \geq 0$ , natural transformations

$$H\mathcal{O}b_n : \mathcal{O}b_i \rightarrow H^\circ(\mathcal{O}b_i)$$

where for  $\Gamma \in CC$  and  $T \in \mathcal{O}b_i(\Gamma)$  one has

$$H\mathcal{O}b_n(T) = H_{Ob}(T)$$

and

$$H\widetilde{\mathcal{O}}b_n : \widetilde{\mathcal{O}}b_i \rightarrow H^\circ(\widetilde{\mathcal{O}}b_i)$$

where for  $\Gamma \in CC$  and  $o \in \widetilde{\mathcal{O}}b_n(\Gamma)$  one has

$$H\widetilde{\mathcal{O}}b_n(o) = H_{Mor}(o)$$

**Definition 4.21** [2016.09.13.def1] *Let  $H : CC \rightarrow CC'$  be a homomorphism of  $C$ -systems. Let  $(\Pi, \lambda)$  and  $(\Pi', \lambda')$  be pre- $(\Pi, \lambda)$ -structures on  $CC$  and  $CC'$  respectively.*

*Then  $H$  is called a  $(\Pi, \lambda)$ -homomorphism if the following two squares commute*

$$\begin{array}{ccc} \mathcal{O}b_2 & \xrightarrow{\Pi} & \mathcal{O}b_1 \\ H\mathcal{O}b_2 \downarrow & & \downarrow H\mathcal{O}b_1 \\ H^\circ(\mathcal{O}b_2) & \xrightarrow{H^\circ(\Pi')} & H^\circ(\mathcal{O}b_1) \end{array} \quad \begin{array}{ccc} \widetilde{\mathcal{O}}b_2 & \xrightarrow{\lambda} & \widetilde{\mathcal{O}}b_1 \\ H\widetilde{\mathcal{O}}b_2 \downarrow & & \downarrow H\widetilde{\mathcal{O}}b_1 \\ H^\circ(\widetilde{\mathcal{O}}b_2) & \xrightarrow{H^\circ(\lambda')} & H^\circ(\widetilde{\mathcal{O}}b_1) \end{array}$$

*If  $(\Pi, \lambda)$  and  $(\Pi', \lambda')$  are  $(\Pi, \lambda)$ -structures then  $H$  is called a  $(\Pi, \lambda)$ -homomorphism if it is a  $(\Pi, \lambda)$ -homomorphism with respect to the corresponding pre- $(\Pi, \lambda)$ -structures.*

Unfolding the definition of  $H\mathcal{O}b_i$  and  $H\widetilde{\mathcal{O}}b_i$  we see that  $H$  is a  $(\Pi, \lambda)$ -homomorphism if and only if for all  $\Gamma \in CC$  one has

1. for all  $T \in \mathcal{O}b_2(\Gamma)$  one has

$$[2016.09.13.eq1] H(\Pi_\Gamma(T)) = \Pi'_{H(\Gamma)}(H(T)) \quad (4.28)$$

2. for all  $o \in \widetilde{\mathcal{O}}b_2(\Gamma)$  one has

$$[2016.09.13.eq2] H(\lambda_\Gamma(o)) = \lambda'_{H(\Gamma)}(H(o)) \quad (4.29)$$

**Theorem 4.22** [2015.03.21.th1] *Let  $(\Phi, \phi, \tilde{\phi})$  be as above and let  $(P, \tilde{P}), (P', \tilde{P}')$  be as in Problem 3.3 for  $\mathcal{C}$  and  $\mathcal{C}'$  respectively.*

*Assume that the squares*

$$\begin{array}{ccc}
 \Phi(I_p(U)) & \xrightarrow{\Phi(P)} & \Phi(U) & & \Phi(I_p(\tilde{U})) & \xrightarrow{\Phi(\tilde{P})} & \Phi(\tilde{U}) \\
 \text{[2015.03.23.sq1]} \quad \xi_{\Phi} \downarrow & & \downarrow \phi & & \tilde{\xi}_{\Phi} \downarrow & & \downarrow \tilde{\phi} \\
 I_{p'}(U') & \xrightarrow{P'} & U' & & I_{p'}(\tilde{U}') & \xrightarrow{\tilde{P}'} & \tilde{U}'
 \end{array} \tag{4.30}$$

*commute. Then the homomorphism*

$$H(\Phi, \phi, \tilde{\phi}) : CC(\mathcal{C}, p) \rightarrow CC(\mathcal{C}', p')$$

*is a homomorphism of  $\mathcal{C}$ -systems with  $(\Pi, \lambda)$ -structures.*

**Proof:** We have to show that for all  $\Gamma \in Ob(CC(\mathcal{C}, p))$ ,  $T \in Ob_2(\Gamma)$  and  $o \in \tilde{Ob}_2(\Gamma)$  the equalities (4.28) and (4.29) hold. We will prove the first equality. The proof of the second is strictly parallel to the proof of the first.

By definition we have:

$$\begin{aligned}
 H(\Pi(T)) &= H(u_1^{-1}(\eta(u_2(T)) \circ P)) = (u_1)^{-1}(\psi(\Gamma) \circ \Phi(\eta(u_2(T)) \circ P) \circ \phi) = \\
 & \quad (u_1)^{-1}(\psi(\Gamma) \circ \Phi(\eta(u_2(T))) \circ \Phi(P) \circ \phi)
 \end{aligned}$$

where the second equality holds by Lemma 4.16(1) and

$$\Pi'(H(T)) = (u_1)^{-1}(u_2(H(T)) \circ P') = (u_1)^{-1}(\eta'(u_2(H(T))) \circ P')$$

Let us show that

$$\eta'(u_2(H(T))) \circ P' = \psi(\Gamma) \circ \Phi(\eta(u_2(T))) \circ \Phi(P) \circ \phi$$

By Lemma 4.20(1) we have

$$\eta'(u_2(H(T))) \circ P' = \psi(\Gamma) \circ \Phi(\eta(u_2(T))) \circ \xi_{\Phi} \circ P'$$

It remains to show that

$$\xi_{\Phi} \circ P' = \Phi(P) \circ \phi$$

which is our assumption about the commutativity of the square first square in (4.30).

## 5 Appendices

The facts discussed and proved in the following appendices are certainly well known. We had to repeat them here because we need to fix notations and because there is a number of facts whose proves I could not find in the literature.

## 5.1 Appendix A. Categories with binary products and binary cartesian closed categories

Let  $\mathcal{C}$  be a category.

**Definition 5.1** [2016.12.02.def1] *A binary product diagram is a pair of morphisms of the form  $(pr_1 : bp \rightarrow X, pr_2 : bp \rightarrow Y)$  such that for all  $A \in \mathcal{C}$  the function*

$$[2016.12.02.eq2a] \text{Mor}_{\mathcal{C}}(A, bp) \rightarrow \text{Mor}_{\mathcal{C}}(A, X) \times \text{Mor}_{\mathcal{C}}(A, Y) \quad (5.1)$$

*given by  $a \mapsto (a \circ pr_1, a \circ pr_2)$  is a bijection.*

*The structure of binary products on  $\mathcal{C}$  is a family, parametrized by pairs of objects  $(X, Y) \in \mathcal{C} \times \mathcal{C}$ , of binary product diagrams  $(pr_1(X, Y) : bp(X, Y) \rightarrow X, pr_2(X, Y) : bp(X, Y) \rightarrow Y)$ .*

Unless another notation is given, as for the binary products in the slice categories considered below, the object  $bp(X, Y)$  is denoted by  $X \times Y$  and the structural morphisms from  $X \times Y$  to  $X$  and  $Y$  by  $pr_1^{X, Y}$  and  $pr_2^{X, Y}$  respectively. We will often abbreviate the notation  $pr_i^{X, Y}$  to  $pr_i$ .

The following lemma expresses the well know “uniqueness” property of the binary products. We need its explicit form because in the next lemma we will need to state and prove that the corresponding “canonical” isomorphisms are natural.

**Lemma 5.2** [2016.12.02.11] *Let  $(pr_{1,i} : bp_i \rightarrow X, pr_{2,i} : bp_i \rightarrow Y)$ , where  $i = 1, 2$ , be two binary product diagrams. Let  $\iota_{1,2} : bp_1 \rightarrow bp_2$  be the morphism such that  $\iota_{1,2} \circ pr_{1,2} = pr_{1,1}$  and  $\iota_{1,2} \circ pr_{2,2} = pr_{2,1}$  and  $\iota_{2,1} : bp_2 \rightarrow bp_1$  be the morphism given by the symmetric condition. Then  $\iota_{1,2}$  and  $\iota_{2,1}$  are mutually inverse isomorphisms.*

**Proof:** To show that  $\iota_{1,2} \circ \iota_{2,1} = Id_{bp_1}$  we need to compare two morphisms whose codomain is a binary product. To do it it is sufficient, because of the injectivity of (5.1), to prove that their compositions with the two projections are equal. This follows by simple rewriting. The same applies to the second composition.

**Lemma 5.3** [2015.04.16.11] *Let  $\mathcal{C}$  be a category. Consider four binary product diagrams  $(pr_{1,i} : bp_i \rightarrow X, pr_{2,i} : bp_i \rightarrow Y)$  and  $(pr'_{1,i} : bp'_i \rightarrow X', pr'_{2,i} : bp'_i \rightarrow Y')$  where  $i = 1, 2$ . Let  $\iota = \iota_{1,2} : pb_1 \rightarrow pb_2$  be as in Lemma 5.2 and similarly  $\iota' : pb'_1 \rightarrow pb'_2$ . Let  $a : X' \rightarrow X$  and  $b : Y' \rightarrow Y$ .*

*Let  $c_i(a, b) : pb'_i \rightarrow pb_i$  be the unique morphisms such that  $c_i(a, b) \circ pr_{1,i} = pr'_{1,i} \circ a$  and  $c_i(a, b) \circ pr_{2,i} = b \circ pr'_{2,i}$ . Then the square*

$$\begin{array}{ccc} pb'_1 & \xrightarrow{c_1(a,b)} & pb_1 \\ \iota' \downarrow & & \downarrow \iota \\ pb'_2 & \xrightarrow{c_2(a,b)} & pb_2 \end{array}$$

*commutes, i.e.,  $c_1(a, b) \circ \iota = \iota' \circ c_2(a, b)$ .*



**Proof:** Since  $(pr_{1,2}, pr_{2,2})$  is a binary product digagram it is sufficient to prove that

$$c_1(a, b) \circ \iota \circ pr_{1,2} = \iota' \circ c_2(a, b) \circ pr_{1,2}$$

and

$$c_1(a, b) \circ \iota \circ pr_{2,2} = \iota' \circ c_2(a, b) \circ pr_{2,2}$$

For the first one we have:

$$c_1(a, b) \circ \iota \circ pr_{1,2} = c_1(a, b) \circ pr_{1,1} = pr'_{1,1} \circ a$$

and

$$\iota' \circ c_2(a, b) \circ pr_{1,2} = \iota' \circ pr'_{1,2} \circ a = pr'_{1,1} \circ a$$

The verification of the second equality is similar.

Given a category with binary products and morphisms  $a : X \rightarrow X'$ ,  $b : Y \rightarrow Y'$  denote by  $a \times b : X \times Y \rightarrow X' \times Y'$  the unique morphism such that  $(a \times b) \circ pr_1 = pr_1 \circ a$  and  $(a \times b) \circ pr_2 = pr_2 \circ b$ .

One has

$$[\mathbf{2016.11.26.eq1}] Id_{X \times Y} = Id_X \times Id_Y \tag{5.2}$$

and for  $a, b$  as above and  $a' : X' \rightarrow X''$ ,  $b' : Y' \rightarrow Y''$  one has

$$[\mathbf{2016.11.26.eq2}] (a \times b) \circ (a' \times b') = (a \circ a') \times (b \circ b') \tag{5.3}$$

One proves these two equalities by composing both sides with  $pr_1$  and  $pr_2$  and using the uniqueness part of the binary product axiom.

From (5.3) one derives

$$[\mathbf{2016.11.28.eq3}] (a \times Id_Y) \circ (a' \times Id_Y) = (a \circ a') \times Id_Y \tag{5.4}$$

and

$$[\mathbf{2016.11.28.eq4}] (Id_X \times b) \circ (Id_X \times b') = Id_X \times (b \circ b') \tag{5.5}$$

The definition of a binary cartesian closed structure given below differs slightly from the definition of the cartesian closed structure given in [?, IV.6] in that, that we do not require the specification of a finite object but only of binary products. The rest of the definition is identical to the one in [?], but written more explicitly in order to introduce the notations that are used in proofs in the main part of the paper.

Since we never use the definition of [?, IV.6] we will often write “cartesian closed” instead of “binary cartesian closed”.

**Definition 5.4** *[2016.11.28.def1] The (binary) cartesian closed structure on a category  $\mathcal{C}$  is a collection of data of the form:*

1. the structure of a category with binary products on  $\mathcal{C}$ ,
2. for all  $X, Y \in \mathcal{C}$  an object  $\underline{Hom}(X, Y)$ ,
3. for all  $X$  and  $b : Y \rightarrow Y'$  a morphism

$$\underline{Hom}(X, b) : \underline{Hom}(X, Y) \rightarrow \underline{Hom}(X, Y')$$

such that for all  $Y$  one has

$$\underline{Hom}(X, Id_Y) = Id_{\underline{Hom}(X, Y)}$$

and for all  $b : Y \rightarrow Y'$ ,  $b' : Y' \rightarrow Y''$  one has

$$\underline{Hom}(X, b \circ b') = \underline{Hom}(X, b) \circ \underline{Hom}(X, b')$$

4. For all  $X, Y$  a morphism

$$ev_Y^X : \underline{Hom}(X, Y) \times X \rightarrow Y$$

such that for all  $W$  the function

$$adj_Y^{W, X} : Mor(W, \underline{Hom}(X, Y)) \rightarrow Mor(W \times X, Y)$$

given by

$$[\mathbf{2016.11.28.eq2}] u \mapsto (u \times Id_X) \circ ev_Y^X \tag{5.6}$$

is a bijection and such that for all  $b : Y \rightarrow Y'$  the square

$$\begin{array}{ccc}
 \underline{Hom}(X, Y) \times X & \xrightarrow{ev_Y^X} & Y \\
 [\mathbf{2016.11.28.eq1}]_{(X, b) \times Id_X} \downarrow & & \downarrow b \\
 \underline{Hom}(X, Y') \times X & \xrightarrow{ev_{Y'}^X} & Y'
 \end{array} \tag{5.7}$$

commutes.

A cartesian closed category is a category together with a cartesian closed structure on it.

By definition the objects  $\underline{Hom}(X, Y)$  are functorial only in  $Y$ . Their functoriality in  $X$  is a consequence of a lemma. For  $X, X', Y$  and  $a : X \rightarrow X'$  let

$$\underline{Hom}(a, Y) : \underline{Hom}(X', Y) \rightarrow \underline{Hom}(X, Y)$$

be the unique morphism such that

$$[\mathbf{2016.11.28.eq5}] adj(\underline{Hom}(a, Y)) = (Id_{\underline{Hom}(X', Y)} \times a) \circ ev_Y^X \tag{5.8}$$

Then one has:

**Lemma 5.5** [2015.04.10.11] *The morphisms  $\underline{Hom}(-, Y)$  satisfy the equalities*

$$\underline{Hom}(a \circ a', Y) = \underline{Hom}(a', Y) \circ \underline{Hom}(a, Y)$$

$$\underline{Hom}(Id_X, Y) = Id_{\underline{Hom}(X, Y)}$$

*making  $\underline{Hom}(-, Y)$  into a contravariant functor from  $\mathcal{C}$  to itself.*

*In addition, for all  $b : Y \rightarrow Y'$  the square*

$$\begin{array}{ccc} \underline{Hom}(X', Y) & \xrightarrow{\underline{Hom}(X, b)} & \underline{Hom}(X', Y') \\ \underline{Hom}(a, Y) \downarrow & & \downarrow \underline{Hom}(a, Y') \\ \underline{Hom}(X, Y) & \xrightarrow{\underline{Hom}(X', b)} & \underline{Hom}(X, Y') \end{array}$$

*commutes.*

**Proof:** It is a particular case of [?, Theorem 3, p.100]. The commutativity of the square is a part of the "bifunctor" claim of the theorem.

**Lemma 5.6** [2015.04.20.12] *In a cartesian closed category let  $X, X', Y$  be objects and let  $a : X \rightarrow X'$  be a morphism. Then the square*

$$\begin{array}{ccc} \underline{Hom}(X', Y) \times X & \xrightarrow{Id_{\underline{Hom}(X', Y)} \times a} & \underline{Hom}(X', Y) \times X' \\ \underline{Hom}(a, Y) \times Id_X \downarrow & & \downarrow ev_Y^{X'} \\ \underline{Hom}(X, Y) \times X & \xrightarrow{ev_Y^X} & Y \end{array}$$

*commutes.*

**Proof:** Let us show that both paths in the square are adjoints to  $\underline{Hom}(a, Y)$ . For the path that goes through the upper right corner it follows from the definition of  $\underline{Hom}(a, Y)$  as the morphism whose adjoint is  $(Id \times a) \circ ev$ . For the path that goes through the lower left corner it follows from the definition of adjoint applied to  $\underline{Hom}(a, Y)$ . Indeed, the adjoint to this morphism is

$$adj(\underline{Hom}(a, Y)) = (\underline{Hom}(a, Y) \times Id_X) \circ ev_Y^X$$

**Lemma 5.7** [2015.05.12.12] *Let  $\mathcal{C}$  be a cartesian closed category. Let  $X, Y, W \in \mathcal{C}$ , then one has:*

1. *Let  $Y'$  be an object and  $b : Y \rightarrow Y'$  a morphism. Then for any  $r \in Mor(W, \underline{Hom}(X, Y'))$  one has*

$$adj(r \circ \underline{Hom}(X, b)) = adj(r) \circ b$$

2. *Let  $X'$  be an object  $a : X \rightarrow X'$  a morphism. Then for any  $r \in Mor(W, \underline{Hom}(X', Y))$  one has*

$$adj(r \circ \underline{Hom}(a, Y)) = (Id_W \times a) \circ adj(r)$$

3. Let  $W'$  be an object  $c : W \rightarrow W'$  a morphism. Then for any  $r \in \text{Mor}(W', \underline{\text{Hom}}(X, Y))$  one has

$$\text{adj}(c \circ r) = (c \times \text{Id}_X) \circ \text{adj}(r)$$

**Proof:** The proof of the first case is given by

$$\begin{aligned} \text{adj}(r \circ \underline{\text{Hom}}(X, b)) &= ((r \circ \underline{\text{Hom}}(X, b)) \times \text{Id}_X) \circ \text{ev}_Y^X = \\ &= (r \times \text{Id}_X) \circ (\underline{\text{Hom}}(X, b)) \times \text{Id}_X \circ \text{ev}_Y^X = \\ &= (r \times \text{Id}_X) \circ \text{ev}_Y^X \circ b = \text{adj}(r) \circ b \end{aligned}$$

where the first equality is by (5.6), second equality by Lemma 5.9, the third equality by the commutativity of (5.7) and the fourth equality again by (5.6).

The proof of the second case is given by the following sequence of equalities where we use the notation  $Hm$  for  $\underline{\text{Hom}}(a, Y)$  as well as a number of other abbreviations:

$$\begin{aligned} \text{adj}(r \circ Hm) &= ((r \circ Hm) \times \text{Id}) \circ \text{ev} = (r \times \text{Id}) \circ (Hm \times \text{Id}) \circ \text{ev} = (r \times \text{Id}) \circ \text{adj}(Hm) = \\ &= (r \times \text{Id}) \circ (\text{Id} \times a) \circ \text{ev} = (r \times a) \circ \text{ev} = (\text{Id} \times a) \circ (r \times \text{Id}) \circ \text{ev} = (\text{Id} \times a) \circ \text{adj}(r) \end{aligned}$$

where the first equality is by (5.6), the second by (5.4), the third by (5.6), the fourth by (5.8), the fifth by (5.3), the sixth by (5.3) and the seventh by (5.6).

The proof of the third case is given by

$$\begin{aligned} \text{adj}(c \circ r) &= ((c \circ r) \times \text{Id}_X) \circ \text{ev}_Y^X = (c \times \text{Id}_X) \circ (r \times \text{Id}_X) \circ \text{ev}_Y^X = \\ &= (c \times \text{Id}_X) \circ \text{adj}(r) \end{aligned}$$

where the first equality is by (5.6), second equality by (5.4) and the third equality by (5.6). Lemma is proved.

## 5.2 Appendix B. Slice categories, pullbacks and locally cartesian closed categories

For a category  $\mathcal{C}$  and  $Z \in \mathcal{C}$  one denotes by  $\mathcal{C}/Z$  the slice category of  $\mathcal{C}$  over  $Z$ . When one works in the set theory one has to choose one of the several possible definitions of  $\mathcal{C}/Z$ . Indeed, the set of objects of  $\mathcal{C}/Z$  can be defined as the set of pairs  $(X, f)$  where  $X \in \mathcal{C}$  and  $f : X \rightarrow Z$  or as the set of morphisms  $f \in \text{Mor}(\mathcal{C})$  such that  $\text{codom}(f) = Z$ . There is an obvious bijection between these two sets but they are not equal. We define  $\text{Ob}(\mathcal{C}/Z)$  as the set of pairs  $(X, f)$ . Even more choices exist in the definition of the set of morphisms of  $\mathcal{C}/Z$ . One definition is the set of triples  $((X, f), (Y, g), a)$  where  $(X, f), (Y, g) \in \text{Ob}(\mathcal{C}/Z)$  and  $a : X \rightarrow Y$  is such that  $f = a \circ g$ . Another one is the set of pairs  $(a, g)$  where  $a, g \in \text{Mor}(\mathcal{C})$  are such that  $\text{codom}(a) = \text{dom}(g)$  and  $\text{codom}(f) = Z$ . Again, these sets are obviously isomorphic but not equal. Various other choices are possible. We will use the second option. We denote the pair  $(a, g)$  by  $a^g$ .

The mappings  $(X, f) \mapsto X$  and  $a^g \mapsto a$  define a functor  $\mathcal{C}/Z \rightarrow \mathcal{C}$  that we denote by  $\pi_{Z,\#}$ . We will rarely write the functions  $(\pi_{Z,\#})_{Ob}$  and  $(\pi_{Z,\#})_{Mor}$  explicitly using them instead as “coercions”. Formally speaking, we will assume that  $(\pi_{Z,\#})_{Ob}$  (resp.  $(\pi_{Z,\#})_{Mor}$ ) is inserted in our notation whenever an object (resp. a morphism) of  $\mathcal{C}/Z$  is specified where an object (resp. a morphism) of  $\mathcal{C}$  is required.

We will say that  $a : X \rightarrow Y$  is a morphism over  $Z$  if  $a \circ g = f$ . For given  $(X, f)$  and  $(Y, g)$ , the function

$$[\mathbf{2016.11.26.eq3}] a^g \mapsto a \tag{5.9}$$

defines a bijection between morphisms  $(X, f) \rightarrow (Y, g)$  in  $\mathcal{C}/Z$  and morphisms  $X \rightarrow Y$  over  $Z$  in  $\mathcal{C}$ .

In a category with binary products the morphism  $Id_Z \times b$  satisfies the equality

$$(Id_Z \times b) \circ pr_1 = pr_1$$

and therefore defines a morphism from  $(Z \times Y, pr_1)$  to  $(Z \times Y', pr_1)$  in  $\mathcal{C}/Z$ . We will denote this morphism in the slice category by  $Z \times b$ . Since (5.9) is injective, the equalities (5.2) and (5.3) imply that

$$[\mathbf{2016.11.30.eq1}] Z \times Id_Y = Id_{(Z \times Y, pr_1)} \tag{5.10}$$

and

$$[\mathbf{2016.11.30.eq2}] Z \times (b \circ b') = (Z \times b) \circ (Z \times b') \tag{5.11}$$

that is, that the mappings  $X \mapsto (Z \times Y, pr_1)$ ,  $b \mapsto Z \times b$  define a functor  $Z \times -$  from  $\mathcal{C}$  to  $\mathcal{C}/Z$ .

The same holds for morphisms of the form  $a : X \rightarrow X'$ . We denote the morphism in  $\mathcal{C}/Z$  corresponding to the morphism  $a \times Id_Z$  by  $a \times Z$  and the resulting functor  $\mathcal{C} \rightarrow \mathcal{C}/Z$  by  $- \times Z$ .

**Lemma 5.8** *[2016.12.16.11]* *Let*

$$[\mathbf{2016.12.16.eq1}] \begin{array}{ccc} X & \xrightarrow{a} & Y \\ \downarrow & & \downarrow g \\ Y' & \xrightarrow{g'} & Z \end{array} \tag{5.12}$$

*be a commutative square of morphisms in  $\mathcal{C}$  and  $f = a \circ g = a' \circ g'$ . Then*

$$[\mathbf{2016.12.16.eq2}] \begin{array}{ccc} (X, f) & \xrightarrow{a^g} & (Y, g) \\ \downarrow (a')^{g'} & & \\ (Y, g') & & \end{array} \tag{5.13}$$

*is a binary product diagram in  $\mathcal{C}/Z$  if and only if (5.12) is a pullback in  $\mathcal{C}$ .*

**Proof:** Assume that (5.13) is a binary product diagram. Let  $W \in \mathcal{C}$  and let  $d : W \rightarrow Y$ ,  $d' : W \rightarrow Y'$  be such that  $d \circ g = d' \circ g'$ . Let  $e = d \circ g$ . Then  $d^g : (W, e) \rightarrow (Y, g)$  and  $(d')^{g'} : (W, e) \rightarrow (Y', g')$  are morphisms in  $\mathcal{C}/Z$  and therefore there exists  $c^f : (W, e) \rightarrow (X, f)$  such that  $c^f \circ a^g = d^g$  and  $c^f \circ (a')^{g'} = (d')^{g'}$  in  $\mathcal{C}/Z$ , that is,  $c \circ a = d$  and  $c \circ a' = d'$  in  $\mathcal{C}$ . Let  $c' : W \rightarrow X$  be another morphism in  $\mathcal{C}$  such that  $c' \circ a = d$  and  $c' \circ a' = d'$ . Then  $e = d \circ g = c' \circ a \circ g = c' \circ f$  and therefore  $(c')^f$  is a morphism  $(W, e) \rightarrow (X, f)$  in  $\mathcal{C}/Z$ . Next,  $(c')^f \circ a^g = (c' \circ a)^g = d^g$  and  $(c')^f \circ (a')^{g'} = (c' \circ a')^{g'} = (d')^{g'}$ . Therefore  $(c')^f = c^f$ , that is,  $c = c'$ . This shows that (5.12) is a pullback in  $\mathcal{C}$ .

Similar reasoning shows that if (5.12) is a pullback in  $\mathcal{C}$  then (5.13) is a binary product diagram in  $\mathcal{C}/Z$ .

Lemma 5.8, combined with a related statement about commutative squares, implies that a choice of binary product structures on all the slice categories  $\mathcal{C}/Z$  is “the same as” the choice of pullbacks for all pairs of morphisms with the common codomain in  $\mathcal{C}$ .

To be precise we have to say that how to construct a bijection between the set of families of binary product structures on the categories  $\mathcal{C}/Z$  for all  $Z$  and the set of pullback structures on  $\mathcal{C}$ .

We usually denote the distinguished binary product of  $(X, f)$  and  $(Y, g)$  in  $\mathcal{C}/Z$  by  $(X, f) \times_Z (Y, g)$  and the canonical morphism from  $(X, f) \times_Z (Y, g)$  to  $Z$  by  $f \diamond g$ .

For  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$ , the two commutative triangles formed by  $pr_1 : (X, f) \times_Z (Y, g) \rightarrow (X, f)$ ,  $f$ ,  $f \diamond g$  and  $pr_2 : (X, f) \times_Z (Y, g) \rightarrow (Y, g)$ ,  $g$ ,  $f \diamond g$  are adjacent and define the familiar commutative square of the pullback of  $f$  and  $g$ .

This defines a function in one direction.

For  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$ , the diagonal of the pullback square based on  $f$  and  $g$  is an object over  $Z$  and the two projections define morphisms from this object to  $(X, f)$  and  $(Y, g)$  respectively. The corresponding pair of morphisms in  $\mathcal{C}/Z$  is a binary product diagram. This defines a morphism in the other direction.

The fact that these morphisms are inverse to each other follows readily from the construction.

Given a binary products structure on  $\mathcal{C}/Z$ , morphisms  $f : X \rightarrow Z$ ,  $g : Y \rightarrow Z$  and morphisms  $a : X' \rightarrow X$ ,  $b : Y' \rightarrow Y$  we have a morphism  $a^f \times_Z b^g$  which is the unique morphism in  $\mathcal{C}/Z$  of the form

$$a^f \times_Z b^g : (X', a \circ f) \times_Z (Y', b \circ g) \rightarrow (X, f) \times_Z (Y, g)$$

such that

$$[\mathbf{2016.11.24.eq1}](a^f \times_Z b^g) \circ pr_1 = pr_1 \circ a^f \tag{5.14}$$

and

$$[\mathbf{2016.11.24.eq2}](a^f \times_Z b^g) \circ pr_2 = pr_2 \circ b^g \tag{5.15}$$

**Lemma 5.9** [2015.05.14.11] *In the setting introduced above one has:*

$$1. \text{Id}_{(X,f) \times_Z (Y,g)} = \text{Id}_{(X,f)} \times_Z \text{Id}_{(Y,g)},$$

2. suppose that we have in addition morphisms  $a' : X'' \rightarrow X'$  and  $b' : Y'' \rightarrow Y'$ . Then

$$((a')^{a \circ f} \times_Z (b')^{b \circ g}) \circ (a^f \times_Z b^g) = (a' \circ a)^f \times_Z (b' \circ b)^g$$

**Proof:** It is a particular case of (5.2) and (5.3).

Following the general case considered in Appendix 5.1 we will write  $(X, f) \times_Z b^g$  (resp.  $a^f \times_Z (Y, g)$ ) for the morphism in  $\mathcal{C}/X$  (resp.  $\mathcal{C}/Y$ ) corresponding to  $\text{Id}_{(X,f)} \times_Z b^g$  (resp.  $a^f \times_Z \text{Id}_{(Y,g)}$ ).

In view of Lemma 5.9 and (5.14), for any  $(X, f : X \rightarrow Z)$ , the functions

$$(Y, g) \mapsto ((X, f) \times_Z (Y, g), pr_1)$$

$$(b^g : (Y', g') \rightarrow (Y, g)) \mapsto (X, f) \times_Z b^g$$

form a functor from  $\mathcal{C}/Z$  to  $\mathcal{C}/X$  and similarly by Lemma 5.9 and (5.15), for any  $(Y, g : Y \rightarrow Z)$  the functions

$$(X, f) \mapsto ((X, f) \times_Z (Y, g), pr_2)$$

$$(a^f : (X', f') \rightarrow (X, f)) \mapsto a^f \times_Z (Y, g)^f$$

form a functor from  $\mathcal{C}/Z$  to  $\mathcal{C}/Y$ .

**Definition 5.10** [2015.03.27.def1] *A locally cartesian closed structure on a category  $\mathcal{C}$  is a family of (binary) cartesian closed structures on the categories  $\mathcal{C}/Z$  for all  $Z \in \mathcal{C}$ .*

*We usually denote the binary product on  $\mathcal{C}/Z$  as above.*

*We usually denote the internal-hom objects in  $\mathcal{C}/Z$  by  $\underline{\text{Hom}}_Z((X, f), (Y, g))$  and the canonical morphisms from  $\underline{\text{Hom}}_Z((X, f), (Y, g))$  to  $Z$  by  $f \Delta g$ .*

*The rest of the notations ( $\underline{\text{Hom}}_Z((X, f), b^g)$ ,  $ev_{(Y,g)}^{(X,f)}$ ,  $adj_{(Y,g)}^{(W,h),(X,f)}$ ,  $\underline{\text{Hom}}_Z(a^f, (Y, g))$ ) immediately follow from the ones introduced previously.*

*A locally cartesian closed category is a category together with a locally cartesian closed structure on it.*

The name “locally cartesian closed” follows naturally from this definition and the intuition based on the example of the category of open sets of a topological space or a Grothendieck site. If only the subsets of the open sets of a particular covering are known then one sometimes says that the space is known only locally, but the global structure that arises from gluing of all these subsets together is not known. Hence the “local” structure of a category is given by the structure of its slice categories.

**Example 5.11** [2015.05.20.ex1] The following example shows that there can be many different structures of a category with pullbacks on a category and also many locally cartesian closed structures.

Let us take as our category the category  $F$  whose objects are natural numbers and

$$Mor(n, m) = Fun(\{0, \dots, n - 1\}, \{0, \dots, m - 1\})$$

Since every isomorphism class contains exactly one object every auto-equivalence of this category is an automorphism. Let  $\Phi$  be such an automorphism. It is easy to see that it must be identity on the set of objects. Let  $X = \{0, 1\}$ . Consider  $\Phi$  on  $End(X)$ . Since  $\Phi$  must respect identities and compositions,  $\Phi$  must take  $Aut(X)$  to itself and must act on it by identity. If  $1$  and  $\sigma$  are the two elements of  $Aut(X)$  we conclude that  $\Phi(1) = 1$  and  $\Phi(\sigma) = \sigma$ .

Let us choose now any structure of a category with pullbacks on  $F$  and let us consider two new structures  $str_1$  and  $str_\sigma$  that are obtained by modifying pullbacks as follows. In both structures we set all pullbacks to be as they were except for the pullback of the pair of morphisms  $(Id_X, Id_X)$ . For this pair we set the pullbacks to be as follows:

$$\begin{array}{ccc}
 X & \xrightarrow{Id_X} & X & & X & \xrightarrow{\sigma} & X \\
 \text{[2015.05.20.sq1]} \downarrow & & \downarrow Id_X & \text{for } str_1 \text{ and} & \sigma \downarrow & & \downarrow Id_X & \text{for } str_\sigma. & (5.16) \\
 X & \xrightarrow{Id_X} & X & & X & \xrightarrow{Id_X} & X
 \end{array}$$

The preceding discussion of the auto-equivalences of  $F$  shows that there is no auto-equivalence which would transform  $str_1$  into  $str_\sigma$ .

The category  $F$  also has a locally cartesian closed structure and it can be shown that it can be modified so that its pullback components are  $str_1$  and  $str_\sigma$ . This shows that  $F$  has at least two locally cartesian closed structures that are not equivalent modulo the auto-equivalences of  $F$ .

The solution to this seeming paradox is that there is a category structure on the set of pullback structures (resp. locally cartesian closed structures) on a category. Any two pullback structures (resp. lcc structures) are isomorphic in this category and in this sense pullbacks on a category are “unique”.

**Remark 5.12** [2015.05.20.rem1] The previous example has a continuation in the univalent foundations where there is a notion of a category and pre-category. There the types of pullback structures and of locally cartesian closed structures on a category (as opposed to those on a general pre-category) are of h-level 1, i.e., classically speaking are either empty or contain only one element.

In addition any such structure on a pre-category should define a structure of the same kind on the Rezk completion of this pre-category with all the different structures on the pre-category becoming equal on the Rezk completion. In the case of the previous example the Rezk completion of  $F$  is the category  $FSets$  of finite sets and in view of the univalence axiom for finite sets the two pullbacks of 5.16 will become equal in  $FSets$ .



## 6 Acknowledgements

I am grateful to the Department of Computer Science and Engineering of the University of Gothenburg and Chalmers University of Technology for its the hospitality during my work on the paper.

Work on this paper was supported by NSF grant 1100938.