

THE GEOMETRY OF $C^{1,\alpha}$ FLAT ISOMETRIC IMMERSIONS

CAMILLO DE LELLIS AND MOHAMMAD REZA PAKZAD

ABSTRACT. We show that any isometric immersion of a flat plane domain into \mathbb{R}^3 is developable provided it enjoys $C_{loc}^{1,\alpha}$ Hölder regularity for some $\alpha > 2/3$. The proof is based on showing the existence of a weak notion of second fundamental form for such immersions, the analysis of the Gauss-Codazzi-Mainardi equations in this weak setting, and a parallel result on the very weak solutions to the Monge-Ampère equation which appeared in [43].

1. INTRODUCTION

1.1. Background. It often happens, in differential geometry, as in other contexts where nonlinear PDEs appear, that the solutions to a given nonlinear system of equations satisfy also systems of higher order equations carrying significant geometric or analytic information. A very basic example is the Gauss-Codazzi-Mainardi equations satisfied by any smooth isometric immersion of a 2 dimensional Riemannian manifold into \mathbb{R}^3 . Note that, in their turn, these equations can be used to prove certain rigidity statements for the immersion, e.g. to show that any isometric immersion of a flat domain is developable, i.e. roughly speaking, at any given point p its image contains a segment of \mathbb{R}^3 passing through p .

The subtlety which is often glossed over in the context of classical differential geometry, where all the mappings are assumed smooth, or at least twice or thrice continuously differentiable, is that, due to nonlinearity, the passage from say a given set of first order equations (e.g. the system of isometric immersion equations in our example), to the higher order equations requires a minimum of regularity. There is no guarantee that in the absence of this regularity the geometric information hidden in the higher order equations would be accessible. The above mentioned subtlety could be a source of confusion. Coming back to our example, it is often assumed that all isometric images of flat domains must be ruled, as the authors have recently noticed implicitly stated in a Wikipedia post on developable surfaces. The latter statement is untrue. Already, at the dawn of modern differential geometry, no less figures than Lebesgue and Picard intuited the existence of a surface which could be flattened onto the plane with no distortion of relative distances, but nowhere contained a straight segment. To quote Picard¹:

According to general practice, we suppose in the preceding analysis, as in all infinitesimal geometry of curves and surfaces, the existence of derivatives which we need in the calculus. It may seem premature to entertain a theory of surfaces in which one does not make such hypotheses. However, a curious result has been pointed out by Mr Lebesgue (Comptes Rendus, 1899 and thesis)²; according to which one may, by the aid of continuous functions, obtain surfaces corresponding to a plane, of such sort that every rectifiable line of the plane has a corresponding rectifiable line of the same length of the surface, nevertheless the surfaces obtained are no

2010 *Mathematics Subject Classification.* 53C24, 53C21, 57N35, 53A05.

Key words and phrases. Nash-Kuiper, rigidity of isometric immersions, differential geometry at low regularity.

¹[51, Page 555], translated and quoted in [6].

²[40]. See also [20, Pages 164-165] and the discussion therein of Lebesgue's heuristic construction.

longer ruled. If one takes a sheet of paper, and crumples it by hand, one obtains a surface applicable to the plane and made up of a finite number of pieces of developable surfaces, joined two and two by lines, along which they form a certain angle. If one imagines that the pieces become infinitely small, the crumpling being pushed everywhere to the limit, one may arrive at the conception of surfaces applicable to the plane and yet not developable [in the sense that there is no envelope of a family of planes of one parameter and not ruled].

The suggested constructions by Lebesgue and then Picard fall short of being totally satisfactory as they point to continuous but yet not differentiable mappings. However, they were perhaps the heralds of a celebrated result due to Kuiper [39], who based on the work by John Nash [47, 48] proved that any Riemannian manifold can be C^1 isometrically embedded into a Riemannian manifold of one dimension higher. As a consequence, the flat disk can be C^1 -isometrically embedded into any given small ball of \mathbb{R}^3 and hence the image cannot be ruled. Actually, to make the matters worse, by Nash and Kuiper's construction, C^1 isometric embeddings can approximate any short immersion, and they can be constructed such that their images do not include any straight segment of \mathbb{R}^3 in any scale! In contrast, and returning to our original developability statement in a higher regularity setting where the Gauss-Codazzi equations hold true, Hartman and Nirenberg [27] proved, incidentally almost at the same time Nash and Kuiper showcased their results, that a C^2 -regular isometric immersion of a flat domain must have a locally ruled surface as its image. We have come full circle, and this discussed dichotomy leaves us with a fundamental question any analyst would like to ask: **At what regularity threshold or thresholds between C^1 and C^2 the unruled isometries transition into the ruled regime?**

To put the above picture in a broader context, we note that the the above dichotomy -known in the recent literature as *flexibility* vs. *rigidity*- is not specific to the case of isometric immersions. The Nash and Kuiper scheme for creating highly oscillatory anomalous solutions of typically lower regularity through iterations could be studied under the broader topic of convex integration [24, 53] for the differential inclusions (or PDEs re-cast in this framework), and the involved density or flexibility related results are usually referred to as h-principle. The existence of such h-principle is usually accompanied by a parallel rigidity property which indicates that the construction cannot be carried out in high regularity scales. A most famous example is the recent discovery by the first author and Székelyhidi [15]: They elegantly showed that the system of Euler equations in fluid dynamics is given to the convex integration method and that Nash and Kuiper's iterations can be adapted in this case in order to create non-conservative compactly supported continuous flows approximating an open set of subsolutions (see [54, 16] for a thorough discussion of the connection between the two problems of turbulence and isometric immersions). Their results, and the subsequent work in improving the regularity of the anomalous solutions stood in contrast with the rigidity result reflected in the conservation of energy for solutions passing a certain regularity threshold ($\alpha > 1/3$ for $C^{1,\alpha}$ -Hölder continuous solutions) in [10, 18]), as conjectured by Onsager [49]. Their approach finally lead to the complete resolution of this conjecture by Isett in [31]; see [17] for a review of the history of the problem and the intermediate results.

Coming back to the question of isometric immersions, a parallel question is whether the images of isometric immersions into \mathbb{R}^3 of elliptic 2d manifolds (i.e. 2-manifolds with positive Gaussian curvature) lie, locally, on convex 3d bodies. This convexity property fails for $C^{1,\alpha}$ isometric immersions if $\alpha < 1/5$, in which regime the above mentioned h-principle for isometric immersions holds true, as shown in [14], improving on the results by [5, 11]. On the other hand, Conti, De Lellis and Székelyhidi proved in [11] that the image must be convex in this setting when $\alpha > 2/3$. Mischa Gromov conjectures in [25, Section 3.5.5.C, Open Problem 34-36] that the transition threshold is $\alpha = 1/2$ in this case. The best evidence so far for this conjecture is provided in [13], where the

authors show that $C^{1,1/2}$ regularity could be the borderline regularity for the identity of intrinsic covariant derivative of the 2-manifold and its geometric interpretation.

This article concerns the fundamental question of the sub- C^2 regularity scales for which, given an isometric immersion of a flat domain, one is able to make sense of fundamentally geometric objects such as the second fundamental form and the Gauss-Codazzi equations. The final goal is to show that the developability property of such isometric immersions (as defined below in Definition 1 and proven for C^2 isometries in [27]) survives for $C^{1,\alpha}$ isometric immersions if $\alpha > 2/3$. As we shall see, parts of the project, e.g. the definition of the second fundamental form, can be carried over for $\alpha > 1/2$, but our analysis falls short of proving a weak variant of the Codazzi-Mainardi equations for $1/2 < \alpha < 2/3$. In this manner, the question of the optimal threshold remains open in the flat case as it is for the elliptic case.

Before proceeding, it should be mentioned that the developability of flat isometric immersions have been discussed in the literature in other regularity regimes. Pogorelev [52, Theorem 1, p. 603] shows this developability under the rather weak assumptions of C^1 regularity and the existence and vanishing of the total extrinsic curvature (defined as a measure). This result lies at the background of our conclusion through the analysis made in [43] and [37, 38]. On the other hand parallel results have been shown by the second author for $W^{2,2}$ isometric immersions [50], where a slight C^1 regularity gain in this case is also proven, and by Jerrard for the class of Monge-Ampère functions [34]. The former result, and the subsequent statement regarding the smooth density of isometric immersions were generalized to higher dimensional domains for the co-dimension 1 case in [44]. The latter approach by Jerrard [23, 33, 34], anchored in geometric measure theory, opened the path for proving C^1 regularity and a full range of developability results for isometric immersions of any dimension and co-dimension of sub-critical Sobolev regularity [35].

1.2. The main result. We first begin by a definition to clarify the notion of developability as we understand it.

Definition 1. Let $\Omega \subset \mathbb{R}^2$ be a domain. We say $u \in C^1_{loc}(\Omega, \mathbb{R}^k)$ is developable if for all $x \in \Omega$, either u is affine in a neighborhood of x , or there exists a line L_x containing x such that the Jacobian derivative ∇u is constant on l_x , the connected component of x in $L_x \cap \Omega$.

Remark 1.1. For bounded domains, l_x is always a segment whose both ends lie on $\partial\Omega$. We allow for unbounded domains, and hence l_x could be a complete line, a ray, or an open segment. With some abuse of notation, we will refer to all of them as “segments” throughout the paper.

There are other equivalent formulations for the developability whose equivalence we will show further on in Proposition 2.1. In particular, local and global developability, as laid out in Corollary 2.2, are equivalent. Also, from Lemma 2.8 and Corollary 2.7 it follows that when u is developable, the segments l_x are uniquely determined outside the maximal affine region; they do neither intersect each other nor pass through the local constancy regions of the Jacobian derivative.

In this article we prove:

Theorem 1. *Let $\alpha > 2/3$, and let $\Omega \subset \mathbb{R}^2$ be a domain. If $u \in C^{1,\alpha}_{loc}(\Omega, \mathbb{R}^3)$ is an isometric immersion, then u is developable.*

Per [52, Theorem 1, p. 603], what we need is to show is that the image of \vec{n} is of measure zero. Following [11, Corollary 5] it is possible to prove that the Brouwer degrees $\deg(y, V, \vec{n})$ of the Gauss map \vec{n} of the immersion vanish for all open sets $V \subset \Omega$ and $y \in \mathbb{R}^2$, when defined. However, contrary to the positive curvature case in [11], this is not sufficient to conclude with $|\vec{n}(\Omega)| = 0$: Through a similar approach as in [45, Section 5], for each $\alpha \in (0, 1)$, one can construct a map in $C^{0,\alpha}(\Omega, \mathbb{R}^2)$ whose local degree vanishes everywhere, but whose image is onto

the unit square. The zero degree Gauss map could still have an image of positive measure. This obstacle necessitates another strategy. Indeed, our argument uses the parallel result proved for the degenerate *very weak Monge-Ampère equation* in [43], which uses the degree formula for both the gradient of the solution, and its affine perturbations of positive degree.

The very weak Hessian determinant [32, 19] of a given function $v \in W_{loc}^{1,2}(\Omega)$ is defined to be

$$\mathcal{D}et D^2v := -\frac{1}{2} \left(\partial_{11}(\partial_2v)^2 + \partial_{22}(\partial_1v)^2 - 2\partial_{12}(\partial_1v\partial_2v) \right) = -\frac{1}{2} \text{curl}^T \text{curl} (\nabla v \otimes \nabla v)$$

in the sense of distributions. The operator $\mathcal{D}et D^2$ coincides with the usual Monge-Ampère operator $\det \nabla^2$ when $v \in C^2$ and ∇^2v stands for the Hessian matrix field of v . (A C^3 regularity is needed for a straightforward calculation). In [43], Lewicka and the second author proved:

Theorem 2. [43, Theorem 1.3] *Let $\Omega \subset \mathbb{R}^2$ be an open bounded domain, $2/3 < \alpha < 1$ and let $v \in C^{1,\alpha}(\bar{\Omega})$ be such that*

$$\mathcal{D}et D^2v = 0,$$

in Ω . Then v is developable.

Remark 1.2. Per Corollary 2.2 the assumption can be easily relaxed to the local $C^{1,\alpha}$ regularity for v in any bounded or unbounded domain.

The main strategy of our proof is hence to relate the given isometric immersion u to a scalar function v which satisfies the assumptions of Theorem 2: In a first step, we will show that a notion of the second fundamental form A in the weak sense exists for immersions of $C^{1,\alpha}$ regularity if $\alpha > 1/2$. Next, the Codazzi-Mainardi equations will be used to show that A must be curl free, which implies that it must be the Hessian matrix of a scalar function, namely the sought after function v . We will then prove the required regularity for v , and proceed to show using the Gauss equation that v satisfies the very weak degenerate Monge-Ampère equation as required by the assumptions of Theorem 2. Finally, we will need to prove that developability of v , as derived from Theorem 2, implies the developability of the isometric immersion u . The $\alpha > 2/3$ Hölder exponent assumption is only required for proving that a weak version of Codazzi-Mainardi equations hold true for u , i.e. when we need to show that A is curl free.

The article is organized as follows: In Section 2, we will present and prove a few statements regarding the developability properties of C^1 mappings. In Section 3, we will gather a few analytical tools which deal with properties of Hölder continuous functions and with quadratic differential expressions in terms of functions of low regularity. In particular, our basic proposition 3.2 will allow us to define a second fundamental form for the immersions of Hölder type regularity. The subsequent section is dedicated to the definition and properties of this second fundamental form. Finally, in Section 5, building on the previous sections, we provide a proof of Theorem 1.

1.3. Acknowledgments. This project was based upon work supported by the National Science Foundation and was partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation). M.R.P. was supported by the NSF award DMS-1813738 and by the DFG via SFB 1060 - Project Id 211504053. The second author would like to thank Stefan Müller for the opportunity to work at IAM, University of Bonn, Germany, where part of this work was completed, and for his kind support.

2. PRELIMINARIES: DEVELOPABLE MAPPINGS

We have gathered in this section a few statements we will need, and their proofs, regarding the developable mappings in two dimensions. Most of the material in this section are well known and can be found in one form or another in the literature on the topic [52, 9, 36, 55, 50, 46, 37, 38, 28,

29, 34, 44, 35, 43], but the authors do not know of any instance where the following statements are explicitly formulated. Note that, as already observed in [28], much of the developability properties of a mapping concern the level sets of its Jacobian derivative, so we formulate our statements having the mappings $f = \nabla u$ in mind, where u is a developable mapping.

2.1. Global vs. local developability. Developability can be defined by local or global formulations, which turn out to be equivalent. There is a risk of confusion, which must be avoided, due to slight differences between the possible formulations (see e.g. the footnote on p. 875 in [38]). Also, it would be useful to have a set of equivalent statements at hand to streamline the arguments. The following proposition states three equivalent conditions which the Jacobian derivative of a C^1 mapping can satisfy to be developable:

Proposition 2.1. *Let $\Omega \subset \mathbb{R}^2$ be any domain and let $f : \Omega \rightarrow \mathbb{R}^m$. If f is continuous, then the following three conditions are equivalent.*

- (a) *For all $x \in \Omega$, either f is constant in a neighborhood of x , or there exists a line L containing x such that f is constant on the connected component of x in $L \cap \Omega$.*
- (b) *For all $x \in \Omega$, there exists an open disk $B_x \subset \Omega$ centered at x with the following property: To all $p \in B_x$, we can associate a line L_p containing p such that f is constant on the segment $\tilde{l}_p = L_p \cap B_x$. Moreover, for any $y, z \in B_x$, $\tilde{l}_y \cap \tilde{l}_z = \emptyset$, or $\tilde{l}_y = \tilde{l}_z$.*
- (c) *For all $x \in \Omega$, there exists an open disk B_x centered at x in Ω such that for all $p \in B_x$, there exists a line $L \subset \mathbb{R}^2$ containing p so that f is constant on the segment $L \cap B_x$.*

Corollary 2.2. *$u \in C^1_{loc}(\Omega, \mathbb{R}^k)$ is developable if and only if for any point in $x \in \Omega$, there exists a neighborhood V of x on which the restriction $u|_V \in C^1_{loc}(V, \mathbb{R}^k)$ is developable. As a consequence, the assumptions of Theorems 1 and 2 can be weakened to local regularity while keeping the global conclusion intact.*

Remark 2.3. The continuity of f is necessary for the implications (a) \Rightarrow (b), (c) in Proposition 2.1 as the following example demonstrates. The function

$$f_1 : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f_1(s, t) := \begin{cases} 0 & st \neq 0 \\ 1 & st = 0, \end{cases}$$

satisfies condition (a) but not (b) or (c). Also, the following example shows the necessity of the distinction between the points in the local constancy regions from other points in condition (a):

$$f_2 : B^1 \rightarrow \mathbb{R}, \quad f_2(s, t) := \begin{cases} 0 & \max\{|s|, |t|\} \leq \frac{\sqrt{2}}{2} \\ |t| - \frac{\sqrt{2}}{2} & |t| > \frac{\sqrt{2}}{2}, \\ |s| - \frac{\sqrt{2}}{2} & |s| > \frac{\sqrt{2}}{2}. \end{cases}$$

In other words, a global formulation of condition (b) is not a statement equivalent to (b): The continuous function f_2 satisfies conditions (a) and (b), but there are $x \in B^1$ for which f cannot be constant on the intersection $L \cap B^1$ for any line L .

Remark 2.4. In the recent literature on similar problems, global (and hence inequivalent) versions of condition (b) are formulated as “developability of f ” in [36, Definition 2.29] and “ Ω -developability of f on Ω ” in [28, Section 3.1]. On the other hand condition (a) appears in the statement of [50, Theorem II], and then in [46] as the “condition (L)”. It is also equivalent to f being “countably developable” according to [28, Definition 1] and [29]. (Note that the assumption (24) in [28] is redundant for continuous f ; see Lemma 2.8). Finally, it is condition (c) that is stated as the property of the gradients (with empty-interior images) of C^1 mappings in the main result

of [37, 38]. (This local result was an ingredient of the proof of Theorem 2 in [43]). Generalizations of (a) to weaker regularity or higher dimensional settings can be found in [55, 34, 35].

Remark 2.5. The developability of a C^1 mapping on Ω , as formulated in Theorems 1 and 2, translates therefore to its Jacobian derivative satisfying condition (a). To clarify some discrepancy in the literature, we emphasize that we prefer to reserve the term *developable* for those mappings whose Jacobian derivatives are constant along lower dimensional flat strata in one form or another [44, 35], rather than directly using it for their derivative mappings as done in [36, Definition 2.29] or [28, 29]. This is for historical reasons, since this term is conventionally used to refer to (ruled) smooth surfaces of zero Gaussian curvature³[8].

Proof.

(a) \Rightarrow (b):

This is implicitly proved in [44, Lemma 3.7]. We pursue a slightly different strategy. For any $f : \Omega \rightarrow \mathbb{R}$, let C_f be the set on which f is locally constant:

$$C_f := \{x \in \Omega; f \text{ is constant in a neighborhood of } x\}.$$

Note that f is constant on the connected components of the open set C_f . If f satisfies the condition (a) of the proposition, for each $x \in \Omega \setminus C_f$ we choose a line L_x so that f is constant on the connected component l in $L_x \cap \Omega$, denoted by l_x .

We begin by the following simple but useful lemma.

Lemma 2.6. *Let $f \in C_{loc}^0(\Omega, \mathbb{R}^m)$ satisfy condition (a) in Proposition 2.1. Let $\Delta \subset \Omega$ be a closed triangular domain. If f is constant on two edges of Δ , then f is constant on Δ .*

Proof. Let x, y, z be the vertices of Δ and assume that f is constant on $[x, y]$ and $[x, z]$. For any point $w \in \Delta \setminus C_f$, l_w must cross one of the two segments $[x, y]$ or $[x, z]$. This implies $f(w) = f(x)$. If $w \in \Delta \cap C_f$, and if the segment $[w, x] \subset C_f$, then once again $f(w) = f(x)$. Finally, if $[w, x] \not\subset C_f$, let \tilde{w} be the closest element to w on the segment $[w, x]$ not in C_f . We already proved that $f(\tilde{w}) = f(x)$. But since \tilde{w} is in the closure of the connected component of w in C_f , and f is continuous, we have also $f(w) = f(\tilde{w}) = f(x)$. We conclude that $f|_{\Delta} \equiv f(x)$. ■

We now make a crucial observation on the constancy directions L_x .

Corollary 2.7. *Let $f \in C^0$ be continuous and satisfy condition (a) in Proposition 2.1. If $x \in \Omega \setminus C_f$, there exists only one line L for which f is constant on the connected component l of x in $L \cap \Omega$.*

Proof. Let $x \in \Omega \setminus C_f$ and assume that for two lines $L \neq \Lambda$, f is constant on the connected components l and λ of x in respectively $L \cap \Omega$ and $\Lambda \cap \Omega$. Choose a disk $B(x, \rho) \subset B(x, 2\rho) \subset \Omega$ and note that $\partial B(x, \rho)$ intersects l and λ on 4 points $x_1, x_2 \in l$ and $\chi_1, \chi_2 \in \lambda$. Now the assumptions of Lemma 2.6 is satisfied for the four closed triangular domains with vertices $x, x_i, \chi_j, i, j \in \{1, 2\}$. This implies that $x \in C_f$, which contradicts the first assumption on x . ■

As a consequence, when f is continuous, for each $x \in \Omega \setminus C_f$, there is only one choice for L_x and there is no ambiguity in the notation. We further observe:

Lemma 2.8. *Let the continuous mapping $f : \Omega \rightarrow \mathbb{R}^m$ satisfy condition (a) in Proposition 2.1. Then for all $x \in \Omega \setminus C_f$, $l_x \subset \Omega \setminus C_f$. Moreover, for all $y, z \in \Omega \setminus C_f$, $l_y \cap l_z = \emptyset$, or $l_y = l_z$.*

³Such surfaces were historically called developable since they can be “developed” (flattened out) on the plane. Perhaps it is the same terminology that was used in pre-digital age photography: A “developed” photo was one that was transferred from the cylindrical film onto the flat printing material without distorting the relative distances.

Proof. For showing the first conclusion, we argue by contradiction: Let $y \in C_f \cap l_x$ and let V be an open neighborhood of y in Ω on which f is constant. We observe that for $\delta > 0$ small enough $B(x, \delta) \setminus l_x$ cannot entirely lie in C_f , since otherwise, $B(x, \delta) \setminus l_x$ having only two connected components on both sides of l_x , and f being continuous, the value of f would be constant on $B(x, \delta)$, contradicting $x \notin C_f$. Therefore, there is a sequence $x_k \in \Omega \setminus C_f$ converging to x such that $f(x_k) \neq f(x)$. Since the value of f on x_k and x differ, $l_{x_k} \cap l_x = \emptyset$ for all k . As $x_k \rightarrow x$, we deduce that l_{x_k} must locally uniformly converge to l_x . This implies that $l_{x_k} \cap V$ is non-empty for k large enough and hence $f(x_k) = f(y) = f(x)$, which is a contradiction.

To prove the second statement, also assume by contradiction that $l_y \neq l_z$ and that $x \in l_y \cap l_z \neq \emptyset$. Choose a disk $B(x, \rho) \subset B(x, 2\rho) \subset \Omega$ and note that $\partial B(x, \rho)$ intersects l_y and l_z on 4 points $y_1, y_2 \in l_y$ and $z_1, z_2 \in l_z$. Now the assumptions of Lemma 2.6 is satisfied for the four closed triangular domains with vertices $x, y_i, z_j, i, j \in \{1, 2\}$. This implies that $x \in C_f$, which contradicts the first statement, as $y \notin C_f$ but $x \in l_y \cap C_f$. ■

We are now ready to prove (b) assuming (a). We first note that the condition (b) is obvious if $x \in C_f$. If $x \notin C_f$, without loss of generality, and through rotations, dilations and translations, we can assume that $x = (0, 0)$ and that l_x is parallel to the horizontal axis. We show that we can find $\delta^-, \delta^+ > 0$ such that the condition (b) holds true for $B^+(x, \delta^+)$ and $B^-(x, \delta^-)$ in the upper and lower planes, where the open half disks are defined by

$$B^\pm(x, \delta) := B(x, \delta) \cap \{(s, t) \in \mathbb{R}^2; \pm t > 0\}.$$

Then we can choose $B = B(x, \min\{\delta^+, \delta^-\})$ and the proof is complete.

Without loss of generality we concentrate on the upper half plane. If there exists $\delta^+ > 0$ such that $B(x, \delta^+) \subset C_f$, the conclusion is obvious: We choose l_y always parallel to the horizontal axis. Otherwise there exists a sequence $x_k \notin C_f$ in the upper open half plane converging to x . Note that by Lemma 2.8, any two (possibly unbounded) segments l_y, l_z do not intersect within Ω unless they are the same. This implies that $l_{x_k} \cap l_x = \emptyset$ and that the l_{x_k} must converge locally uniformly to l_x . We choose $\rho > 0$ such that the closed rectangular box

$$S_\rho^+ := \{(s, t) \in \mathbb{R}^2; |s| \leq \rho, 0 \leq t \leq \rho\}$$

is a subset of Ω . We let $x_0^\pm := (\pm\rho, 0)$, $x^\pm = (\pm\rho, \rho)$ and choose k large enough such that for $y = x_k$, $l_y = l_{x_k}$ intersects the two segments $[x_0^\pm, x^\pm]$ in their respective interior points y^- and y^+ . The closure of the convex open quadrilateral P created by the four vertices x_0^-, x_0^+, y^+, y^- lies in S_ρ^+ and we have that $[y^-, y^+] \subset l_y$ and $[x_0^-, x_0^+] \subset l_x$.

We claim that the condition (b) is valid for P (standing for B_x), which completes the proof since there exists some $\delta^+ > 0$, such that $B^+(x, \delta^+) \subset P$. For any $z \in P$, if $z \notin C_f$, then we can choose $\tilde{l}_z = l_z \cap P$, which lies between $\tilde{l}_{x_0^-}$ and \tilde{l}_{y^+} . If, on the other hand, $z \in C_f$, let the points $z_1 \in [y^-, y^+]$ and $z_0 \in [x_0^-, x_0^+]$ be those points on ∂P which lie vertically above and below z . Since $z_0, z_1 \notin C_f$, we can choose the two closest elements of $P \setminus C_f$ to z on the segment $[z_0, z_1]$, which we name respectively by \tilde{z}_0, \tilde{z}_1 . We observe that $l_{\tilde{z}_0}$ and $l_{\tilde{z}_1}$ cannot intersect l_x and l_y , which contain the upper and lower boundaries of P , and hence will intersect the two vertical boundaries of P between x_0^\pm and y^\pm in \tilde{z}_i^\pm , $i = 0, 1$. We claim that f is constant in the region X enclosed between $l_{\tilde{z}_0}$ (which is above \tilde{l}_x and below $l_{\tilde{z}_1}$) and $l_{\tilde{z}_1}$ (which is above \tilde{l}_{z_0} and below \tilde{l}_{y^+}) in P . Note that X is the convex quadrilateral created by $\tilde{z}_0^-, \tilde{z}_0^+, \tilde{z}_1^+, \tilde{z}_1^-$ and contains z in its interior.

The latter claim about the constancy of f on X , if proven, completes the proof of the former claim regarding P , since we are free to choose our non-intersecting constancy segments in this region X (in particular for z) in a manner that no such two segments intersect within P : If the directions $l_{\tilde{z}_0}$ and $l_{\tilde{z}_1}$ are parallel, choose the line L_p for all points $p \in X$ to be the line parallel to

them passing from p . If, on the other hand, the extensions of $l_{\tilde{z}_0}$ and $l_{\tilde{z}_1}$ meet at a point q (outside of Ω), for all $p \in X$ we choose the line L_p to be the one passing through p and q . If \tilde{l}_p , for any $p \in X$, were to intersect any other $\tilde{l}_{\tilde{p}}$ for $\tilde{p} \in P \setminus X$, it would have to first intersect one of the two $\tilde{l}_{\tilde{z}_i}$, $i = 0, 1$, which does not happen by construction. Note that X is the connected component of $C_f \cap P$ containing z . We can therefore foliate P in constancy segments by exhausting all the connected components of $C_f \cap P$.

To prove our final claim, that is the constancy of f on the region X , we note that f is constant on the segment $\tilde{l}_{\tilde{z}_1} = [\tilde{z}_1^-, \tilde{z}_1^+]$, and on the segment $[\tilde{z}_0, \tilde{z}_1]$, whose interior lies in C_f . Since $\tilde{z}_1 \in [\tilde{z}_1^-, \tilde{z}_1^+]$, by Lemma 2.6, f must be constant in the triangular domain created by $\tilde{z}_0, \tilde{z}_1^-, \tilde{z}_1^+$. But then this last assertion implies that f is constant on the segments $[\tilde{z}_0, \tilde{z}_1^-]$ and $[\tilde{z}_0, \tilde{z}_1^+]$, which combined with Lemma 2.6 again, using the constancy of f on segments $[\tilde{z}_0, \tilde{z}_0^-]$ and $[\tilde{z}_0, \tilde{z}_0^+] \subset l_{\tilde{z}_0}$, proves the claim that f is constant on X . The proof of (a) \Rightarrow (b) is complete.

(b) \Rightarrow (c):

It is logically obvious.

(c) \Rightarrow (a):

The proof can be found in the last page of [43]. We reproduce it here for completeness. Let $x \in \Omega$ and consider a Ball B_x over which the condition (c) is satisfied. In particular, f is constant on $L_x \cap B_x$ for some line L_x passing through x . Now, consider the (possibly infinite) segment

$$c_x := \text{The connected component of } x \text{ in } \{y \in L_x \cap \Omega; f(y) = f(x)\}.$$

We claim that c_x , which contains the segment $L_x \cap B_x$, is either equal to l_x , i.e. the connected component of x in $L_x \cap \Omega$, or $x \in C_f$. This proves that f is constant either on l_x , or in a neighborhood of x , and so (a) holds true.

We assume that $c_x \neq l_x$, and prove $x \in C_f$. In this case c_x must admit at least one endpoint within Ω . Let $y \in \Omega$ be that endpoint. Consider the open disk B_y centered at y in which all points admit constancy segments within B_y . Let the line L_y passing through y be such that f is constant on the segment $L_y \cap B_y$ and let z, w be the endpoints of this segment. L_y and L_x cannot be parallel, since then c_x can be extended along $L_x = L_y$ to include either z or w , which contradicts the fact that it must be a maximal connected constancy subset of l_x . Therefore, we can choose an element $\tilde{y} \in B_y \cap c_x$ to form an open triangle Δ with vertices \tilde{y}, z, w . f is constant on $[y, z]$ and on $[\tilde{y}, y]$, where $y \in (z, w)$. Since no segment $L_p \cap B_y$ departing from a point $p \in \Delta$ can reach ∂B_y on its both ends without crossing $[w, z]$ or $[\tilde{y}, y]$, we deduce that $f|_{\Delta} \equiv f(y) = f(x)$.

We observe that C_f contains Δ , and hence also the open segment (\tilde{y}, y) . We have thus found a non-empty interval in c_x from which we can propagate our local constancy property and so reach to the desired conclusion of $x \in C_f$. We argue by contradiction: If $x \notin C_f$, let \tilde{x} be the closest point to y on $[x, y]$ which is not in C_f . Certainly $C_f \supset (\tilde{x}, y) \supset (\tilde{y}, y)$ and $(x, y) \supset (\tilde{x}, y)$. Consider once more the open disk $B_{\tilde{x}}$ according to the condition (c). Since C_f is open, for some $\hat{y} \in (\tilde{x}, y)$ close enough to \tilde{x} , f is constant on a segment $[\hat{z}, \hat{w}] \subset B_{\tilde{x}} \cap C_f$ orthogonal to c_x at \hat{y} , with $\hat{y} \in (\hat{z}, \hat{w})$. Also note that for $\hat{x} \in [x, \hat{y}]$, f is constant on the portion $[\hat{x}, \hat{y}] = [x, \hat{y}] \cap B_{\tilde{x}} \supset [\tilde{x}, \hat{y}]$ of the segment $[x, \hat{y}] \subset c_x$. Since f takes the value $f(x)$ on both of $[\hat{x}, \hat{y}]$ and $[\hat{z}, \hat{w}]$, both within the set $B_{\tilde{x}}$, a similar argument as above shows that f is constant on the open triangle $\hat{\Delta}$ with vertices $\hat{x}, \hat{z}, \hat{w}$, and so $\hat{\Delta} \subset C_f$. But then $\tilde{x} \in (\hat{x}, \hat{y}) \subset \hat{\Delta} \subset C_f$, which contradicts its choice. \blacksquare

In view of Proposition 2.1, and as already observed in [36, Proposition 2.30], we have:

Corollary 2.9. *Let $\Omega \subset \mathbb{R}^2$ be a domain and let $f \in C_{loc}^0(\Omega, \mathbb{R}^m)$ satisfy any of the equivalent conditions in Proposition 2.1. Then for all $x \in \Omega$, there exists $L > 0$, an open disk B centered at*

x , and a unit Lipschitz vector field $\vec{\eta} : B \rightarrow \mathbb{R}^2$ with $\text{Lip } \vec{\eta} \leq L$, for which

$$\forall y \in B \quad \vec{\eta}(y) = \vec{\eta}(y + s\vec{\eta}(y)) \quad \text{and} \quad f(y) = f(y + s\vec{\eta}(y)) \quad \text{for all } s \text{ for which } y + s\vec{\eta}(y) \in B.$$

Proof. We choose a disk $B_x = B(x, 2\delta) \subset \Omega$ according to the condition (b) and we let $B = B(x, \delta)$. For any $p \in B$, there exists a line L_p such that f is constant on $\tilde{l}_p = L_p \cap B_x$, and no two such lines meet within B_x . We define a the mapping $\mathbf{\Lambda} : B \rightarrow \mathbb{RP}^1$ which associates to each point $p \in B$ the element of the real projective line \mathbb{RP}^1 determined by the direction of L_p , and we note that it is constant along the segments $L_p \cap B$ and continuous. Since B is simply connected, $\mathbf{\Lambda}$ can be lifted to a continuous mapping $\vec{\eta} : B \rightarrow \mathbb{S}^1$. By constancy of $\mathbf{\Lambda}$ along the segments $L_p \cap B$, $\vec{\eta}$ can only take two distinct values along them, and so its continuity implies that $\vec{\eta}$ is constant along these directions, which are now determined by $\vec{\eta}$ itself. Since two distinct lines L_p and L_q , for $p, q \in B$, do not meet except possibly at an at least a δ -distance from ∂B , we conclude that $\text{Lip } \vec{\eta} \leq 1/\delta$. \blacksquare

2.2. Developability through test functions. The following lemma will allow us to translate the developability properties of $v \in C^1$ into a property for the distribution $\nabla^2 v$:

Lemma 2.10. *Let $B \subset \mathbb{R}^2$ be an open disk and let $f \in C_{loc}^0(B, \mathbb{R}^m)$. Also, let $\vec{\eta} : B \rightarrow \mathbb{R}^2$ be a unit Lipschitz vector field with $\text{Lip } \vec{\eta} \leq L$, such that the following property holds true:*

$$\forall x \in B \quad \vec{\eta}(x) = \vec{\eta}(x + s\vec{\eta}(x)) \quad \text{for all } s \text{ for which } x + s\vec{\eta}(x) \in B.$$

Then the following two properties are equivalent:

- (a) $\forall x \in B \quad f(x) = f(x + s\vec{\eta}(x))$ for all s for which $x + s\vec{\eta}(x) \in B$,
- (b) $\forall \psi \in C_c^\infty(B) \quad \int_B f \text{div}(\psi \vec{\eta}) = 0$.

Proof. Let $\vec{\xi} = \vec{\eta}^\perp$ and $x \in B$. Choose $T_0 > 0$ such that $\gamma : (-T_0, T_0) \rightarrow B$ is a solution to the ODE $\gamma'(t) = \vec{\xi}(\gamma(t))$ in B with initial value $\gamma(0) = x$. Let $k(t) = -\gamma''(t) \cdot \vec{\eta}(\gamma(t))$ and note that $\|k\|_{L^\infty(-T_0, T_0)} \leq L$. It is straightforward to see by contradiction that there exists $0 < t_0 \leq T_0/2$ such that for all $t, t' \in [-t_0, t_0]$, and all $s, s' \in \mathbb{R}$,

$$\gamma(t) + s\vec{\eta}(\gamma(t)), \gamma(t') + s'\vec{\eta}(\gamma(t')) \in B \implies \gamma(t) + s\vec{\eta}(\gamma(t)) \neq \gamma(t') + s'\vec{\eta}(\gamma(t')),$$

unless $t = t'$ and $s = s'$. Since $\gamma([-t_0, t_0])$ is compact in B , we choose $0 < s_0 < \frac{1}{2L}$ such that the image of $[-t_0, t_0] \times [-s_0, s_0]$ under

$$\Phi(t, s) := \gamma(t) + s\vec{\eta}(\gamma(t)), \quad t \in [-t_0, t_0], s \in [-s_0, s_0],$$

lies compactly in B . We note that Φ is one-to-one and Lipschitz on $U := (-t_0, t_0) \times (-s_0, s_0)$ and that we have

$$\det \nabla \Phi(t, s) = 1 + sk(t) \geq \frac{1}{2} \quad \text{a.e. in } U.$$

We conclude that $\Phi : U \rightarrow \Phi(U) =: V_x \ni x$ is a bilipschitz change of variable.

We now calculate for any $\psi \in C_c^\infty(V_x)$:

$$\text{div}(\psi \vec{\eta}) = \partial_{\vec{\eta}}(\psi \vec{\eta}) \cdot \vec{\eta} + \partial_{\vec{\xi}}(\psi \vec{\eta}) \cdot \vec{\xi} = \nabla \psi \cdot \vec{\eta} + \psi \partial_{\vec{\xi}} \vec{\eta} \cdot \vec{\xi},$$

which gives for $\tilde{\psi} = \psi \circ \Phi$:

$$\text{div}(\psi \vec{\eta})(\Phi(t, s)) = \partial_s \tilde{\psi} + \tilde{\psi} \frac{k(t)}{1 + sk(t)}.$$

Therefore, letting $\tilde{f}(t, s) = f \circ \Phi(s, t) = f(\gamma(t) + s\vec{\eta}(\gamma(t)))$ we obtain

$$\begin{aligned}
(2.1) \quad \int_{V_x} f \operatorname{div}(\psi \vec{\eta}) &= \int_U f \operatorname{div}(\psi \vec{\eta}) \circ \Phi(s, t) (1 + sk(t)) \, ds dt \\
&= \int_U \tilde{f}(s, t) \left(\partial_s \tilde{\psi} + \tilde{\psi} \frac{k(t)}{1 + sk(t)} \right) (1 + sk(t)) \, ds dt \\
&= \int_U \tilde{f}(t, s) \left(k(t) \tilde{\psi} + \partial_s \tilde{\psi} + sk(t) \partial_s \tilde{\psi} \right) \, ds dt \\
&= \int_U \tilde{f}(t, s) \partial_s ((1 + sk(t)) \tilde{\psi}) \, ds dt.
\end{aligned}$$

If (a) is satisfied, then $\tilde{f}(t, s) = \tilde{f}(t)$ for all values of s for which the function is defined, and hence, since $\tilde{\psi}$ is compactly supported in U , we obtain:

$$(2.2) \quad \forall \psi \in C_c^\infty(V_x) \quad \int_{V_x} f \operatorname{div}(\psi \vec{\eta}) = 0.$$

Now for $\psi \in C_c^\infty(B)$, we let $K := \operatorname{supp}(\psi) \subset B$. K is compact and so it admits a finite covering of open sets of the form V_{x_i} , for $i = 1, \dots, n$. We consider a partition of unity associated with this covering

$$\theta_i \in C_c^\infty(V_{x_i}), \quad \sum_{i=1}^n \theta_i = 1 \text{ on } K,$$

and we conclude with (b) by (2.2):

$$\int_B f \operatorname{div}(\psi \vec{\eta}) = \sum_{i=1}^n \int_B f \operatorname{div}(\theta_i \psi \vec{\eta}) = \sum_{i=1}^n \int_{V_{x_i}} f \operatorname{div}(\theta_i \psi \vec{\eta}) = 0.$$

To prove the converse, assume (b) and let $\varphi \in L^1(U)$, be such that

$$(2.3) \quad \int_{-s_0}^{s_0} \varphi(t, s) \, ds = 0,$$

and define

$$\phi(t, s') := \frac{1}{1 + s'k(t)} \int_{-s_0}^{s'} \varphi(t, s) \, ds.$$

We can construct a sequence $\psi_k \in W_c^{1,\infty}(V_x)$ such that

$$\psi_k \circ \Phi \rightarrow \phi \quad \text{in } L^1(U) \quad \text{and} \quad \partial_s(\psi_k \circ \Phi) \rightarrow \partial_s \phi \quad \text{in } L^1(U).$$

Therefore by (b)

$$\int_{V_x} f \operatorname{div}(\psi \vec{\eta}) = 0 \quad \forall \psi \in C_c^\infty(V_x) \implies \int_{V_x} f \operatorname{div}(\psi_k \vec{\eta}) = 0,$$

which in view of (2.1) implies

$$\int_U \tilde{f} \varphi \, ds dt = \int_U \tilde{f}(t, s) \partial_s ((1 + sk(t)) \phi) \, ds dt = \lim_{k \rightarrow \infty} \int_U \tilde{f}(t, s) \partial_s ((1 + sk(t)) \psi_k \circ \Phi) \, ds dt = 0.$$

This fact being true for all $\varphi \in L^1(U)$ satisfying (2.3) implies that

$$\forall t \in (-t_0, t_0), \forall s \in (-s_0, s_0) \quad \tilde{f}(t, s) = \int_{-s_0}^{s_0} \tilde{f}(t, s) \, ds.$$

This means that for $|s| < s_0$:

$$f(x + s\vec{\eta}(x)) = \tilde{f}(0, s) = \tilde{f}(0, 0) = f(x).$$

The global property (a) is a direct consequence of this local property around each $x \in B$, and the fact that $\vec{\eta}$ itself is constant on $x + s\vec{\eta}(x) \in B$ for all s . \blacksquare

3. A HÖLDER CONTINUITY TOOLBOX

3.1. Notations and estimates on convolutions. For any open set $\omega \subset \mathbb{R}^n$, we will denote by $\|\cdot\|_{k;\omega}$ the norm of a function in the Banach space $C^k(\omega)$ and by $[\cdot]_k$ the $C^0(\Omega)$ norm of its k th (multi-index) derivatives. For $0 < \alpha < 1$, the corresponding Hölder norm and seminorms, and Banach spaces are shown by the standard conventions $\|\cdot\|_{k,\alpha;\omega}$, $[\cdot]_{k,\alpha;\omega}$ and

$$C^{k,\alpha}(\omega) := \{f : \omega \rightarrow \mathbb{R}; \|f\|_{k,\alpha} < \infty\}.$$

(This Banach space is sometimes also denoted by $C^{1,\alpha}(\bar{\omega})$ in the literature). Throughout this article, the constant $c = O(1)$ might change but is independent of all the data, unless specified by an argument. We will drop the subscript ω when denoting these norms for the specific case of an open set named ω , hence $\|\cdot\| = \|\cdot\|_{\cdot;\omega}$, etc.

For a mollifying kernel $\phi \in C_c^\infty(B^1)$ with $\int \phi = 1$, and $x \in \mathbb{R}^n$, we denote the standard convolution of a given mapping f with ϕ over the length scale ε by

$$f_\varepsilon(x) := f * \phi_\varepsilon(x) = \int_{\mathbb{R}^n} f(x-y)\phi_\varepsilon(y)dy,$$

where

$$\forall x \phi_\varepsilon(x) := \varepsilon^{-n}\phi\left(\frac{x}{\varepsilon}\right).$$

Naturally $f_\varepsilon(x)$ is defined for ε small enough provided f is integrable in some neighborhood of x .

We will extensively use the following well-known convolution estimates:

Lemma 3.1. *Let $V \Subset \omega$ be an open set compactly contained in ω . Assume that $0 < \alpha < 1$ and $f, h \in C^{0,\alpha}(\omega)$. Then for $\varepsilon < \text{dist}(\bar{V}, \partial\omega)$:*

- (i) $\|f_\varepsilon - f\|_{0;V} \leq [f]_{0,\alpha}\varepsilon^\alpha$,
- (ii) $\|\nabla f_\varepsilon\|_{0;V} \leq [f]_{0,\alpha}\varepsilon^{\alpha-1}$,
- (iii) $\|f_\varepsilon h_\varepsilon - (fh)_\varepsilon\|_{1;V} \leq c[f]_{0,\alpha}[h]_{0,\alpha}\varepsilon^{2\alpha-1}$.

The proof follows the same lines as in [11, Lemma 1] and is left to the reader. The commutator estimate Lemma 3.1-(iii) will be crucial for our analysis. It will be used to establish the quadratic estimate (4.3) ([11, Proposition 1]) for the pull-back metric of mollified isometric immersions.

3.2. A distributional product. The following statement is more or less known by the community at large and similar statements have appeared in various contexts, e.g. in the discussion of the Young integral, or implicitly when the weak Jacobian determinant is studied. For examples, see [2, Theorem 2.52] and [22, Theorem 13.16], or compare with [3, Corollary 1]. We have formulated a rather straightforward version which is more fitting to our needs.

Proposition 3.2. *Let $V \Subset \omega$ be an open set compactly contained in ω . Let $0 < \alpha < 1$ and $f, h \in C^{0,\alpha}(\omega)$. If $\alpha > 1/2$, then the sequence $f_\varepsilon \partial_j h_\varepsilon$ converges in the sense of distributions to a distribution on V which we denote by $f \partial_j h$. In particular, if $\psi \in W_0^{1,1}(V)$*

$$f \partial_j h[\psi] := \lim_{\varepsilon \rightarrow 0} \int_V f_\varepsilon(x) \partial_j h_\varepsilon(x) \psi(x) dx$$

exists. Moreover

$$\left| \int_V f_\varepsilon(x) \partial_j h_\varepsilon(x) \psi(x) dx - f \partial_j h[\psi] \right| \leq 2 \|f\|_{0,\alpha} [h]_{0,\alpha} \left(\varepsilon^{2\alpha-1} \|\psi\|_{L^1} + \varepsilon^\alpha \|\partial_j \psi\|_{L^1} \right)$$

and

$$|f \partial_j h[\psi]| \leq c(\omega, V) \|f\|_{0,\alpha} [h]_{0,\alpha} \|\psi\|_{L^1}^\alpha \|\partial_j \psi\|_{L^1}^{1-\alpha}.$$

Proof. Let $\psi \in W_0^{1,1}(V)$, and for $\varepsilon < \text{dist}(\bar{V}, \partial\omega)$ we define:

$$a^{(\varepsilon)} := \int_V f_\varepsilon(x) \partial_j h_\varepsilon(x) \psi(x) dx.$$

We first prove that $a^{(\varepsilon)}$ is a Cauchy sequence. We have for $\varepsilon, \varepsilon'$:

$$\begin{aligned} |a^{(\varepsilon)} - a^{(\varepsilon')}| &= \left| \int_V (f_\varepsilon \partial_j h_\varepsilon - f_{\varepsilon'} \partial_j h_{\varepsilon'}) \psi \right| \\ &\leq \left| \int_V f_\varepsilon (\partial_j h_\varepsilon - \partial_j h_{\varepsilon'}) \psi \right| + \left| \int_V (f_\varepsilon - f_{\varepsilon'}) \partial_j h_{\varepsilon'} \psi \right| \\ (3.1) \quad &= \left| - \int_V \partial_j f_\varepsilon (h_\varepsilon - h_{\varepsilon'}) \psi - \int_V f_\varepsilon (h_\varepsilon - h_{\varepsilon'}) \partial_j \psi \right| + \left| \int_V (f_\varepsilon - f_{\varepsilon'}) \partial_j h_{\varepsilon'} \psi \right| \\ &\leq \left| \int_V \partial_j f_\varepsilon (h_\varepsilon - h_{\varepsilon'}) \psi \right| + \left| \int_V f_\varepsilon (h_\varepsilon - h_{\varepsilon'}) \partial_j \psi \right| + \left| \int_V (f_\varepsilon - f_{\varepsilon'}) \partial_j h_{\varepsilon'} \psi \right| \\ &= I_1(\varepsilon, \varepsilon') + I_2(\varepsilon, \varepsilon') + I_3(\varepsilon, \varepsilon') \end{aligned}$$

We estimate for $\varepsilon, \varepsilon' \leq \delta$:

$$\begin{aligned} I_1(\varepsilon, \varepsilon') &= \left| \int_V \psi(x) \left(\int_{\mathbb{R}^n} (f(x-y) \partial_j \phi_\varepsilon(y)) dy \right) \left(\int_{\mathbb{R}^n} (h(x-z) (\phi_\varepsilon(z) - \phi_{\varepsilon'}(z))) dz \right) dx \right| \\ &= \left| \int_V \psi(x) \int_{\mathbb{R}^n} (f(x-y) - f(x)) \partial_j \phi_\varepsilon(y) dy \int_{\mathbb{R}^n} (h(x-z) - h(x)) (\phi_\varepsilon(z) - \phi_{\varepsilon'}(z)) dz dx \right| \\ &\leq \|\psi\|_{L^1} (\|f\|_{0,\alpha;\omega} \varepsilon^{\alpha-1}) ([h]_{0,\alpha} \delta^\alpha) \leq \|\psi\|_{L^1} [h]_{0,\alpha} [f]_{0,\alpha} \delta^{2\alpha-1}. \end{aligned}$$

By symmetry, a similar estimate is valid for $I_3(\varepsilon, \varepsilon')$:

$$I_3(\varepsilon, \varepsilon') \leq \|\psi\|_{L^1} [h]_{0,\alpha} [f]_{0,\alpha} \delta^{2\alpha-1}.$$

We also have:

$$\begin{aligned} I_2(\varepsilon, \varepsilon') &= \left| \int_V f_\varepsilon (h_\varepsilon - h_{\varepsilon'}) \partial_j \psi \right| \\ &\leq \|\partial_j \psi\|_{L^1} \|f\|_0 \sup_{x \in V} \left| \int_{\mathbb{R}^n} h(x-y) (\phi_\varepsilon(y) - \phi_{\varepsilon'}(y)) dy \right| \\ &= \|\partial_j \psi\|_{L^1} \|f\|_0 \sup_{x \in V} \left| \int_{\mathbb{R}^n} (h(x-y) - h(x)) (\phi_\varepsilon(y) - \phi_{\varepsilon'}(y)) dy \right| \\ &\leq \|\partial_j \psi\|_{L^1} \|f\|_0 [h]_{0,\alpha} \delta^\alpha. \end{aligned}$$

In view of (3.1), adding the three last estimates implies that $a^{(\varepsilon)}$ is Cauchy sequence. Hence the limit

$$f \partial_j h[\psi] := a = \lim_{\varepsilon \rightarrow 0} a^{(\varepsilon)},$$

exists. Now, we estimate for a fixed $0 < \varepsilon$, as $\varepsilon' \rightarrow 0$:

$$\left| \int_V f_\varepsilon(x) \partial_j h_\varepsilon(x) \psi(x) dx - f \partial_j h[\psi] \right| = |a^{(\varepsilon)} - a| \leq 2 \|\psi\|_{L^1} [f]_{0,\alpha} [h]_{0,\alpha} \varepsilon^{2\alpha-1} + \|\partial_j \psi\|_{L^1} \|f\|_0 [h]_{0,\alpha} \varepsilon^\alpha,$$

which yields:

$$\begin{aligned} |f\partial_j h[\psi]| &\leq |a^{(\varepsilon)}| + |a^{(\varepsilon)} - a| \leq |a^{(\varepsilon)}| + 2\varepsilon^{2\alpha-1}[f]_{0,\alpha}[h]_{0,\alpha}\|\psi\|_{L^1} + \varepsilon^\alpha\|f\|_0[h]_{0,\alpha}\|\partial_j\psi\|_{L^1} \\ &\leq \varepsilon^{\alpha-1}\|f\|_0[h]_{0,\alpha}\|\psi\|_{L^1} + 2\varepsilon^{2\alpha-1}[f]_{0,\alpha}[h]_{0,\alpha}\|\psi\|_{L^1} + \varepsilon^\alpha\|f\|_0[h]_{0,\alpha}\|\partial_j\psi\|_{L^1} \\ &\leq 3\|f\|_{0,\alpha}[h]_{0,\alpha}\left(\varepsilon^{\alpha-1}\|\psi\|_{L^1} + \varepsilon^\alpha\|\partial_j\psi\|_{L^1}\right). \end{aligned}$$

If the inequality were valid for all $\varepsilon > 0$, we could let $\varepsilon = \|\psi\|_{L^1}/\|\partial_j\psi\|_{L^1}$ to obtain the desired estimate. Here, we let

$$\varepsilon = \left(1 + \frac{\text{diam}(V)}{\text{dist}(\bar{V}, \partial\omega)}\right)^{-1} \frac{\|\psi\|_{L^1}}{\|\partial_j\psi\|_{L^1}} < \text{dist}(\bar{V}, \partial\omega).$$

by the Poincaré inequality

$$\|\psi\|_{L^1} \leq \text{diam}(V)\|\partial_j\psi\|_{L^1},$$

on V , to conclude the proof:

$$|f\partial_j h[\psi]| \leq 6\left(1 + \frac{\text{diam}(V)}{\text{dist}(\bar{V}, \partial\omega)}\right)^{1-\alpha} \|f\|_{0,\alpha}[h]_{0,\alpha}\|\psi\|_{L^1}^\alpha \|\partial_j\psi\|_{L^1}^{1-\alpha}.$$

■

3.3. A criteria for Hölder continuity.

Proposition 3.3. *Assume that $\omega \subset \mathbb{R}^n$ is an open coordinate rectangular box (i.e. a rectangular cuboid with edges parallel to the coordinate system) and that given $f \in L^1(\omega)$, for all $j = 1, \dots, n$, we have*

$$(3.2) \quad \text{for all } j \in \{1, \dots, n\}, \forall \psi \in C_c^\infty(\omega) \quad \left| \int_\omega f \partial_j \psi \right| \leq \|\psi\|_{L^1(\omega)}^\alpha \|\partial_j \psi\|_{L^1(\omega)}^{1-\alpha}.$$

Then $f \in C^{0,\alpha}(\omega)$ and $[f]_{0,\alpha} \leq 2^{n+1}$.

Proof. We prove the statement by induction over the dimension.

First let $n = 1$ and $\omega \subset \mathbb{R}$ and be an open interval and let $I \subset \omega$ be any open subinterval. Note that for any $\phi \in C_c^\infty(I)$, we can construct a sequence $\phi_k \in C_c^\infty(I)$ with $\int_I \phi_k = 0$, converging strongly in L^1 to $\phi - \int_I \phi$. Let $\psi_k(t) = \int_{-\infty}^t \phi_k(s) ds$ and note that $\psi_k \in W_0^{1,1}(I)$. We have therefore for all $\phi \in L^1(I)$:

$$\begin{aligned} \left| \int_I (f - \int_I f) \phi \right| &= \left| \int_I f (\phi - \int_I \phi) \right| = \lim_{k \rightarrow \infty} \left| \int_I f \phi_k \right| = \lim_{k \rightarrow \infty} \left| \int_I f \psi_k' \right| \leq \lim_{k \rightarrow \infty} \|\psi_k\|_{L^1}^\alpha \|\psi_k'\|_{L^1}^{1-\alpha} \\ &\leq \lim_{k \rightarrow \infty} |I|^\alpha \|\psi_k'\|_{L^1}^\alpha \|\psi_k'\|_{L^1}^{1-\alpha} = \lim_{k \rightarrow \infty} |I|^\alpha \|\phi_k - \int_I \phi_k\|_{L^1} = |I|^\alpha \|\phi - \int_I \phi\|_{L^1} \\ &\leq 2|I|^\alpha \|\phi\|_{L^1}, \end{aligned}$$

where we used the Poincaré inequality on I . Since $(L^1)' = L^\infty$, and $C_c^\infty(I)$ is dense in $L^1(I)$, we conclude that

$$(3.3) \quad \|f - \int_I f\|_{L^\infty(I)} \leq 2|I|^\alpha.$$

Now, let $y \in \omega$, and let $d = \text{dist}(y, \partial\omega)$, and for $z \in (y - d/2, y + d/2)$ and $r < d/2$ define

$$h_r(z) := \int_{z-r}^{z+r} f(x) dx.$$

h_r is continuous in z and converges a.e. to f on $(y - d/2, y + d/2)$ as $r \rightarrow 0$. On the other hand, for all $r' < r < d/2$:

$$\left| h_r(z) - h_{r'}(z) \right| \leq \int_{(z-r, z+r) \times (z-r', z+r')} |f(x) - f(x')| dx dx' \leq 4|(z-r, z+r)|^\alpha = 2^{\alpha+2} r^\alpha,$$

where we applied the bound (3.3) to $I = (z-r, z+r)$. As a consequence, h_r is a Cauchy sequence in the uniform norm and f is continuous as the uniform limit of the h_r for $r \rightarrow 0$. Now, for any $x, y \in \omega$, applying once again (3.3) to $I = (x, y) \subset \omega$ we obtain that

$$|f(x) - f(y)| \leq 4|x - y|^\alpha,$$

which implies that $f \in C^{0,\alpha}(\omega)$ and that $[f]_{0,\alpha} \leq 4$ as required.

Now, assume that $n > 1$ and that the statement is true for $n-1$. Let $Q \subset \omega$ be any coordinate rectangular box, i.e.

$$Q := \prod_{j=1}^n I_j \subset \omega,$$

with the open intervals $I_k \subset \mathbb{R}$. Let $\hat{x} := (x_1, \dots, x_{n-1})$ and note that for all $h \in L^1(Q)$,

$$\text{for a.e. } \hat{x} \in Q^{n-1} := \prod_{j=1}^{n-1} I_j, \quad \hat{h}(\hat{x}) := \int_{I_n} h(x) dx_n$$

is well defined and belongs to $L^1(Q)$. We claim that:

$$(3.4) \quad \text{ess sup}_{x \in Q} \left| f(x) - \hat{f}(\hat{x}) \right| \leq 2|I_n|^\alpha.$$

Let us first show that the conclusion holds assuming the claim (3.4) is true for all coordinate boxes $Q \subset \omega$. Let $\psi \in C_c^\infty(Q^{n-1})$ and for a, b the extremities of I_n , choose $0 \leq \theta_k \leq 1$ in $C_c^\infty(I_n)$ such that $\theta_k \equiv 1$ on $(a + 1/k, b - 1/k)$. For $1 \leq j \leq n-1$ we have by our main assumption on f :

$$\begin{aligned} \left| \int_{Q^{n-1}} \hat{f} \partial_j \psi \right| &= \frac{1}{|I_n|} \left| \int_Q f(x) \partial_j \psi(\hat{x}) dx \right| = \lim_{k \rightarrow \infty} \frac{1}{|I_n|} \left| \int_Q f(x) \theta_k(x_n) \partial_j \psi(\hat{x}) dx \right| \\ &= \lim_{k \rightarrow \infty} \frac{1}{|I_n|} \left| \int_Q f \partial_j(\theta_k \psi) \right| \leq \lim_{k \rightarrow \infty} \frac{1}{|I_n|} \|\theta_k \psi\|_{L^1(Q)}^\alpha \|\partial_j(\theta_k \psi)\|_{L^1(Q)}^{1-\alpha} \\ &= \lim_{k \rightarrow \infty} \frac{1}{|I_n|} \|\theta_k \psi\|_{L^1(Q)}^\alpha \|\theta_k \partial_j(\psi)\|_{L^1(Q)}^{1-\alpha} = \frac{1}{|I_n|} \|\psi\|_{L^1(Q)}^\alpha \|\partial_j(\psi)\|_{L^1(Q)}^{1-\alpha} \\ &= \|\psi\|_{L^1(Q^{n-1})}^\alpha \|\partial_j \psi\|_{L^1(Q^{n-1})}^{1-\alpha}. \end{aligned}$$

Applying the induction assumption, we deduce that

$$(3.5) \quad \hat{f} \in C^{0,\alpha}(Q^{n-1}), \quad [\hat{f}]_{0,\alpha} \leq 2^n.$$

To prove that f is continuous, fix $y \in \omega$ and consider a box $Q_d = Q^{n-1} \times (y_n - d, y_n + d) \subset \omega$ containing y . For all $z \in Q^{n-1} \times (y_n - d/2, y_n + d/2)$, and $r < d/2$ we define

$$h_r(z) := \int_{z_n-r}^{z_n+r} f(\hat{z}, s) ds$$

Note that, if r is fixed, the vertical averages of f are continuous in \hat{z} as established in (3.5), and so h_r is continuous in z . Applying (3.4) to $Q = Q^{n-1} \times (z_n - r, z_n + r)$ we have for $r' < r < d/2$:

$$\left| h_r(z) - h_{r'}(z) \right| \leq \int_{(z-r, z+r) \times (z-r', z+r')} |f(\hat{z}, s) - f(\hat{z}, s')| ds ds' \leq 4|(z-r, z+r)|^\alpha = 2^{\alpha+2} r^\alpha,$$

which implies that h_r locally uniformly converge to their limit, which happens to be f . Hence f is continuous in ω .

To obtain a Hölder estimate, let $x, y \in \omega$. First, assume $x_j = y_j$ for some $1 \leq j \leq n$. We re-arrange the coordinates so that $j = n$ and note that for any sequence of coordinate rectangular boxes Q_k of height $1/k$, containing x and y , such that $x_n = y_n$ is the midpoint of $I_{n,k}$, we have by (3.5)

$$|f(x) - f(y)| = \lim_{k \rightarrow \infty} |\hat{f}_{Q_k}(\hat{x}) - \hat{f}_{Q_k}(\hat{y})| \leq 2^n |\widehat{x^n} - \widehat{y^n}| \leq 2^n |x - y|.$$

If, on the other hand, $x_j \neq y_j$ for all j , we can choose a coordinate rectangular box $Q \subset \omega$ which has x and y as its opposite vertices on the largest diameter. We obtain by applying (3.5) and (3.4) to the now continuous f :

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - \hat{f}(\hat{x})| + |\hat{f}(\hat{x}) - \hat{f}(\hat{y})| + |f(y) - \hat{f}(\hat{y})| \\ &\leq 4|x_n - y_n|^\alpha + 2^n |\hat{x} - \hat{y}|^\alpha \leq (4 + 2^n) |x - y|^\alpha \\ &\leq 2^{n+1} |x - y|^\alpha, \end{aligned}$$

which is the desired bound. Note that the sequence Q_k in the first case, and the box Q in the second case, exist since ω is assumed to be a coordinate box itself.

It remains to prove the claim (3.4). For any $\phi \in C_c^\infty(Q)$, consider a sequence $\tilde{\phi}_k \in C_c^\infty(Q)$ converging strongly in L^1 to $\phi - \hat{\phi}(\hat{x})$. As a consequence,

$$\lim_{k \rightarrow \infty} \int_{I_n} \tilde{\phi}_k(x) dx_n = 0 \quad \text{in } L^1(Q^{n-1}).$$

Choose a $\varphi \in C_c^\infty(I_n)$ such that $\int_{I_n} \varphi = 1$ and define

$$\phi_k(x) = \tilde{\phi}_k(x) - \left(\int_{I_n} \tilde{\phi}_k(\hat{x}, s) ds \right) \varphi(x_n).$$

Then $\phi_k \in C_c^\infty(Q)$ converges in L^1 to $\phi - \hat{\phi}(\hat{x})$ and $\hat{\phi}_k(\hat{x}) = \int_{I_n} \phi_k(x) dx_n = 0$. Letting $\psi_k(\hat{x}_n, t) = \int_{-\infty}^t \phi_k(\hat{x}, s) ds$ and noting that $\psi_k \in W_0^{1,1}(Q)$, we obtain for all $\phi \in C_c^\infty(Q)$:

$$\begin{aligned} \left| \int_Q (f - \hat{f}(\hat{x})) \phi \right| &= \left| \int_Q f(\phi - \hat{\phi}(\hat{x})) \right| = \lim_{k \rightarrow \infty} \left| \int_Q f \phi_k \right| = \lim_{k \rightarrow \infty} \left| \int_Q f \partial_n \psi_k \right| \\ &\leq \lim_{k \rightarrow \infty} \|\psi_k\|_{L^1}^\alpha \|\partial_n \psi_k\|_{L^1}^{1-\alpha} \leq \lim_{k \rightarrow \infty} |I_n|^\alpha \|\partial_n \psi_k\|_{L^1}^\alpha \|\partial_n \psi_k\|_{L^1}^{1-\alpha} \\ &= \lim_{k \rightarrow \infty} |I_n|^\alpha \|\phi_k - \int_{I_n} \phi_k(x) dx_n\|_{L^1} = |I_n|^\alpha \|\phi - \int_{I_n} \phi(x) dx_n\|_{L^1(Q)} \\ &\leq 2|I_n|^\alpha \|\phi\|_{L^1(Q)}, \end{aligned}$$

where we used the Poincaré inequality on I_n . Once again, using the fact that L^∞ is the dual of L^1 , and the density of $C_c^\infty(Q)$ in $L^1(Q)$, we conclude with (3.4). ■

4. A SECOND FUNDAMENTAL FORM FOR $C^{1,\alpha}$ ISOMETRIES

Let $u \in C_{loc}^{1,\alpha}(\Omega, \mathbb{R}^3)$ be an isometric immersion with $\alpha > 2/3$ and let $V \Subset \Omega$ be any smooth simply-connected domain compactly contained in Ω , e.g. an open disk. We intend to show the following three statements, which we will formulate more precisely later on. We shall prove:

1. A weak notion of the second fundamental form $A_{ij} = -\partial_i u \cdot \partial_j \vec{n}$ makes sense for the immersion u on V . This part can be carried out for $\alpha > 1/2$.

2. $A = [A_{ij}]$ is symmetric and curl free, and is equal to $\nabla^2 v$ for a scalar function $v \in C_{loc}^{1,\alpha}(V)$.
3. $\mathcal{D}et(D^2 v) = -\frac{1}{2} \text{curl}^T \text{curl}(\nabla v \otimes \nabla v) = 0$ in $\mathcal{D}'(V)$.

We choose an open domain ω such that $V \Subset \omega \Subset \Omega$. Let $\phi \in C_c^\infty(B^1)$ be a nonnegative function with $\int \phi = 1$. Let $u_\varepsilon := u * \phi_\varepsilon : V \rightarrow \mathbb{R}^3$ be the standard mollification of u on V with $\varepsilon < \text{dist}(\bar{V}, \partial\omega)$. If ε is small enough, u_ε is an immersion since ∇u_ε converges uniformly to ∇u on V , and

$$|\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon| \geq 1/2.$$

We define

$$N^\varepsilon := \frac{\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon}{|\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon|}$$

to be the unit normal corresponding to u_ε , and we let

$$\vec{n}_\varepsilon := \vec{n} * \phi_\varepsilon$$

to be the mollification of the unit normal $\vec{n} = \partial_1 u \times \partial_2 u$ of immersion u . A direct application of Lemma 3.1 yields:

$$(4.1) \quad \|\partial_i u_\varepsilon - \partial_i u\|_{0;V} \leq \|u\|_{1,\alpha} \varepsilon^\alpha, \quad \|u_\varepsilon\|_{2;V} \leq \|u\|_{1,\alpha} \varepsilon^{\alpha-1}; \quad \|\vec{n}_\varepsilon - \vec{n}\|_{0;V} \leq \|u\|_{1,\alpha}^2 \varepsilon^\alpha.$$

We also note that

$$(4.2) \quad \|N^\varepsilon - \vec{n}\|_{0;V} \leq c \|u\|_{1,\alpha}^2 \varepsilon^\alpha, \quad \|\nabla N^\varepsilon\|_{0;V} \leq c \|u\|_{1,\alpha}^2 \varepsilon^{\alpha-1}.$$

To see the first estimate we observe that

$$\begin{aligned} \|N^\varepsilon - \vec{n}\|_{0;V} &\leq \left\| \frac{\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon}{|\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon|} - (\partial_1 u \times \partial_2 u) \right\|_{0;V} \\ &= \frac{1}{|\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon|} \left\| \partial_1 u_\varepsilon \times \partial_2 u_\varepsilon - |\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon| (\partial_1 u \times \partial_2 u) \right\|_{0;V} \\ &\leq 2 \left(\left\| \partial_1 u_\varepsilon \times \partial_2 u_\varepsilon - \partial_1 u \times \partial_2 u \right\|_{0;V} + \left\| |\partial_1 u_\varepsilon \times \partial_2 u_\varepsilon| - 1 \right\|_{0;V} \right) \\ &\leq 4 \left\| \partial_1 u_\varepsilon \times \partial_2 u_\varepsilon - \partial_1 u \times \partial_2 u \right\|_{0;V} \leq c \|u\|_{1,\alpha}^2 \varepsilon^\alpha, \end{aligned}$$

where we used (4.1). The second estimate also follows from (4.1) and a direct calculation.

We denote the pull-back metric induced by u_ε by $\mathcal{G}^\varepsilon := (\nabla u_\varepsilon)^T \nabla u_\varepsilon = u_\varepsilon^* \mathcal{E}_3$, where \mathcal{E}_3 is the Euclidean metric of \mathbb{R}^3 . Remember that $g := (\nabla u)^T (\nabla u)$ is the Euclidean metric \mathcal{E}_2 on Ω . By the commutator estimate Lemma 3.1-(iii) we have

$$\|\mathcal{G}^\varepsilon - (g * \phi_\varepsilon)\|_{1;V} = \|(\nabla u_\varepsilon)^T \nabla u_\varepsilon - ((\nabla u)^T \nabla u)_\varepsilon\|_{1;V} \leq c \|u\|_{1,\alpha}^2 \varepsilon^{2\alpha-1}.$$

So, in view of the fact that $\mathcal{E}_2 * \phi_\varepsilon = \mathcal{E}_2$ on V we obtain:

$$(4.3) \quad \|\mathcal{G}^\varepsilon - \mathcal{E}_2\|_{1;V} \leq c \|u\|_{1,\alpha}^2 \varepsilon^{2\alpha-1}.$$

Compare with [11, Proposition 1], which is stated under more general settings and where the mollifier needs to be symmetric.

We finally denote by the symmetric matrix field $A^\varepsilon := [A_{ij}^\varepsilon]$ the second fundamental form associated with the immersion u_ε in the local coordinates, i.e.

$$A_{ij}^\varepsilon := \partial_{ij} u_\varepsilon \cdot N^\varepsilon = -\partial_i u_\varepsilon \cdot \partial_j N^\varepsilon,$$

which by (4.1) satisfies the obvious bound

$$(4.4) \quad \|A^\varepsilon\|_{0;V} \leq \|u_\varepsilon\|_{2;V} \leq \|u\|_{1,\alpha} \varepsilon^{\alpha-1}.$$

4.1. Definition of the second fundamental form A . We are ready now to prove our main claims about the existence of a second fundamental form for the isometric immersion u and its properties. In what follows, the constant $C = c(\omega, V, \|u\|_{1,\alpha;\omega})$, according to our convention for c .

Proposition 4.1. *If $\alpha > 1/2$, then A_{ij}^ε converges in the sense of distributions to a distribution $A_{ij} \in \mathcal{D}'(V)$. Moreover, we have for all $\psi \in C_c^\infty(V)$:*

$$|(A_{ij}^\varepsilon - A_{ij})[\psi]| \leq C \left(\varepsilon^{2\alpha-1} \|\psi\|_{L^1} + \varepsilon^\alpha \|\partial_j \psi\|_{L^1} \right)$$

and

$$(4.5) \quad |A_{ij}[\psi]| \leq C \|\psi\|_{L^1}^\alpha \|\partial_j \psi\|_{L^1}^{1-\alpha}.$$

Proof. Applying Proposition 3.2 to the components $f := \partial_i u^m$, $h := \vec{n}_m$, for $m \in \{1, 2, 3\}$, we deduce that for a distribution $A_{ij} \in \mathcal{D}'(V)$,

$$B_{ij}^\varepsilon := -\partial_i u_\varepsilon \cdot \partial_j \vec{n}_\varepsilon \rightarrow A_{ij} := -\partial_i u \cdot \partial_j \vec{n} \quad \text{in } \mathcal{D}'(V),$$

and

$$|B_{ij}^\varepsilon[\psi] - A_{ij}[\psi]| \leq C \left(\varepsilon^{2\alpha-1} \|\psi\|_{L^1} + \varepsilon^\alpha \|\partial_j \psi\|_{L^1} \right).$$

On the other hand, we have for all $\psi \in W_0^{1,1}(V)$:

$$\begin{aligned} |A_{ij}^\varepsilon[\psi] - A_{ij}[\psi]| &\leq |B_{ij}^\varepsilon[\psi] - A_{ij}[\psi]| + \left| \int_B \partial_i u_\varepsilon(x) \cdot (\partial_j \vec{n}_\varepsilon(x) - \partial_j N^\varepsilon(x)) \psi(x) dx \right| \\ &\leq C \left(\varepsilon^{2\alpha-1} \|\psi\|_{L^1} + \varepsilon^\alpha \|\partial_j \psi\|_{L^1} \right) \\ &\quad + \left| \int_V \partial_{ij} u_\varepsilon \cdot (\vec{n}_\varepsilon - N^\varepsilon) \psi dx \right| + \left| \int_V \partial_i u_\varepsilon \cdot (\vec{n}_\varepsilon - N^\varepsilon) \partial_j \psi \right| \\ &\leq C \left(\varepsilon^{2\alpha-1} \|\psi\|_{L^1} + \varepsilon^\alpha \|\partial_j \psi\|_{L^1} \right), \end{aligned}$$

where we used Proposition 3.2, (4.1) and (4.2). The second estimate follows as in Proposition 3.2. \blacksquare

Definition 2. We define the weak second fundamental form of u as the distribution

$$A : C_c^\infty(V) \rightarrow \mathbb{R}_{sym}^{2 \times 2}, \quad A[\psi] := [A_{ij}[\psi]]_{2 \times 2}.$$

4.2. A satisfies Gauss-Codazzi-Mainardi equations (in a weak sense).

Proposition 4.2. *We have the uniform estimate*

$$(4.6) \quad \|\text{curl } A^\varepsilon\|_{0;V} \leq C \varepsilon^{3\alpha-2}.$$

In particular, if $\alpha > 2/3$, then $\text{curl } A = 0$ in $\mathcal{D}'(V)$.

Proof. We remind that the Christoffel symbols associated with a metric g are given by

$$\Gamma_{jk}^i(g) = \frac{1}{2} g^{im} (\partial_k g_{jm} + \partial_j g_{km} - \partial_m g_{jk}),$$

where the Einstein summation convention is used. Hence, in view of (4.3), we have

$$\Gamma_{kj}^{i,\varepsilon} := \Gamma_{jk}^i(\mathcal{G}^\varepsilon) \rightarrow 0$$

uniformly, if $\alpha > 1/2$, with the estimate

$$(4.7) \quad \|\Gamma_{kj}^{i,\varepsilon}\|_{0;V} \leq C \varepsilon^{2\alpha-1}.$$

Writing the Codazzi-Mainardi equations [26, Equation (2.1.6)] for the immersion u_ε we have

$$\begin{aligned}\partial_2 A_{11}^\varepsilon - \partial_1 A_{12}^\varepsilon &= A_{11}^\varepsilon \Gamma_{12}^{1,\varepsilon} + A_{12}^\varepsilon (\Gamma_{12}^{2,\varepsilon} - \Gamma_{11}^{1,\varepsilon}) - A_{22}^\varepsilon \Gamma_{11}^{2,\varepsilon}, \\ \partial_2 A_{12}^\varepsilon - \partial_1 A_{22}^\varepsilon &= A_{11}^\varepsilon \Gamma_{22}^{1,\varepsilon} + A_{12}^\varepsilon (\Gamma_{22}^{2,\varepsilon} - \Gamma_{12}^{1,\varepsilon}) - A_{22}^\varepsilon \Gamma_{12}^{2,\varepsilon}.\end{aligned}$$

Hence, by (4.4) and (4.7),

$$\|\partial_2 A_{i1}^\varepsilon - \partial_1 A_{i2}^\varepsilon\|_{0;V} \leq C\varepsilon^{\alpha-1}\varepsilon^{2\alpha-1},$$

which completes the proof. \blacksquare

Before proceeding to the next step, we need to establish a further property of the sequence A^ε . Observe that if $u \in C^2(\Omega, \mathbb{R}^3)$ were an isometric immersion, then the determinant of its second fundamental form A would vanish, since, in this particular case, it would be equal to the Gaussian curvature of the Euclidean metric. In our situation, A does not enjoy the sufficient regularity for that $\det A$ is defined directly. But once again it can be shown that for $\alpha > 1/2$, the $\det A^\varepsilon$ converge in the sense of distributions to 0, that with a rate which will be crucial in our analysis. The following statement is a slight variant of [11, Proposition 7] recast for our purposes:

Proposition 4.3. $\forall \psi \in C_c^\infty(V) \quad \left| \int_V (\det A^\varepsilon) \psi \right| \leq C\varepsilon^{2\alpha-1} \|\psi\|_{W^{1,1}}.$

Proof. In view of the formula for the (0,4)-Riemann curvature tensor [26, Equation (2.1.2)]

$$R_{iljk} = g_{lm}(\partial_k \Gamma_{ij}^m - \partial_j \Gamma_{ik}^m + \Gamma_{ks}^m \Gamma_{ij}^s - \Gamma_{js}^m \Gamma_{ik}^s),$$

and by the Gauss equation [26, Equation (2.1.7)], (4.3) and (4.7) we have on V :

$$\begin{aligned}\det A^\varepsilon &= R_{1212}(\mathcal{G}^\varepsilon) = \mathcal{G}_{1m}^\varepsilon (\partial_1 \Gamma_{22}^{m,\varepsilon} - \partial_2 \Gamma_{21}^{m,\varepsilon} + \Gamma_{1s}^{m,\varepsilon} \Gamma_{22}^{s,\varepsilon} - \Gamma_{2s}^{m,\varepsilon} \Gamma_{21}^{s,\varepsilon}) \\ &= \partial_1 (\mathcal{G}_{1m}^\varepsilon \Gamma_{22}^{m,\varepsilon}) - \partial_2 (\mathcal{G}_{1m}^\varepsilon \Gamma_{21}^{m,\varepsilon}) + O(\varepsilon^{2\alpha-1}) \\ &= 2\partial_{12} \mathcal{G}_{12}^\varepsilon - \partial_{11} \mathcal{G}_{22}^\varepsilon - \partial_{22} \mathcal{G}_{11}^\varepsilon + O(\varepsilon^{2\alpha-1}) \\ &= -\operatorname{curl}^T \operatorname{curl} \mathcal{G}^\varepsilon + O(\varepsilon^{2\alpha-1}).\end{aligned}$$

The conclusion follows by an integration by parts involving the first term and applying (4.3). \blacksquare

4.3. A as the Hessian of a scalar function.

Proposition 4.4. *There exists $v \in C_{loc}^{1,\alpha}(V) \cap W^{1,2}(V)$ such that $\nabla^2 v = A$ in the sense of distributions. Moreover*

$$\operatorname{Det} D^2 v = -\frac{1}{2} \operatorname{curl}^T \operatorname{curl}(\nabla v \otimes \nabla v) = 0 \quad \text{in } \mathcal{D}'(V).$$

Remark 4.5. In our setting, it can also be shown that $v \in C^{1,\alpha}(V)$ by following the analysis on a larger open set containing V . We will not need this fact.

Proof. Remember that V is assumed to be simply connected and smooth. Let A^ε be as above and let F^ε be the solution to the Neumann problem

$$\begin{cases} \Delta F^\varepsilon = \operatorname{div} A^\varepsilon & \text{in } V \\ \partial_\nu F^\varepsilon = A^\varepsilon \cdot \nu & \text{on } \partial V \end{cases}$$

Since $\operatorname{div}(\nabla F^\varepsilon - A^\varepsilon) = 0$, there exists a vector field E^ε such that $A^\varepsilon = \nabla F^\varepsilon + \nabla^\perp E^\varepsilon$. E^ε solves

$$\begin{cases} \Delta E^\varepsilon = \operatorname{curl}(A^\varepsilon - \nabla F^\varepsilon) = \operatorname{curl} A^\varepsilon & \text{in } V \\ E^\varepsilon = \operatorname{const.} & \text{on } \partial V \end{cases}$$

Since A^ε is symmetric, we have $\partial_2(F_1^\varepsilon + E_2^\varepsilon) = \partial_1(F_2^\varepsilon - E_1^\varepsilon)$, and thus we derive the existence of a scalar function $v^{(\varepsilon)} \in C^\infty(V)$ satisfying

$$\partial_1 v^{(\varepsilon)} = F_1^\varepsilon + E_2^\varepsilon, \quad \partial_2 v^{(\varepsilon)} = F_2^\varepsilon - E_1^\varepsilon,$$

i.e. $\nabla v^{(\varepsilon)} = F^\varepsilon + (E_1^\varepsilon, -E_2^\varepsilon)$. As a consequence

$$A^\varepsilon = \nabla^2 v^{(\varepsilon)} - \nabla(E^\varepsilon)^\perp + \nabla^\perp E^\varepsilon.$$

Standard elliptic estimates imply that for any $p < \infty$

$$(4.8) \quad \|\nabla E^\varepsilon\|_{W^{1,p}(V)} \leq c(p, V) \|\operatorname{curl} A^\varepsilon\|_{0;V}.$$

As a consequence, fixing $p > 2$, and in view of (4.6), $\|A^\varepsilon - \nabla^2 v^{(\varepsilon)}\|_{0;V} \leq C\varepsilon^{3\alpha-2} \rightarrow 0$ and therefore $\nabla^2 v^{(\varepsilon)}$ converges to A in the sense of distributions.

Now, for all $\varphi \in C_c^\infty(V)$, there exists a vector field Ψ in V , vanishing on ∂V , such that $\operatorname{div} \Psi = \varphi - \int_V \varphi$, for which $\|\Psi\|_{W^{1,2}(V)} \leq c(V) \|\varphi\|_{L^2(V)}$ (see e.g. [1]). By adjusting the $\nabla v^{(\varepsilon)}$ so that it is of average 0 over V , and in view of Proposition 4.1, we obtain that for all $\varepsilon, \varepsilon' \leq \delta$:

$$\begin{aligned} \left| \int_V (\nabla v^{(\varepsilon)} - \nabla v^{(\varepsilon')}) \varphi \right| &= \left| \int_V (\nabla v^{(\varepsilon)} - \nabla v^{(\varepsilon')}) (\varphi - \int_V \varphi) \right| = \left| \int_V (\nabla v^{(\varepsilon)} - \nabla v^{(\varepsilon')}) \operatorname{div} \Psi \right| \\ &= \left| \int_V (\nabla^2 v^{(\varepsilon)} - \nabla^2 v^{(\varepsilon')}) \Psi \right| \\ &\leq \left| \int_V (A^\varepsilon - A^{\varepsilon'}) \Psi \right| + \left| \int_V (\nabla^2 v^{(\varepsilon)} - A^\varepsilon) \Psi \right| + \left| \int_V (\nabla^2 v^{(\varepsilon')} - A^{\varepsilon'}) \Psi \right| \\ &\leq C\delta^{3\alpha-2} \|\Psi\|_{W^{1,1}} \leq C\delta^{3\alpha-2} \|\varphi\|_{L^2(V)}. \end{aligned}$$

By duality, we conclude that $\nabla v^{(\varepsilon)}$ is converging in $L^2(V)$ to a vector field F and that $\nabla F = A$. It is now straightforward to see that, if necessary by adjusting the $v^{(\varepsilon)}$ by constants, $v^{(\varepsilon)}$ converges strongly in $W^{1,2}$ to some $v \in W^{1,2}(V)$ and that $A = \nabla^2 v$.

To complete the proof, it is only necessary to show that a) $\mathcal{D}et D^2 v = 0$ and b) $v \in C_{loc}^{1,\alpha}(V)$. In order to show a), we first prove:

Lemma 4.6. *If $\alpha > 2/3$, we have for all $\psi \in C_c^\infty(V)$ and arbitrary i, j, k and l :*

$$\lim_{\varepsilon \rightarrow 0} \int_V A_{ij}^\varepsilon (\partial_k E_l^\varepsilon) \psi \rightarrow 0.$$

Proof. We have, using Proposition 4.1 and (4.8)

$$\begin{aligned} \left| \int_V A_{ij}^\varepsilon (\partial_k E_l^\varepsilon) \psi \right| &\leq \left| (A_{ij}^\varepsilon - A_{ij}) [(\partial_k E_l^\varepsilon) \psi] \right| + \left| A [(\partial_k E_l^\varepsilon) \psi] \right| \\ &\leq C\varepsilon^{2\alpha-1} \|(\partial_k E_l^\varepsilon) \psi\|_{W^{1,1}} + C \|(\partial_k E_l^\varepsilon) \psi\|_{W^{1,1}} \\ &\leq C \|\psi\|_1 \varepsilon^{2\alpha-1} \varepsilon^{3\alpha-2} + C \|\psi\|_1 \varepsilon^{3\alpha-2} \rightarrow 0. \end{aligned}$$

■

Now, we have, since $\nabla v^{(\varepsilon)} \rightarrow \nabla v$ in $L^2(V)$, for all $\psi \in C_c^\infty(V)$:

$$\begin{aligned}
-\frac{1}{2} \operatorname{curl}^T \operatorname{curl}(\nabla v \otimes \nabla v)[\psi] &= \lim_{\varepsilon \rightarrow 0} -\frac{1}{2} \operatorname{curl}^T \operatorname{curl}(\nabla v^{(\varepsilon)} \otimes \nabla v^{(\varepsilon)})[\psi] = \lim_{\varepsilon \rightarrow 0} \det \nabla^2 v^{(\varepsilon)}[\psi] \\
&= \lim_{\varepsilon \rightarrow 0} \int_V \det(A^\varepsilon + \nabla(E^\varepsilon)^\perp - \nabla^\perp E^\varepsilon) \psi \\
&= \lim_{\varepsilon \rightarrow 0} \int_V (\det A^\varepsilon) \psi + \lim_{\varepsilon \rightarrow 0} \sum_{\text{some indices } i,j,k,l} \int_V A_{ij}^\varepsilon (\partial_k E_l^\varepsilon) \psi \\
&\quad + \lim_{\varepsilon \rightarrow 0} \int_V \det(\nabla(E^\varepsilon)^\perp - \nabla^\perp E^\varepsilon) \psi.
\end{aligned}$$

The third term obviously converges to 0, and so does the second term by Lemma 4.6. The vanishing of the first term is less obvious, but faster. It directly follows from Proposition 4.3.

It remains to prove b) $v \in C_{loc}^{1,\alpha}(V)$. In view of (4.5) and since $\nabla^2 v = A$ is symmetric in i, j , we have $\partial_{ji} v = \partial_{ij} v = A_{ij}$, as distributions, and hence for all $\psi \in W_0^{1,1}(V)$:

$$\left| \int_V \partial_i v \partial_j \psi \right| = |A_{ij}[\psi]| \leq C \|\psi\|_{L^1(V)}^\alpha \|\partial_j \psi\|_{L^1(V)}^{1-\alpha}.$$

For any $x \in V$, we apply Proposition 3.3 to a coordinate rectangular box containing x and compactly included in V to conclude that $\partial_i v \in C_{loc}^{0,\alpha}(V)$ for $i = 1, 2$, which yields $v \in C_{loc}^{1,\alpha}(V)$. \blacksquare

5. DEVELOPABILITY: A PROOF OF THEOREM 1

In order to prove our main Theorem, and in view of Proposition 2.1, it is sufficient to show that ∇u satisfies condition (c). We first fix an open disk $V \ni x$ compactly contained in Ω and note the existence of the function $v \in C_{loc}^{1,\alpha}(V)$ as defined in Proposition 4.4. We apply the key developability result Theorem 2 to v to obtain that ∇v satisfies any of the three equivalent conditions of Proposition 2.1 in V . We apply Corollary 2.9 to ∇v , and choose accordingly the open disk B around x and the Lipschitz unit vector field $\vec{\eta}: B \rightarrow \mathbb{R}^2$. Therefore

$$(5.1) \quad \forall y \in B \quad \vec{\eta}(y) = \vec{\eta}(y + s\vec{\eta}(y)) \quad \text{and} \quad \nabla v(y) = \nabla v(y + s\vec{\eta}(y)) \quad \text{for all } s \text{ for which } y + s\vec{\eta}(y) \in B.$$

We claim that the Jacobian derivative ∇u of the isometric immersion u is indeed constant along the same directions defined by $\vec{\eta}$ in B . This would immediately imply that ∇u satisfies condition (c) of Proposition 2.1 and completes the proof.

In order to prove the claim, we consider the following identity [26] (also known as the Gauss equation in the literature), which is valid for the mollified sequence of smooth immersions u_ε :

$$(5.2) \quad \partial_{ij} u_\varepsilon = \Gamma_{ij}^{k,\varepsilon} \partial_k u_\varepsilon + A_{ij}^\varepsilon N^\varepsilon,$$

where the standard Einstein summation convention is used.

We apply Lemma 2.10 to $f = \partial_j v$ and $B = B_x$ to obtain for any $\psi \in C_c^\infty(B)$:

$$\int_B (\partial_j v) \operatorname{div}(\psi \vec{\eta}) = 0, \quad j = 1, 2.$$

We denote the coefficients of u and N^ε respectively by u^m and $N^{\varepsilon,m}$ and calculate for $m = 1, 2, 3$:

$$\begin{aligned}
(5.3) \quad \int_B \operatorname{div}(\psi \vec{\eta})(\nabla u^m) &= \lim_{\varepsilon \rightarrow 0} \int_B \operatorname{div}(\psi \vec{\eta})(\nabla u_\varepsilon^m) = \lim_{\varepsilon \rightarrow 0} - \int_B \psi \vec{\eta} \cdot \nabla^2 u_\varepsilon^m \\
&= \lim_{\varepsilon \rightarrow 0} - \int_B \psi \vec{\eta} \cdot \left(\Gamma^{k,\varepsilon} \partial_k u_\varepsilon^m + A^\varepsilon N^{\varepsilon,m} \right) = \lim_{\varepsilon \rightarrow 0} - \int_B \psi \vec{\eta} \cdot (A^\varepsilon N^{\varepsilon,m}),
\end{aligned}$$

where we used (4.7). Note that in view of $A = \nabla^2 v$, (5.1), and Lemma 2.10 we have for all ε , in view of the fact that $N^{\varepsilon,m}\psi \in C_c^\infty(B)$:

$$A[N^{\varepsilon,m}\psi\vec{\eta}] = - \int_B \operatorname{div}(N^{\varepsilon,m}\psi\vec{\eta})\nabla v = 0.$$

Hence by Proposition 4.1 we obtain:

$$\begin{aligned} \left| \int_B \psi\vec{\eta} \cdot (A^\varepsilon N^{\varepsilon,m}) \right| &= \left| A^\varepsilon [N^{\varepsilon,m}\psi\vec{\eta}] \right| = \left| (A^\varepsilon - A)[N^{\varepsilon,m}\psi\vec{\eta}] \right| + \left| A[N^{\varepsilon,m}\psi\vec{\eta}] \right| = \left| (A^\varepsilon - A)[N^{\varepsilon,m}\psi\vec{\eta}] \right| \\ &\leq C \left(\varepsilon^{2\alpha-1} \|N^{\varepsilon,m}\psi\vec{\eta}\|_{L^1} + \varepsilon^\alpha \|\nabla(N^{\varepsilon,m}\psi\vec{\eta})\|_{L^1} \right) \\ &\leq C \varepsilon^{2\alpha-1} \|\psi\vec{\eta}\|_{L^1} + \varepsilon^\alpha \|\nabla(\psi\vec{\eta})\|_{L^1} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

where we used (4.2). Therefore, (5.3) implies

$$\int_B \operatorname{div}(\psi\vec{\eta})(\nabla u^m) = 0.$$

Once again applying Lemma 2.10 and (5.1) implies that ∇u^m , $m = 1, 2, 3$, is constant along the segments generated by the vector field $\vec{\eta}$ in B . Our claim is proved and as already mentioned, this concludes the proof of Theorem 1 in view of Proposition 2.1.

6. APPENDIX: DEVELOPABILITY OF EACH COMPONENT CAN BE SHOWN INDEPENDENTLY

In this section we will directly show a weaker statement on the developability of $C^{1,\alpha}$ isometries:

Theorem 3. *Assume $\Omega \subset \mathbb{R}^2$ is an arbitrary domain, $\alpha > 2/3$ and let $u \in C_{loc}^{1,\alpha}(\Omega, \mathbb{R}^3)$ be an isometric immersion. Then for each $m = 1, 2, 3$, the component u^m is developable in Ω .*

Remark 6.1. Obviously this statement does not guarantee that the constancy lines or regions of ∇u^m are the same for the three components. If this fact is independently shown (e.g. through geometric arguments) Theorem 1 will also be proved through this approach.

Proof. In view of Theorem 2 and Proposition 2.1, it suffices to show that $\operatorname{Det} D^2 u^m = 0$ for each m . Once again, we consider the mollifications $u_\varepsilon = u * \phi_\varepsilon$ defined on a suitably chosen domain $V \Subset \Omega$ and use the (5.2) to calculate for $m=1,2,3$ and $\psi \in C_c^\infty(V)$:

$$\begin{aligned} -\frac{1}{2} \operatorname{curl}^T \operatorname{curl}(\nabla u^m \otimes \nabla u^m)[\psi] &= \lim_{\varepsilon \rightarrow 0} -\frac{1}{2} \operatorname{curl}^T \operatorname{curl}(\nabla u_\varepsilon^m \otimes \nabla u_\varepsilon^m)[\psi] = \lim_{\varepsilon \rightarrow 0} \det \nabla^2 u_\varepsilon^m[\psi] \\ &= \lim_{\varepsilon \rightarrow 0} \int_\Omega \det(\Gamma^{k,\varepsilon} \partial_k u_\varepsilon^m + A^\varepsilon N^{\varepsilon,m}) \psi \\ &= \lim_{\varepsilon \rightarrow 0} \int_\Omega \det(A^\varepsilon) (N^{\varepsilon,m})^2 \psi + \lim_{\varepsilon \rightarrow 0} \int_\Omega \det(\Gamma^{k,\varepsilon} \partial_k u_\varepsilon^m) \psi \\ &\quad + \lim_{\varepsilon \rightarrow 0} \sum_{\text{some indices } i,j,r,s} \int_\Omega A_{ij}^\varepsilon N^{\varepsilon,m} (\Gamma_{rs}^{k,\varepsilon} \partial_k u_\varepsilon^m) \psi. \end{aligned}$$

The second term obviously converges to 0 in view of (4.7). For the third term we use (4.1), (4.4) and (4.7) to prove:

$$\left| \int_\Omega A_{ij}^\varepsilon N^{\varepsilon,m} (\Gamma_{rs}^{k,\varepsilon} \partial_k u_\varepsilon^l) \psi \right| \leq C \|A^\varepsilon\|_{0;V} \|\Gamma^{k,\varepsilon}\|_{0;V} \|\psi\|_{L^1} \leq C \varepsilon^{3\alpha-2} \|\psi\|_{L^1} \rightarrow 0.$$

Finally, the vanishing of the first term follows from Propostion 4.3 and (4.2) since

$$\begin{aligned} \left| \int_{\Omega} \det(A^\varepsilon)(N^{\varepsilon,m})^2 \psi \right| &\leq C\varepsilon^{2\alpha-1} \|(N^{\varepsilon,m})^2 \psi\|_{W^{1,1}} \\ &\leq C\varepsilon^{2\alpha-1} \left(\|\psi\|_{W^{1,1}} + \|\nabla N^\varepsilon\|_{0;V} \|\psi\|_{L^1} \right) \\ &\leq C\varepsilon^{3\alpha-2} \|\psi\|_{W^{1,1}} \rightarrow 0. \end{aligned}$$

■

REFERENCES

- [1] D.N. Arnold, L.R. Scott and M. Vogelius, *Regular inversion of the divergence operator with Dirichlet boundary conditions on a polygon*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. Serie IV, **15** no. 2, (1988), 169–192.
- [2] H. Bahouri, J.Y. Chemin and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 343. Springer, Heidelberg, 2011.
- [3] H. Brezis, and H.-M. Nguyen, *The Jacobian determinant revisited*, Invent. Math. **185** no. 1 (2011), 17–54.
- [4] Yu. F. Borisov, *The parallel translation on a smooth surface. III.*, Vestnik Leningrad. Univ. **14** no. 1, (1959), 34–50.
- [5] Yu. F. Borisov, *Irregular surfaces of the class $C^{1,\beta}$ with an analytic metric*, (Russian) Sibirsk. Mat. Zh. **45** no. 1, (2004), 25–61; translation in Siberian Math. J. **45** no. 1, (2004), 19–52.
- [6] F. Cajori, *Generalisations in Geometry as Seen in the History of Developable Surfaces*, The American Mathematical Monthly **36**, 8, (1929), 431–437.
- [7] W. Cao and L. Székelyhidi Jr, *Very weak solutions to the two-dimensional Monge-Ampère equation*, Science China Mathematics, **62** (6), (2019) 1041–1056.
- [8] M.P. do Carmo, *Differential geometry of curves and surfaces*, Translated from the Portuguese. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1976.
- [9] S.S. Chern and R.K. Lashof, *On the Total Curvature of Immersed Manifolds*, American Journal of Mathematics, **79** No. 2., (1957), 306–318.
- [10] P. Constantin, W. E and E. S. Titi, *Onsager’s conjecture on the energy conservation for solutions of Euler’s equation*, Comm. Math. Phys. **165** no. 1, (1994), 207–209.
- [11] S. Conti, C. De Lellis and L. Székelyhidi Jr., *h-principle and rigidity for $C^{1,\alpha}$ isometric embeddings*, Nonlinear Partial Differential Equations, Proceedings of the Abel Symposium, Abel Symposia Volume **7**, (2012), 83–116.
- [12] S. Conti and F. Maggi, *Confining thin sheets and folding paper*, Arch. Ration. Mech. Anal. **187** no. 1, (2008), 1–48
- [13] C. De Lellis, D. Inauen, *$C^{1,\alpha}$ isometric embeddings of polar caps*, Preprint (2018), <https://arxiv.org/pdf/1809.04161>.
- [14] C. De Lellis, D. Inauen and L. Székelyhidi Jr., *A Nash-Kuiper theorem for $C^{1,\frac{1}{5}-\delta}$ immersions of surfaces in 3 dimensions*, Revista matemática iberoamericana, **34** (3), (2018) 1119–1152.
- [15] C. De Lellis and L. Székelyhidi Jr., *Dissipative continuous Euler flows*, Invent. Math. **193** no. 2, (2013), 377–407.
- [16] C. De Lellis and L. Székelyhidi Jr., *High dimensionality and h-principle in PDE*, Bull. Amer. Math. Soc. (N.S.) **54** no. 2 (2017), 247–282.
- [17] C. De Lellis and L. Székelyhidi Jr., *On turbulence and geometry: from Nash to Onsager*, Notices Amer. Math. Soc. **66** no. 5, (2019), 677–685.
- [18] G. L. Eyink, *Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer*. Phys. D, **78** (1994) no. 3-4, 222–240.
- [19] I. Fonseca and J. Malý, *From Jacobian to Hessian: distributional form and relaxation*, Riv. Mat. Univ. Parma **7** 4*, (2005), 45–74.
- [20] M. Friedman, *A history of folding in mathematics: Mathematizing the margins*, Science Networks. Historical Studies, 59. Birkhäuser/Springer, Cham, 2018.
- [21] G. Friesecke, R.D. James and S. Müller, *A hierarchy of plate models, derived from nonlinear elasticity by Gamma-convergence*, Arch. Rational Mech. Anal. **180**, (2006), 183–236.
- [22] P. Friz and M. Hairer, *A course on rough paths. With an introduction to regularity structures*, Universitext. Springer, Cham, 2014.
- [23] J.H.G. Fu, *Monge-Ampère functions I*, Indiana Univ. Math. Jour., **38** no. 3, (1989), 745–771.

- [24] M. Gromov, *Partial differential relations*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 9. Springer-Verlag, Berlin, 1986.
- [25] M. Gromov, *Geometric, algebraic, and analytic descendants of Nash isometric embedding theorems*, Bull. Amer. Math. Soc., **54** (2017), 173–245.
- [26] Q. Han and J.-X. Hong, *Isometric embedding of Riemannian manifolds in Euclidean spaces*, Mathematical Surveys and Monographs, 130, American Mathematical Society, Providence, RI, (2006).
- [27] P. Hartman and L. Nirenberg, *On spherical image maps whose Jacobians do not change sign*, Amer. J. Math., **81** (1959), 901–920.
- [28] P. Hornung, *Fine level set structure of flat isometric immersions*, Arch. Rational Mech. Anal., **199** (2011), 943–1014.
- [29] P. Hornung, *Approximation of flat $W^{2,2}$ isometric immersions by smooth ones*, Arch. Rational Mech. Anal., **199** (2011), 1015–1067.
- [30] P. Hornung and I. Velčić, *Regularity of intrinsically convex H^2 surfaces and the derivation of homogenized bending shell models*, Journal de Mathématiques Pures et Appliquées **115**, (2018), 1–23.
- [31] P. Isett, *A proof of Onsager’s conjecture* Annals of Mathematics, **188** No. 3, (2018), 871–963.
- [32] T. Iwaniec, *On the concept of weak Jacobian and Hessian*, Report Univ. Jyväskylä **83**, (2001), 181–205.
- [33] R.L. Jerrard, *Some remarks on Monge-Ampère functions*, Singularities in PDE and the calculus of variations, CRM Proc. Lecture Notes, **44**, (2008), 89–112.
- [34] R.L. Jerrard, *Some rigidity results related to Monge-Ampère functions*, Canad. J. Math. **62** no. 2, (2010), 320–354
- [35] R.L. Jerrard and M.R. Pakzad, *Sobolev Spaces of isometric immersions of arbitrary dimension and co-dimension*, Ann. Mat. Pura Appl. (4), **196** no. 2, (2017), 687–716.
- [36] B. Kirchheim, *Geometry and rigidity of microstructures*, habilitation thesis, University of Leipzig, (2001). (MPI-MIS Lecture Notes 16/ 2003 <http://www.mis.mpg.de/preprints/ln/index.html>).
- [37] M.V. Korobkov, *Properties of the C^1 -smooth functions with nowhere dense gradient range*, Sibirsk. Mat. Zh. **48** no. 6, (2007), 1272–1284; translation in Siberian Math. J. **48** no. 6, (2007), 1019–1028
- [38] M.V. Korobkov, *Properties of C^1 -smooth mappings with one-dimensional gradient range* Sibirsk. Mat. Zh. **50** no. 5, (2009), 874–886; translation in Siberian Math. J. **50** no. 5, (2009), 874–886.
- [39] N. H. Kuiper, *On C^1 -isometric imbeddings. I, II.*, Nederl. Akad. Wetensch. Proc. Ser. A. **58**, (1955) 545–556, 683–689.
- [40] H. Lebesgue, *Sur quelques surfaces non réglés applicables sur le plan*, Comptes Rendus de l’Académie des Sciences CXXVIII, (1899), 1502–1505.
- [41] M. Lewicka, L. Mahadevan and M.R. Pakzad, *The Monge-Ampère constraint: matching of isometries, density and regularity and elastic theories of shallow shells*, Annales de l’Institut Henri Poincaré (C) Non Linear Analysis, Vol **34** Issue 1, (2017), 45–67.
- [42] M. Lewicka and M.R. Pakzad, *Prestrained elasticity: from shape formation to Monge-Ampère anomalies*, Notices of the AMS, vol. **63** no. 1, (January 2016).
- [43] M. Lewicka and M.R. Pakzad, *Convex integration for the Monge-Ampère equation in two dimensions*, Analysis and PDE, **10** No. 3, (2017), 695–727.
- [44] Z. Liu and M.R. Pakzad, *Regularity, rigidity and density of smooth mappings for co-dimension 1 Sobolev isometric immersions*, Ann. Sc. Norm. Sup. Pisa - Classe di Scienze. Vol. XIV (3), (2015), 76–817.
- [45] J. Malý and O. Martio, *Lusin’s condition (N) and mappings of the class $W^{1,n}$* , J. Reine Angew. Math., **458**, (1995), 19–36.
- [46] S. Müller and M.R. Pakzad, *Regularity properties of isometric immersions*, Math. Zeit. **251** (2), (2005), 313–331.
- [47] J. Nash, *The imbedding problem for Riemannian manifolds*, Ann. Math., **63**, (1956), 20–63.
- [48] J. Nash, *C^1 isometric imbeddings*, Ann. Math., **60**, (1954), 383–396.
- [49] L. Onsager, *Statistical hydrodynamics. II*, Nuovo Cimento (1943-1954), **6** (2), (1949), 279–287.
- [50] M.R. Pakzad, *On the Sobolev space of isometric immersions*, Journal of Differential Geometry, **66**, (2004), 47–69.
- [51] E. Picard, *Traité d’analyse*, 3rd ed. Paris: Gauthier-Villars et fils, 1922.
- [52] A.V. Pogorelov, *Extrinsic geometry of convex surfaces*, Translations of mathematical monographs vol. 35, American Math. Soc., 1973.
- [53] D. Spring, *Convex integration theory. Solutions to the h-principle in geometry and topology*, Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 2010.

- [54] L. Székelyhidi, *From Isometric Embeddings to Turbulence*, MPI-MIS Lecture note 41/2012, <http://www.mis.mpg.de/publications/other-series/ln/lecturenote-4112.html>.
- [55] S. C. Venkataramani, T. A. Witten, E. M. Kramer, and R. P. Geroch, *Limitations on the smooth confinement of an unstretchable manifold*, J. Math. Phys. **41** (7), (2000), 5107–5128.

CAMILLO DE LELLIS, SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, 1 EINSTEIN DR., PRINCETON NJ 05840, USA

Email address: `camillo.delellis@math.ias.edu`

MOHAMMAD REZA PAKZAD, IAM, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, BONN 53111, GERMANY, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PA, USA

Email address: `pakzad@iam.uni-bonn.de`, `pakzad@pitt.edu`