

ONSAGER'S CONJECTURE FOR ADMISSIBLE WEAK SOLUTIONS

TRISTAN BUCKMASTER, CAMILLO DE LELLIS, LÁSZLÓ SZÉKELYHIDI JR., AND VLAD VICOL

ABSTRACT. We prove that given any $\beta < 1/3$, a time interval $[0, T]$, and given any smooth energy profile $e: [0, T] \rightarrow (0, \infty)$, there exists a weak solution v of the three-dimensional Euler equations such that $v \in C^\beta([0, T] \times \mathbb{T}^3)$, with $e(t) = \int_{\mathbb{T}^3} |v(x, t)|^2 dx$ for all $t \in [0, T]$. Moreover, we show that a suitable h -principle holds in the regularity class $C_{t,x}^\beta$, for any $\beta < 1/3$. The implication of this is that the dissipative solutions we construct are in a sense typical in the appropriate space of subsolutions as opposed to just isolated examples.

1. INTRODUCTION

In this paper we consider the incompressible Euler equations

$$\begin{cases} \partial_t v + v \cdot \nabla v + \nabla p = 0 \\ \operatorname{div} v = 0, \end{cases} \quad (1.1)$$

in the periodic setting $x \in \mathbb{T}^3 = \mathbb{R}^3 \setminus \mathbb{Z}^3$, where v is a vector field representing the velocity of the fluid and p is the pressure. We study weak (distributional) solutions v which are Hölder continuous in space, i.e. such that¹

$$|v(x, t) - v(y, t)| \leq C|x - y|^\beta \quad \text{for all } t \in [0, T] \quad (1.2)$$

for some constant C which is independent of time t .

In his famous 1949 note on statistical hydrodynamics Lars Onsager [Ons49] conjectured that the threshold regularity for the validity of the energy conservation of weak solutions of (1.1) is the exponent $1/3$: in particular he asserted that for larger Hölder exponents any weak solution would conserve the energy, whereas for any smaller exponent there are solutions which do not. The first assertion was fully proved by Constantin, E and Titi in [CET94], after a partial result of Eyink in [Eyi94] (see also [CCFS08] for a sharper criterion in L^3 -based spaces). Concerning the second assertion, the first proof of the existence of a square summable weak solution which does not preserve the energy is due to Scheffer in his pioneering paper [Sch93]. A different proof has been later given by Shnirelman in [Shn97]. In [DLS09] the second and third author realized that techniques from the theory of differential inclusions could be applied very efficiently to produce bounded weak solutions which violate the energy conservation in several forms (see also [DL10, Chi14, LXX16], [CFG11, Shv11, Szé12, CCF16], and [CS14] for related work on the compressible Euler equations, non-odd active scalar equations, and the stationary Euler equations, respectively). Pushed by the analogy of these constructions with the famous C^1 solutions of Nash and Kuiper for the isometric embedding problem (cf. [Nas54] and [Kui55]), the second and third authors proposed to approach the remaining statement of the Onsager's conjecture in a similar way (cf. [DLS12]). Indeed in [DLS13] and [DLS14] they were able to give the first examples of, respectively, continuous and Hölder continuous solutions which dissipate the energy, reaching the threshold exponent $1/10$ (see

Date: October 16, 2021.

¹The smallest constant C satisfying (1.2) will be denoted by $[v]_\beta$, cf. Appendix A. We will write $v \in C^\beta(\mathbb{T}^3 \times [0, T])$ when v is Hölder continuous in the whole space-time.

also [IV15, BSV16] for constructions of Hölder continuous weak solutions for other hydrodynamic models such as the IPM and SQG equations). After a series of important partial results improving the threshold and the techniques from several points of view, cf. [Ise13a, BDLS13, Buc14, BDLISJ15, IO16, Buc15, BDLS16], in his recent paper Isett [Ise16] has been able to finally reach the Onsager exponent $1/3$. The proof of Isett combines previous ideas with two new important ingredients, one developed by Daneri and the third author in [DS17] (the introduction of Mikado flows, see Section 2.6) and one introduced by Isett himself in the aforementioned paper (the gluing technique, see Section 2.5).

However, the solutions produced in [Ise16] are only shown to be nonconservative and in fact for those solutions the total kinetic energy fails to be monotonic on any interval of time. Thus Isett's theorem left open the question whether it is possible or not to construct solutions which *dissipate* the kinetic energy (i.e. with strictly monotonic decreasing energy). In fact the latter is a relevant point for at least two reasons: dissipative solutions satisfy the weak-strong uniqueness property [Lio96, BDLS11], and Onsager, in his work, conjectures the existence of dissipative solutions. Indeed, in [Ons49] Onsager states:

It is of some interest to note that in principle, turbulent dissipation as described could take place just as readily without the final assistance by viscosity.

In this note we suitably modify the approach of Isett in order to show the following theorem.

Theorem 1.1. *Assume $e : [0, T] \rightarrow \mathbb{R}$ is a strictly positive smooth function. Then for any $0 < \beta < 1/3$ there exists a weak solution $v \in C^\beta(\mathbb{T}^3 \times [0, T])$ to (1.1) such that*

$$\int_{\mathbb{T}^3} |v(x, t)|^2 dx = e(t).$$

We are indeed able to prove a stronger statement than Theorem 1.1, namely an h -principle in the sense of [DS17]. Following [DS17] we introduce smooth strict subsolutions of the Euler equations.

Definition 1.2. *A smooth strict subsolution of (1.1) on $\mathbb{T}^3 \times [0, T]$ is a smooth triple $(\bar{v}, \bar{p}, \bar{R})$ with \bar{R} a symmetric 2-tensor, such that*

$$\begin{cases} \partial_t \bar{v} + \operatorname{div}(\bar{v} \otimes \bar{v}) + \nabla \bar{p} = -\operatorname{div} \bar{R} \\ \operatorname{div} \bar{v} = 0, \end{cases} \quad (1.3)$$

and $\bar{R}(x, t)$ is positive definite for all (x, t) .

We then can prove that any smooth strict subsolution can be suitably approximated by C^β solutions for any $\beta < 1/3$. More precisely:

Theorem 1.3. *Let $(\bar{v}, \bar{p}, \bar{R})$ be a smooth strict subsolution of the Euler equations on $\mathbb{T}^3 \times [0, T]$ and let $\beta < 1/3$. Then there exists a sequence (v_k, p_k) of weak solutions of (1.1) such that $v_k \in C^\beta(\mathbb{T}^3 \times [0, T])$,*

$$v_k \xrightarrow{*} \bar{v} \quad \text{and} \quad v_k \otimes v_k \xrightarrow{*} \bar{v} \otimes \bar{v} + \bar{R} \quad \text{in} \quad L^\infty$$

uniformly in time, and furthermore for all $t \in [0, T]$

$$\int_{\mathbb{T}^3} |v_k|^2 dx = \int_{\mathbb{T}^3} (|\bar{v}|^2 + \operatorname{tr} \bar{R}) dx. \quad (1.4)$$

Theorem 1.1 can be concluded as a simple corollary of Theorem 1.3. However, we give an alternative, simpler and self-contained argument for Theorem 1.1. Indeed the proof of Theorem 1.3 invokes some results of [DS17], whereas the argument for Theorem 1.1 is entirely contained in our note,

aside from technical propositions which are classical statements in the literature, all collected in the Appendix.

The most important differences in our proof compared to that of [Ise16] rely on the estimates for the “gluing step” of Isett’s proof (we refer to Section 2.5 for more details) and in a simple remark concerning the regions where the perturbation is added (see Section 2.6). We note that, even without the extra benefit of imposing the energy profile and achieving the more general h -principle statement, the proof proposed here is considerably shorter than that of [Ise16].

Acknowledgments. The work of T.B. has been partially supported by the National Science Foundation grant DMS-1600868. The research of C.D.L. has been supported by the grant 200021_159403 of the Swiss National Foundation. L.Sz. gratefully acknowledges the support of the ERC Grant Agreement No. 277993. V.V. was partially supported by the National Science Foundation grant DMS-1514771 and by an Alfred P. Sloan Research Fellowship.

2. OUTLINE OF THE PROOF

As already mentioned, although Theorem 1.1 can be recovered as a corollary of Theorem 1.3, in this section we outline an independent proof, reducing it to a suitable iterative procedure, summarized in Proposition 2.1 below. The same iteration procedure can be used to prove Theorem 1.3, as shown in Section 7 at the end of the note, but the corresponding argument will need some results from [DS17], which we state without proof. In contrast, the proof of Theorem 1.1 is completely self-contained.

2.1. Inductive proposition. First of all, we impose for the moment that

$$\sup_{t \in [0, T]} \left| \frac{d}{dt} e(t) \right| \leq 1 \quad (2.1)$$

(we will see later that this can be done without losing generality).

Let then $q \geq 0$ be a natural number. At a given step q we assume to have a triple $(v_q, p_q, \mathring{R}_q)$ to the Euler-Reynolds system (1.3), namely such that

$$\begin{cases} \partial_t v_q + \operatorname{div}(v_q \otimes v_q) + \nabla p_q = \operatorname{div} \mathring{R}_q \\ \operatorname{div} v_q = 0, \end{cases} \quad (2.2)$$

to which we add the constraints that

$$\operatorname{tr} \mathring{R}_q = 0 \quad (2.3)$$

and that

$$\int_{\mathbb{T}^3} p_q(x, t) dx = 0 \quad (2.4)$$

(which uniquely determines the pressure).

The size of the approximate solution v_q and the error \mathring{R}_q will be measured by a frequency λ_q and an amplitude δ_q , which are given by

$$\lambda_q = 2\pi \lceil a^{(b^q)} \rceil \quad (2.5)$$

$$\delta_q = \lambda_q^{-2\beta} \quad (2.6)$$

where $\lceil x \rceil$ denotes the smallest integer $n \geq x$, $a > 1$ is a large parameter, $b > 1$ is close to 1 and $0 < \beta < 1/3$ is the exponent of Theorem 1.1. The parameters a and b are then related to β .

We proceed by induction, assuming the estimates

$$\left\| \mathring{R}_q \right\|_0 \leq \delta_{q+1} \lambda_q^{-3\alpha} \quad (2.7)$$

$$\|v_q\|_1 \leq M \delta_q^{1/2} \lambda_q \quad (2.8)$$

$$\|v_q\|_0 \leq 1 - \delta_q^{1/2} \quad (2.9)$$

$$\delta_{q+1} \lambda_q^{-\alpha} \leq e(t) - \int_{\mathbb{T}^3} |v_q|^2 dx \leq \delta_{q+1} \quad (2.10)$$

where $0 < \alpha < 1$ is a small parameter to be chosen suitably (which will depend upon β), and M is a universal constant (which is fixed throughout the iteration and whose choice depends on certain geometric properties of the space of symmetric matrices and on the “squiggling” regions of the perturbation step, cf. Remark 5.2, Lemma 5.5 and Definition 5.6). We refer to Appendix A for the definitions of the Hölder norms used above, where we take into account only *space regularity*.

Proposition 2.1. *There is a universal constant M with the following property. Assume $0 < \beta < 1/3$ and*

$$1 < b < \frac{1 - \beta}{2\beta}. \quad (2.11)$$

Then there exists an α_0 depending on β and b , such that for any $0 < \alpha < \alpha_0$ there exists an a_0 depending on β , b , α and M , such that for any $a \geq a_0$ the following holds: Given a strictly positive energy function $e : [0, T] \rightarrow \mathbb{R}$ satisfying (2.1), and a triple $(v_q, \mathring{R}_q, p_q)$ solving (2.2)–(2.4) and satisfying the estimates (2.7)–(2.10), then there exists a solution $(v_{q+1}, \mathring{R}_{q+1}, p_{q+1})$ to (2.2)–(2.4) satisfying (2.7)–(2.10) with q replaced by $q + 1$. Moreover, we have

$$\|v_{q+1} - v_q\|_0 + \frac{1}{\lambda_{q+1}} \|v_{q+1} - v_q\|_1 \leq M \delta_{q+1}^{1/2}. \quad (2.12)$$

The proof of Proposition 2.1 is summarized in the Sections 2.3, 2.4, 2.5 and 2.6, but its details will occupy most of the paper and will be completed in Section 6 below. We show next that this proposition immediately implies Theorem 1.1.

2.2. Proof of Theorem 1.1. First of all, we fix any Hölder exponent $\beta < 1/3$ and also the parameters b and α , the first satisfying (2.11) and the second smaller than the threshold given in Proposition 2.1. Next we show that, without loss of generality, we may further assume the energy profile satisfies

$$\inf_t e(t) \geq \delta_1 \lambda_0^{-\alpha}, \quad \sup_t e(t) \leq \delta_1, \quad \text{and} \quad \sup_t e'(t) \leq 1, \quad (2.13)$$

provided the parameter a is chosen sufficiently large.

To see this, we first note that the Euler equations are invariant under the transformation

$$v(x, t) \mapsto \Gamma v(x, \Gamma t) \quad \text{and} \quad p(x, t) \mapsto \Gamma^2 p(x, \Gamma t).$$

Thus if we choose

$$\Gamma = \left(\frac{\delta_1}{\sup_t e(t)} \right)^{1/2},$$

then using the scaling invariance, the stated problem reduces to finding a solution with the energy profile given by

$$\tilde{e}(t) = \Gamma^2 e(\Gamma t),$$

for which we have

$$\inf_t \tilde{e}(t) \geq \frac{\delta_1 \inf_t e(t)}{\sup_t e(t)}, \quad \sup_t \tilde{e}(t) \leq \delta_1, \quad \text{and} \quad \sup_t \tilde{e}'(t) \leq \left(\frac{\delta_1}{\sup_t e(t)} \right)^{3/2} \sup_t e'(t).$$

If a is chosen sufficiently large then we can ensure

$$\sup_t \tilde{e}'(t) \leq \left(\frac{\delta_1}{\sup_t e(t)} \right)^{3/2} \sup_t e'(t) \leq 1, \quad \text{and} \quad \frac{\inf_t e(t)}{\sup_t e(t)} \geq \lambda_0^{-\alpha}.$$

Now we apply Proposition 2.1 iteratively with $(v_0, R_0, p_0) = (0, 0, 0)$. Indeed the pair (v_0, R_0) trivially satisfies (2.7)–(2.9), whereas the estimate (2.10) and (2.1) follows as a consequence of (2.13). Notice that by (2.12) v_q converges uniformly to some continuous v . Moreover, we recall that the pressure is determined by

$$\Delta p_q = \operatorname{div} \operatorname{div}(-v_q \otimes v_q + \mathring{R}_q) \quad (2.14)$$

and (2.4) and thus p_q is also converging to some pressure p (for the moment only in L^r for every $r < \infty$). Since $\mathring{R}_q \rightarrow 0$ uniformly, the pair (v, p) solves the Euler equations.

Observe that using (2.12) we also infer²

$$\sum_{q=0}^{\infty} \|v_{q+1} - v_q\|_{\beta'} \lesssim \sum_{q=0}^{\infty} \|v_{q+1} - v_q\|_0^{1-\beta'} \|v_{q+1} - v_q\|_1^{\beta'} \lesssim \sum_{q=0}^{\infty} \delta_{q+1}^{\frac{1-\beta'}{2}} \left(\delta_{q+1}^{1/2} \lambda_q \right)^{\beta'} \lesssim \sum_{q=0}^{\infty} \lambda_q^{\beta' - \beta}$$

and hence that v_q is uniformly bounded in $C_t^0 C_x^{\beta'}$ for all $\beta' < \beta$. To recover the time regularity, we could use the Euler equations and the general result in [Ise13b]. Nevertheless, we believe that the following short and self-contained proof of the time-regularity may be of independent interest:

Fix a smooth standard mollifier ψ in space, let $q \in \mathbb{N}$, and consider $\tilde{v}_q := v * \psi_{2^{-q}}$, where $\psi_\ell(x) = \ell^{-3} \psi(x\ell^{-1})$. From standard mollification estimates we have

$$\|\tilde{v}_q - v\|_0 \lesssim \|v\|_{\beta'} 2^{-q\beta'}, \quad (2.15)$$

and thus $\tilde{v}_q - v \rightarrow 0$ uniformly as $q \rightarrow \infty$. Moreover, \tilde{v}_q obeys the following equation

$$\partial_t \tilde{v}_q + \operatorname{div}(v \otimes v) * \psi_{2^{-q}} + \nabla p * \psi_{2^{-q}} = 0.$$

Next, since

$$-\Delta p * \psi_{2^{-q}} = \operatorname{div} \operatorname{div}(v \otimes v) * \psi_{2^{-q}},$$

using Schauder's estimates, for any fixed $\varepsilon > 0$ we get

$$\|\nabla p * \psi_{2^{-q}}\|_0 \leq \|\nabla p * \psi_{2^{-q}}\|_\varepsilon \lesssim \|v \otimes v\|_{\beta'} 2^{q(1+\varepsilon-\beta')} \lesssim \|v\|_{\beta'}^2 2^{q(1+\varepsilon-\beta')},$$

(where the constant in the estimate depends on ε but not on q). Similarly,

$$\|(v \otimes v) * \psi_{2^{-q}}\|_1 \lesssim \|v \otimes v\|_{\beta'} 2^{q(1-\beta')} \lesssim \|v\|_{\beta'}^2 2^{q(1-\beta')}.$$

Thus the above estimates yield

$$\|\partial_t \tilde{v}_q\|_0 \lesssim \|v\|_{\beta'}^2 2^{q(1+\varepsilon-\beta')}. \quad (2.16)$$

Next, for $\beta'' < \beta'$ we conclude from (2.15) and (2.16) that

$$\begin{aligned} \|\tilde{v}_q - \tilde{v}_{q+1}\|_{C_x^0 C_t^{\beta''}} &\lesssim (\|\tilde{v}_q - v\|_0 + \|\tilde{v}_{q+1} - v\|_0)^{1-\beta''} (\|\partial_t \tilde{v}_q\|_0 + \|\partial_t \tilde{v}_{q+1}\|_0)^{\beta''} \\ &\lesssim \|v\|_{\beta'}^{1+\beta''} 2^{-q\beta'(1-\beta'')} 2^{q\beta''(1+\varepsilon-\beta')} = \|v\|_{\beta'}^{1+\beta''} 2^{-q(\beta'-(1+\varepsilon)\beta'')} \\ &\lesssim \|v\|_{\beta'}^{1+\beta''} 2^{-q\varepsilon} \end{aligned}$$

²Throughout the manuscript we use the notation $x \lesssim y$ to denote $x \leq Cy$, for a sufficiently large constant $C > 0$, which is independent of a, b , and q , but may change from line to line.

Here we have chosen $\varepsilon > 0$ sufficiently small (in terms of β' and β'') so that that $\beta' - (1 + \varepsilon)\beta'' \geq \varepsilon$. Thus, the series

$$v = \tilde{v}_0 + \sum_{q \geq 0} (\tilde{v}_{q+1} - \tilde{v}_q)$$

converges in $C_x^0 C_t^{\beta''}$. Since we already know $v \in C_t^0 C_x^{\beta'}$, we obtain that $v \in C^{\beta''}([0, T] \times \mathbb{T}^3)$ as desired, with $\beta'' < \beta' < \beta < 1/3$ arbitrary.

Finally, since $\delta_{q+1} \rightarrow 0$ as $q \rightarrow \infty$, from (2.10) we have

$$\int_{\mathbb{T}^3} |v|^2 dx = e(t),$$

which completes the proof of the theorem.

2.3. Stages. Except for Section 7, the rest of the paper is devoted to the proof of Proposition 2.1. It will be useful to make the assumption that α is small enough so to have

$$\lambda_q^{3\alpha} \leq \left(\frac{\delta_q}{\delta_{q+1}} \right)^{3/2} \leq \frac{\lambda_{q+1}}{\lambda_q}, \quad (2.17)$$

which also require that a is large enough to absorb any constant appearing from the ratio $\lambda_q/a^{(b^q)}$, for which we have the elementary bounds

$$2\pi \leq \frac{\lambda_q}{a^{b^q}} \leq 4\pi. \quad (2.18)$$

The proof consists of three stages, in each of which we modify v_q . Roughly speaking, the stages are as follows:

- Mollification: $(v_q, \mathring{R}_q) \mapsto (v_\ell, \mathring{R}_\ell)$;
- Gluing: $(v_\ell, \mathring{R}_\ell) \mapsto (\bar{v}_q, \bar{\mathring{R}}_q)$;
- Perturbation: $(\bar{v}_q, \bar{\mathring{R}}_q) \mapsto (v_{q+1}, \mathring{R}_{q+1})$.

2.4. Mollification step. The first stage is mollification: we mollify v_q at length scale ℓ in order to handle the *loss of derivative* problem, typical of convex integration schemes. To this aim, we fix a standard mollification kernel ψ *in space* and introduce the mollification parameter

$$\ell := \frac{\delta_{q+1}^{1/2}}{\delta_q^{1/2} \lambda_q^{1+3\alpha/2}}, \quad (2.19)$$

and define

$$\begin{aligned} v_\ell &:= v_q * \psi_\ell \\ \mathring{R}_\ell &:= \mathring{R}_q * \psi_\ell - (v_q \otimes v_q) * \psi_\ell + v_\ell \otimes v_\ell \end{aligned}$$

where $f \otimes g$ is the traceless part of the tensor $f \otimes g$. These functions obey the equation

$$\begin{cases} \partial_t v_\ell + \operatorname{div}(v_\ell \otimes v_\ell) + \nabla p_\ell = \operatorname{div} \mathring{R}_\ell \\ \operatorname{div} v_\ell = 0, \end{cases} \quad (2.20)$$

in view of (2.2).

Observe, again choosing α sufficiently small and a sufficiently large we can assume

$$\lambda_q^{-3/2} \leq \ell \leq \lambda_q^{-1}, \quad (2.21)$$

which will be applied repeatedly in order to simplify the statements of several estimates.

From (2.21), standard mollification estimates and Proposition A.2, we obtain the following bounds³

Proposition 2.2.

$$\|v_\ell - v_q\|_0 \lesssim \delta_{q+1}^{1/2} \lambda_q^{-\alpha}, \quad (2.22)$$

$$\|v_\ell\|_{N+1} \lesssim \delta_q^{1/2} \lambda_q \ell^{-N} \quad \forall N \geq 0, \quad (2.23)$$

$$\left\| \mathring{R}_\ell \right\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N+\alpha} \quad \forall N \geq 0. \quad (2.24)$$

$$\left| \int_{\mathbb{T}^3} |v_q|^2 - |v_\ell|^2 dx \right| \lesssim \delta_{q+1} \ell^\alpha. \quad (2.25)$$

Proof of Proposition 2.2. The bounds (2.22) and (2.23) follow from the obvious estimates

$$\|v_\ell - v_q\|_0 \leq \|v_q\|_1 \ell \lesssim \delta_q^{1/2} \lambda_q \ell \lesssim \delta_{q+1}^{1/2} \lambda_q^{-\alpha}$$

and

$$\|v_\ell\|_{N+1} \leq \|v_q\|_1 \ell^{-N} \lesssim \delta_q^{1/2} \lambda_q \ell^{-N}.$$

Next, applying Proposition A.2,

$$\left\| \mathring{R}_\ell \right\|_{N+\alpha} \lesssim \|\mathring{R}_q\|_0 \ell^{-N-\alpha} + \|v_q\|_1^2 \ell^{2-N-\alpha} \lesssim \delta_{q+1} \lambda_q^{-3\alpha} \ell^{-N-\alpha} + \delta_q \lambda_q^2 \ell^2 \ell^{-N-\alpha} \lesssim \delta_{q+1} \lambda_q^{-3\alpha} \ell^{-N-\alpha},$$

on the other hand, by (2.21) $\lambda_q^{-3\alpha} \leq \ell^{2\alpha}$, from which (2.24) follows. Similarly, by Proposition A.2,

$$\left| \int_{\mathbb{T}^3} |v_q|^2 - |v_\ell|^2 dx \right| = \left| \int_{\mathbb{T}^3} (|v_q|^2)_\ell - |v_\ell|^2 dx \right| \lesssim \left\| (|v_q|^2)_\ell - |v_\ell|^2 \right\|_0 \lesssim \|v_q\|_1^2 \ell^2,$$

which implies (2.25). \square

2.5. Gluing step. In the second stage we encounter the new crucial ingredient introduced by Isett in [Ise16]: we glue together exact solutions to the Euler equations in order to produce a new \bar{v}_q , close to v_q , whose associated Reynolds stress error has support in pairwise disjoint temporal regions of length τ_q in time, where

$$\tau_q = \frac{\ell^{2\alpha}}{\delta_q^{1/2} \lambda_q}. \quad (2.26)$$

The parameter τ_q should be compared to the parameter μ^{-1} used in the paper [BDLISJ15]. Indeed, τ_q^{-1} satisfies precisely the same parameter inequalities that μ satisfies in Section 2 of [BDLISJ15]. We note in particular that like in [BDLISJ15] we have the CFL-like condition

$$\tau_q \|v_\ell\|_{1+\alpha} \stackrel{(2.23)}{\lesssim} \tau_q \delta_q^{1/2} \lambda_q \ell^{-\alpha} \lesssim \ell^\alpha \ll 1 \quad (2.27)$$

as long as a is sufficiently large.

More precisely, we aim to construct a new triple $(\bar{v}_q, \mathring{\bar{R}}_q, \bar{p}_q)$ solving the Euler Reynolds equation (2.2) such that the temporal support of $\mathring{\bar{R}}_q$ is contained in pairwise disjoint intervals I_i of length $\sim \tau_q$ and such that the gaps between neighbouring intervals is also of length $\sim \tau_q$. More precisely, for any $n \in \mathbb{Z}$ let

$$t_n = n\tau_q, \quad I_n = [t_n + \frac{1}{3}\tau_q, t_n + \frac{2}{3}\tau_q] \cap [0, T], \quad J_n = (t_n - \frac{1}{3}\tau_q, t_n + \frac{1}{3}\tau_q) \cap [0, T].$$

³In the following, when considering higher order norms $\|\cdot\|_N$ or $\|\cdot\|_{N+1}$, the symbol \lesssim will imply that the constant in the inequality might also depend on N .

We require

$$\text{supp } \overset{\circ}{\bar{R}}_q \subset \bigcup_{n \in \mathbb{N}} I_n \times \mathbb{T}^3. \quad (2.28)$$

Moreover, $(\bar{v}_q, \overset{\circ}{\bar{R}}_q)$ will satisfy the following estimates for any $N \geq 0$

$$\|\bar{v}_q - v_\ell\|_0 \lesssim \delta_{q+1}^{1/2} \ell^\alpha \quad (2.29)$$

$$\|\bar{v}_q\|_{1+N} \lesssim \delta_q^{1/2} \lambda_q \ell^{-N} \quad (2.30)$$

$$\left\| \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N+\alpha} \quad (2.31)$$

$$\left\| \partial_t \overset{\circ}{\bar{R}}_q + (\bar{v}_q \cdot \nabla) \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} \lesssim \delta_{q+1} \delta_q^{1/2} \lambda_q \ell^{-N-\alpha} \quad (2.32)$$

$$\left| \int_{\mathbb{T}^3} |\bar{v}_q|^2 - |v_\ell|^2 dx \right| \lesssim \delta_{q+1} \ell^\alpha \quad (2.33)$$

where the implicit constants depend only on M, α , and N , cf. Propositions 4.3, 4.4 and 4.5.

The gluing procedure will be broken up into two parts: first, we construct a sequence of exact solutions to the Euler equations with appropriate stability estimates in Section 3 and then we glue the solutions together in Section 4 with a partition of unity in order to construct \bar{v}_q satisfying the properties mentioned above. This is indeed the key idea of Isett in [Ise16]. The main difference with [Ise16] is in the construction of the tensor $\overset{\circ}{\bar{R}}_q$: in this paper we use the usual elliptic operators introduced in [DLS13]. This has the advantage that our Reynolds stress remains trace free, in contrast to the one of [Ise16], and in turn this is crucial to control the energy in the perturbation step below. It should be noticed that in [Ise16] the author resorts to a different definition of $\overset{\circ}{\bar{R}}_q$ because he is not able to find efficient estimates. Our main technical improvement is that this difficulty can be overcome employing suitable commutator estimates on the advective derivative of differential operators of negative order, cf. the proof of Proposition 3.4 and Proposition D.1. This remark allows us not only to keep a better control on the energy and a trace-free Reynolds stress with the desired estimate, but it also shortens the arguments considerably compared to [Ise16].

2.6. Perturbation and proof of Proposition 2.1. The gluing procedure can be used to localize the Reynolds stress error $\overset{\circ}{\bar{R}}_q$ to small disjoint temporal regions, but it cannot be used to completely eliminate the error.

First of all note that as a corollary of (2.10), (2.25) and (2.33), by choosing a sufficiently large we can ensure that

$$\frac{\delta_{q+1}}{2\lambda_q^\alpha} \leq e(t) - \int_{\mathbb{T}^3} |\bar{v}_q|^2 dx \leq 2\delta_{q+1}. \quad (2.34)$$

Starting with the solution $(\bar{v}_q, \bar{p}_q, \overset{\circ}{\bar{R}}_q)$ satisfying (2.28) and the estimates (2.29)-(2.34), we then produce a new solution $(v_{q+1}, p_{q+1}, \overset{\circ}{\bar{R}}_{q+1})$ of the Euler-Reynolds system (2.2) with estimates

$$\|v_{q+1} - \bar{v}_q\|_0 + \lambda_{q+1}^{-1} \|v_{q+1} - \bar{v}_q\|_1 \leq \frac{M}{2} \delta_{q+1}^{1/2} \quad (2.35)$$

$$\|\overset{\circ}{\bar{R}}_{q+1}\|_\alpha \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-4\alpha}}. \quad (2.36)$$

$$\left| e(t) - \int_{\mathbb{T}^3} |v_{q+1}|^2 dx - \frac{\delta_{q+2}}{2} \right| \lesssim \frac{\delta_q^{1/2} \delta_{q+1}^{1/2} \lambda_q^{1+2\alpha}}{\lambda_{q+1}}, \quad (2.37)$$

cf. Corollary 5.8 and Propositions 6.1 and 6.2.

As in previous papers [DLS14, Ise13a, BDLS13, Buc14, BDLISJ15, Buc15, BDLS16] the key idea, introduced in [DLS13], for reducing the size of the error is to add a highly oscillatory perturbation w_{q+1} to \bar{v}_q . Previous schemes heavily relied on Beltrami flows, but these seemed insufficient to push the method beyond Hölder exponent $1/5$. A new set of flows, called Mikado flows, with much better properties were introduced in [DS17] and indeed, a key element in the proof of Isett [Ise16] is the observation, already used in [DS17], that Mikado flows behave better under advection by a mean flow.

An important point is that the Mikado flows will not only be used to “cancel” the error \bar{R}_q , but also to “improve the energy” in areas where the error vanishes identically. In particular, the perturbation will be added in spacetime regions which are disjoint and contained in time-slabs of thickness $2\tau_q$, but with the property that their projections on the time axis is a covering of the interval $[0, T]$.

Proof of Proposition 2.1. The estimate (2.12) is a consequence of (2.22), (2.23), (2.29), (2.30) and (2.35):

$$\|v_{q+1} - v_q\|_0 + \lambda_{q+1}^{-1} \|v_{q+1} - v_1\|_1 \leq \frac{M}{2} \delta_{q+1}^{1/2} + C \delta_{q+1}^{1/2} \ell^\alpha + C \delta_q^{1/2} \lambda_q \lambda_{q+1}^{-1},$$

where the constant C depends on α, β, M , but not on a, b and q . In particular, for every fixed b (2.12) holds if a is large enough. For (2.8), we use the induction assumption to get

$$\|v_{q+1}\|_1 \leq M \delta_q^{1/2} \lambda_q + \frac{M}{2} \delta_{q+1}^{1/2} \lambda_{q+1} + C \delta_{q+1}^{1/2} \ell^\alpha \lambda_{q+1} + C \delta_q^{1/2} \lambda_q$$

and again a sufficiently large choice of a will guarantee $\|v_{q+1}\|_0 \leq M \delta_{q+1}^{1/2} \lambda_{q+1}$. Similarly for (2.9), which will follow from

$$\|v_{q+1}\|_0 \leq \|v_q\|_0 + \|v_{q+1} - v_q\|_0 \leq 1 - \delta_q^{1/2} + M \delta_{q+1}^{1/2}.$$

From (2.36) and (2.37), the inequalities (2.7) and (2.10) follow as a consequence of the parameter inequality

$$\frac{\delta_q^{1/2} \delta_{q+1}^{1/2} \lambda_q}{\lambda_{q+1}} \leq \frac{\delta_{q+2}}{\lambda_{q+1}^{8\alpha}}. \quad (2.38)$$

To see this, one divides by the right hand side, takes logarithms and divides by $\log \lambda_q$, to obtain

$$-\beta - \beta b + 1 - b + 2b^2\beta + 8b\alpha + O\left(\frac{1}{\log \lambda_q}\right) \leq 0,$$

where the error term $O\left(\frac{1}{\log \lambda_q}\right)$ is due to the constants in (2.18). From the relation (2.11), if α is sufficiently small we obtain

$$-\beta - \beta b + 1 - b + 2b^2\beta + 8b\alpha < 0. \quad (2.39)$$

Hence fixing b to satisfy (2.11), choosing subsequently α sufficiently small and then a sufficiently large, we obtain (2.38).

Finally, an entirely analogous argument shows (2.10) from (2.37). \square

3. STABILITY ESTIMATES FOR CLASSICAL EXACT SOLUTIONS

3.1. Classical solutions. For each i , let $t_i = i\tau_q$, and consider smooth solutions of the Euler equations

$$\begin{cases} \partial_t v_i + \operatorname{div}(v_i \otimes v_i) + \nabla p_i = 0 \\ \operatorname{div} v_i = 0 \\ v_i(\cdot, t_i) = v_\ell(\cdot, t_i). \end{cases} \quad (3.1)$$

defined over their own maximal interval of existence. Next, recall the following

Proposition 3.1. *For any $\alpha > 0$ there exists a constant $c = c(\alpha) > 0$ with the following property. Given any initial data $u_0 \in C^\infty$, and $T \leq c \|u_0\|_{1+\alpha}^{-1}$, there exists a unique solution $u : \mathbb{R}^3 \times [-T, T] \rightarrow \mathbb{R}^3$ to the Euler equation*

$$\begin{cases} \partial_t u + \operatorname{div}(u \otimes u) + \nabla p = 0 \\ \operatorname{div} u = 0, \\ u(\cdot, 0) = u_0 \end{cases}$$

Moreover, u obeys the bounds

$$\|u\|_{N+\alpha} \lesssim \|u_0\|_{N+\alpha} . \quad (3.2)$$

for all $N \geq 1$, where the implicit constant depends on N and $\alpha > 0$.

Proof of Proposition 3.1. The proof of the existence of a unique solution is standard (see e.g. [MB02, Chapter 4]), and follows from the restriction $T \leq c \|u_0\|_{1+\alpha}^{-1}$. The higher-order bounds (3.2) are also standard, and can be obtained as follows: For any multi-index θ with $|\theta| = N$ we have

$$\partial_t \partial^\theta v + v \cdot \nabla \partial^\theta v + [\partial^\theta, v \cdot \nabla]v + \nabla \partial^\theta p = 0.$$

Using the equation for the pressure $-\Delta p = \operatorname{tr}(\nabla v \nabla v)$ and Schauder estimates we obtain

$$\|\nabla \partial^\theta p\|_\alpha \lesssim \|\operatorname{tr}(\nabla v \nabla v)\|_{N-1+\alpha} \lesssim \|v\|_{1+\alpha} \|v\|_{N+\alpha}.$$

Therefore

$$\|(\partial_t + v \cdot \nabla) \partial^\theta v\|_\alpha \lesssim \|v\|_{1+\alpha} \|v\|_{N+\alpha},$$

and (3.2) follows by applying (B.3) and Grönwall's inequality. □

An immediate consequence is:

Corollary 3.2. *If a is sufficiently large, for $|t - t_i| \leq \tau_q$, we have*

$$\|v_i\|_{N+\alpha} \lesssim \delta_q^{1/2} \lambda_q \ell^{1-N-\alpha} \lesssim \tau_q^{-1} \ell^{1-N+\alpha} \quad \text{for any } N \geq 1. \quad (3.3)$$

Proof of Corollary 3.2. We apply Proposition 3.1 and use the estimate (2.27) to obtain

$$\|v_i\|_{N+\alpha} \lesssim \|v_\ell(t_i)\|_{N+\alpha}$$

for any $N \geq 1$. From (2.23) we then deduce the estimate (3.3). □

3.2. Stability and estimates on $v_i - v_\ell$. We will now show that for $|t_i - t| \leq \tau_q$, v_i is close to v_ℓ and by the identity

$$v_i - v_{i+1} = (v_i - v_\ell) - (v_{i+1} - v_\ell),$$

the vector field v_i is also close to v_{i+1} .

Proposition 3.3. *For $|t - t_i| \leq \tau_q$ and $N \geq 0$ we have*

$$\|v_i - v_\ell\|_{N+\alpha} \lesssim \tau_q \delta_{q+1} \ell^{-N-1+\alpha}, \quad (3.4)$$

$$\|\nabla(p_\ell - p_i)\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N-1+\alpha}, \quad (3.5)$$

$$\|D_{t,\ell}(v_i - v_\ell)\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N-1+\alpha}, \quad (3.6)$$

where we write

$$D_{t,\ell} = \partial_t + v_\ell \cdot \nabla \quad (3.7)$$

for the transport derivative.

Proof of Proposition 3.3. Let us first consider (3.4) with $N = 0$. From (2.20) and (3.1) we have

$$\partial_t(v_\ell - v_i) + (v_\ell \cdot \nabla)(v_\ell - v_i) = (v_i - v_\ell) \cdot \nabla v_i - \nabla(p_\ell - p_i) + \operatorname{div} \mathring{R}_\ell. \quad (3.8)$$

In particular, using

$$\Delta(p_\ell - p_i) = \operatorname{div}(\nabla v_\ell(v_i - v_\ell)) + \operatorname{div}(\nabla v_i(v_i - v_\ell)) + \operatorname{div} \operatorname{div} \mathring{R}_\ell, \quad (3.9)$$

estimates (2.24) and (3.3), and Proposition C.1 (recall that $\partial_i \partial_j (-\Delta)^{-1}$ is given by $1/3 \delta_{ij}$ + a Calderón-Zygmund operator), we conclude

$$\|\nabla(p_\ell - p_i)(\cdot, t)\|_\alpha \leq \delta_q^{1/2} \lambda_q \ell^{-\alpha} \|v_i - v_\ell\|_\alpha + \delta_{q+1} \ell^{-1+\alpha}.$$

Thus, using (2.24) and the definition of τ_q , we have

$$\|D_{t,\ell}(v_\ell - v_i)\|_\alpha \lesssim \delta_{q+1} \ell^{-1+\alpha} + \tau_q^{-1} \|v_\ell - v_i\|_\alpha \quad (3.10)$$

By applying (B.3) we obtain

$$\|(v_\ell - v_i)(\cdot, t)\|_\alpha \lesssim |t - t_i| \delta_{q+1} \ell^{-1+\alpha} + \int_{t_i}^t \tau_q^{-1} \|(v_\ell - v_i)(\cdot, s)\|_\alpha ds.$$

Applying Grönwall's inequality and using the assumption $|t - t_i| \leq \tau_q$ we obtain

$$\|v_i - v_\ell\|_\alpha \lesssim \tau_q \delta_{q+1} \ell^{-1+\alpha}, \quad (3.11)$$

i.e. (3.4) for the case $N = 0$. Then, as a consequence of (3.10) we obtain (3.6) for the case $N = 0$.

Next, consider the case $N \geq 1$ and let θ be a multiindex with $|\theta| = N$. Commuting the derivative ∂^θ with the material derivative $\partial_t + v_\ell \cdot \nabla$ we have

$$\begin{aligned} \|D_{t,\ell} \partial^\theta (v_\ell - v_i)\|_\alpha &\lesssim \|\partial^\theta D_{t,\ell}(v_\ell - v_i)\|_\alpha + \|[v_\ell \cdot \nabla, \partial^\theta](v_\ell - v_i)\|_\alpha \\ &\lesssim \|\partial^\theta D_{t,\ell}(v_\ell - v_i)\|_\alpha + \|v_\ell\|_{N+\alpha} \|v_\ell - v_i\|_{1+\alpha} + \|v_\ell\|_{1+\alpha} \|v_\ell - v_i\|_{N+\alpha} \\ &\lesssim \|\partial^\theta D_{t,\ell}(v_\ell - v_i)\|_\alpha + \|v_\ell\|_{N+1+\alpha} \|v_\ell - v_i\|_\alpha + \|v_\ell\|_{1+\alpha} \|v_\ell - v_i\|_{N+\alpha}, \end{aligned}$$

where in the last inequality we used the standard interpolation inequalities on Hölder norms, cf. (A.1). On the other hand differentiating (3.8) leads to

$$\begin{aligned} \|\partial^\theta D_{t,\ell}(v_\ell - v_i)\|_\alpha &\lesssim \|v_\ell - v_i\|_{N+\alpha} \|v_i\|_{1+\alpha} + \|v_\ell - v_i\|_\alpha \|v_i\|_{N+1+\alpha} + \|p_\ell - p_i\|_{N+1+\alpha} + \|\mathring{R}_\ell\|_{N+1+\alpha} \\ &\lesssim \tau_q^{-1} \|v_\ell - v_i\|_{N+\alpha} + \delta_{q+1} \ell^{-N-1+\alpha} + \|\nabla(p_\ell - p_i)\|_{N+\alpha}, \end{aligned} \quad (3.12)$$

where we have used (3.11). Furthermore, from (3.9) we also obtain, using Corollary 3.2 and (3.11)

$$\begin{aligned} \|\nabla(p_\ell - p_i)\|_{N+\alpha} &\lesssim (\|v_\ell\|_{N+1+\alpha} + \|v_i\|_{N+1+\alpha})\|v_\ell - v_i\|_\alpha \\ &\quad + (\|v_\ell\|_{1+\alpha} + \|v_i\|_{1+\alpha})\|v_\ell - v_i\|_{N+\alpha} + \|\mathring{R}_\ell\|_{N+1+\alpha} \\ &\lesssim \delta_{q+1}\ell^{-N-1+\alpha} + \tau_q^{-1}\|v_\ell - v_i\|_{N+\alpha}. \end{aligned} \quad (3.13)$$

Summarizing, for any multiindex θ with $|\theta| = N$ we obtain

$$\|D_{t,\ell}\partial^\theta(v_\ell - v_i)\|_\alpha \lesssim \delta_{q+1}\ell^{-N-1+\alpha} + \tau_q^{-1}\|v_\ell - v_i\|_{N+\alpha}.$$

Therefore, invoking once more (B.3) we deduce

$$\|(v_\ell - v_i)(\cdot, t)\|_{N+\alpha} \lesssim \tau_q\delta_{q+1}\ell^{-N-1+\alpha} + \int_{t_i}^t \tau_q^{-1}\|(v_\ell - v_i)(\cdot, s)\|_{N+\alpha} ds,$$

and hence, using Grönwall's inequality and the assumption $|t - t_i| \leq \tau_q$ we obtain (3.4). From (3.13) and (3.12) we then also conclude (3.5) and (3.6). \square

3.3. Estimates on vector potentials. Define the vector potentials to the solutions v_i as

$$z_i = \mathcal{B}v_i := (-\Delta)^{-1} \operatorname{curl} v_i, \quad (3.14)$$

where \mathcal{B} is the Biot-Savart operator, so that

$$\operatorname{div} z_i = 0 \quad \text{and} \quad \operatorname{curl} z_i = v_i - \int_{\mathbb{T}^3} v_i \quad (3.15)$$

(recall that, since $\mathbb{T}^3 = \mathbb{R}^3 \setminus \mathbb{Z}^3$, $\int_{\mathbb{T}^3} v_i(x, t) dx$ is the space average of the solution). Our aim is to obtain estimates for the differences $z_i - z_{i+1}$. The heuristic is as follows: from Proposition 3.3 we obtain

$$\|v_i - v_{i+1}\|_{N+\alpha} \lesssim \tau_q\delta_{q+1}\ell^{-N-1+\alpha}.$$

Since the characteristic length-scale of the vectorfields v_i is ℓ (cf. Corollary 3.2), we expect to gain a factor ℓ when passing to first order potentials. This is formalized in Proposition 3.4 below.

Proposition 3.4. *For $|t - t_i| \leq \tau_q$, we have that*

$$\|z_i - z_{i+1}\|_{N+\alpha} \lesssim \tau_q\delta_{q+1}\ell^{-N+\alpha}, \quad (3.16)$$

$$\|D_{t,\ell}(z_i - z_{i+1})\|_{N+\alpha} \lesssim \delta_{q+1}\ell^{-N+\alpha}, \quad (3.17)$$

where $D_{t,\ell}$ is as in (3.7).

Proof of Proposition 3.4. Set $\tilde{z}_i := \mathcal{B}(v_i - v_\ell)$ and observe that $z_i - z_{i+1} = \tilde{z}_i - \tilde{z}_{i+1}$. Hence, it suffices to estimate \tilde{z}_i in place of $z_i - z_{i+1}$.

The estimate on $\|\nabla\tilde{z}_i\|_{N-1+\alpha}$ for $N \geq 1$ follows directly from (3.4) and the fact that $\nabla\mathcal{B}$ is a bounded operator on Hölder spaces:

$$\|\nabla\tilde{z}_i\|_{N-1+\alpha} \leq \|\nabla\mathcal{B}(v_i - v_\ell)\|_{N-1+\alpha} \|v_i - v_\ell\|_{N+\alpha} \lesssim \tau_q\delta_{q+1}\ell^{-N+\alpha}. \quad (3.18)$$

Next, observe that

$$\partial_t(v_i - v_\ell) + v_\ell \cdot \nabla(v_i - v_\ell) + (v_i - v_\ell) \cdot \nabla v_i + \nabla(p_i - p_\ell) + \operatorname{div} \mathring{R}_\ell = 0. \quad (3.19)$$

Since $v_i - v_\ell = \operatorname{curl} \tilde{z}_i$ with $\operatorname{div} \tilde{z}_i = 0$, we have⁴

$$\begin{aligned} v_\ell \cdot \nabla(v_i - v_\ell) &= \operatorname{curl}((v_\ell \cdot \nabla)\tilde{z}_i) + \operatorname{div}((\tilde{z}_i \times \nabla)v_\ell) \\ ((v_i - v_\ell) \cdot \nabla)v_i &= \operatorname{div}((\tilde{z}_i \times \nabla)v_i^T), \end{aligned}$$

⁴Here we use the notation $[(z \times \nabla)v]^{ij} = \epsilon_{ikl}z^k\partial_l v^j$ for vector fields z, v .

so that we can write (3.19) as

$$\operatorname{curl}(\partial_t \tilde{z}_i + (v_\ell \cdot \nabla) \tilde{z}_i) = -\operatorname{div}((\tilde{z}_i \times \nabla) v_\ell + (\tilde{z}_i \times \nabla) v_i^T) - \nabla(p_i - p_\ell) - \operatorname{div} \mathring{R}_\ell. \quad (3.20)$$

Taking the curl of (3.20) the pressure term drops out. Using in addition that $\operatorname{div} \tilde{z}_i = \operatorname{div}(v_i - v_\ell) = 0$ and the identity $\operatorname{curl} \operatorname{curl} = -\Delta + \nabla \operatorname{div}$, we then arrive at

$$-\Delta(\partial_t \tilde{z}_i + (v_\ell \cdot \nabla) \tilde{z}_i) = F,$$

where

$$F = -\nabla \operatorname{div}((\tilde{z}_i \cdot \nabla) v_\ell) - \operatorname{curl} \operatorname{div}((\tilde{z}_i \times \nabla) v_\ell + (\tilde{z}_i \times \nabla) v_i^T) - \operatorname{curl} \operatorname{div} \mathring{R}_\ell.^5$$

Consequently,

$$\begin{aligned} \|\partial_t \tilde{z}_i + (v_\ell \cdot \nabla) \tilde{z}_i\|_{N+\alpha} &\lesssim (\|v_i\|_{N+1+\alpha} + \|v_\ell\|_{N+1+\alpha}) \|\tilde{z}_i\|_\alpha \\ &\quad + (\|v_i\|_{1+\alpha} + \|v_\ell\|_{1+\alpha}) \|\tilde{z}_i\|_{N+\alpha} + \|\mathring{R}_\ell\|_{N+\alpha} \\ &\lesssim \tau_q^{-1} \|\tilde{z}_i\|_{N+\alpha} + \tau_q^{-1} \ell^{-N} \|\tilde{z}_i\|_\alpha + \delta_{q+1} \ell^{-N+\alpha}. \end{aligned} \quad (3.21)$$

Setting $N = 0$ and using (B.3) and Grönwall's inequality we obtain

$$\|\tilde{z}_i\|_\alpha \lesssim \tau_q \delta_{q+1} \ell^\alpha,$$

which together with (3.18) gives (3.16). Using (3.16) into (3.21) we conclude

$$\|\partial_t \tilde{z}_i + (v_\ell \cdot \nabla) \tilde{z}_i\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N+\alpha}.$$

Finally commuting the derivatives in the $N + \alpha$ -norm with $D_{t,\ell}$ as in the proof of Proposition 3.3 and using again (3.16) we achieve (3.17). \square

4. GLUING PROCEDURE

Now we proceed to glue the solutions v_i together in order to construct \bar{v}_q . The stability estimates above will be used in order to ensure that \bar{v}_q remains an approximate solution to the Euler equations.

4.1. Partition of unity and definition of \bar{v}_q . Let

$$t_i = i\tau_q, \quad I_i = [t_i + \frac{1}{3}\tau_q, t_i + \frac{2}{3}\tau_q] \cap [0, T], \quad J_i = (t_i - \frac{1}{3}\tau_q, t_i + \frac{1}{3}\tau_q) \cap [0, T].$$

Note that $\{I_i, J_i\}_i$ is a decomposition of $[0, T]$ into pairwise disjoint intervals. We define a partition of unity $\{\chi_i\}_i$ in time with the following properties:

- The cut-offs form a partition of unity

$$\sum_i \chi_i \equiv 1 \quad (4.1)$$

- $\operatorname{supp} \chi_i \cap \operatorname{supp} \chi_{i+2} = \emptyset$ and moreover

$$\begin{aligned} \operatorname{supp} \chi_i &\subset (t_i - \frac{2}{3}\tau_q, t_i + \frac{2}{3}\tau_q) \\ \chi_i(t) &= 1 \quad \text{for } t \in J_i \end{aligned} \quad (4.2)$$

- For any i and N we have

$$\|\partial_t^N \chi_i\|_0 \lesssim \tau_q^{-N}. \quad (4.3)$$

⁵In deriving the latter equality we have used the identity $\nabla \operatorname{div}((v_\ell \cdot \nabla) \tilde{z}_i) = \nabla \operatorname{div}((\tilde{z}_i \cdot \nabla) v_\ell)$, which follows easily from the fact that both v_ℓ and \tilde{z}_i are divergence free.

We define

$$\begin{aligned}\bar{v}_q &= \sum_i \chi_i v_i \\ \bar{p}_q^{(1)} &= \sum_i \chi_i p_i\end{aligned}$$

Observe that $\operatorname{div} \bar{v}_q = 0$. Furthermore, if $t \in I_i$, then $\chi_i + \chi_{i+1} = 1$ and $\chi_j = 0$ for $j \neq i, i+1$, therefore on I_i :

$$\begin{aligned}\bar{v}_q &= \chi_i v_i + (1 - \chi_i) v_{i+1} \\ \bar{p}_q^{(1)} &= \chi_i p_i + (1 - \chi_i) p_{i+1}\end{aligned}$$

and

$$\begin{aligned}\partial_t \bar{v}_q + \operatorname{div}(\bar{v}_q \otimes \bar{v}_q) + \nabla \bar{p}_q^{(1)} &= \chi_i \partial_t v_i + (1 - \chi_i) \partial_t v_{i+1} + \partial_t \chi_i (v_i - v_{i+1}) \\ &\quad + \operatorname{div}(\chi_i^2 v_i \otimes v_i + (1 - \chi_i)^2 v_{i+1} \otimes v_{i+1}) \\ &\quad + \chi_i (1 - \chi_i) \operatorname{div}(v_i \otimes v_{i+1} + v_{i+1} \otimes v_i) \\ &\quad + \chi_i \nabla p_i + (1 - \chi_i) \nabla p_{i+1} \\ &= \partial_t \chi_i (v_i - v_{i+1}) - \chi_i (1 - \chi_i) \operatorname{div}((v_i - v_{i+1}) \otimes (v_i - v_{i+1})).\end{aligned}$$

On the other hand, if $t \in J_i$ then $\chi_i = 1$ and $\chi_j = 0$ for all $j \neq i$ for all \tilde{t} sufficiently close to t (since J_i is open). Then for all $t \in J_i$ we have

$$\bar{v}_q = v_i, \quad \bar{p}_q^{(1)} = p_i,$$

and, from (3.1),

$$\partial_t \bar{v}_q + \operatorname{div}(\bar{v}_q \otimes \bar{v}_q) + \nabla \bar{p}_q^{(1)} = 0.$$

4.2. The new Reynolds tensor. In order to define the new Reynolds tensor, we recall the operator \mathcal{R} from [DLS13], which can be thought of as an ‘‘inverse divergence’’ operator for symmetric tracefree 2-tensors. The operator is defined as

$$\begin{aligned}(\mathcal{R}f)^{ij} &= \mathcal{R}^{ijk} f^k \\ \mathcal{R}^{ijk} &= -\frac{1}{2} \Delta^{-2} \partial_i \partial_j \partial_k + \frac{1}{2} \Delta^{-1} \partial_k \delta_{ij} - \Delta^{-1} \partial_i \delta_{jk} - \Delta^{-1} \partial_j \delta_{ik}.\end{aligned}\tag{4.4}$$

when acting on vectors f with zero mean on \mathbb{T}^3 . The following statement, taken from [DLS13], can be proved by direct calculation.

Proposition 4.1. *The tensor \mathcal{R} defined in (4.4) is symmetric, and we have*

$$\operatorname{div}(\mathcal{R}f) = f$$

for any f with zero mean on \mathbb{T}^3 .

We define

$$\begin{aligned}\overset{\circ}{R}_q &= \partial_t \chi_i \mathcal{R}(v_i - v_{i+1}) - \chi_i (1 - \chi_i) (v_i - v_{i+1}) \overset{\circ}{\otimes} (v_i - v_{i+1}) \\ \bar{p}_q^{(2)} &= -\chi_i (1 - \chi_i) |v_i - v_{i+1}|^2,\end{aligned}$$

for $t \in I_i$ and $\overset{\circ}{R}_q = 0, \bar{p}_q^{(2)} = 0$ for $t \notin \bigcup_i I_i$.

Remark 4.2. We must however check that $v_i - v_{i+1}$ has zero mean. Recall that v_i solves Euler, so its mean is constant:

$$\frac{d}{dt} \int_{\mathbb{T}^3} v_i = - \int_{\mathbb{T}^3} (\operatorname{div}(v_i \otimes v_i) - \nabla p_i) = 0.$$

For the same reason, the mean of v_ℓ is also constant. Since $v_i - v_\ell$ coincide at the time t_i , we conclude that $v_i - v_\ell$ has zero mean at every time. But then the same applies to $v_{i+1} - v_\ell$ and hence to $v_{i+1} - v_i = (v_{i+1} - v_\ell) - (v_i - v_\ell)$.

Furthermore, we set

$$\bar{p}_q = \bar{p}_q^{(1)} + \bar{p}_q^{(2)}$$

It follows from the preceding discussion and Proposition 4.1 that

- $\overset{\circ}{R}_q$ is a smooth symmetric and traceless 2-tensor;
- For all $(x, t) \in \mathbb{T}^3 \times [0, T]$

$$\begin{cases} \partial_t \bar{v}_q + \operatorname{div}(\bar{v}_q \otimes \bar{v}_q) + \nabla \bar{p}_q = \operatorname{div} \overset{\circ}{R}_q, \\ \operatorname{div} \bar{v}_q = 0; \end{cases}$$

- $\operatorname{supp} \overset{\circ}{R}_q \subset \mathbb{T}^3 \times \bigcup_i I_i$.

4.3. Estimates on \bar{v}_q . Next, we estimate the various Hölder norms of \bar{v}_q and $\overset{\circ}{R}_q$ in order to obtain (2.29)–(2.32).

Proposition 4.3. *The velocity field \bar{v}_q satisfies the following estimates*

$$\|\bar{v}_q - v_\ell\|_\alpha \lesssim \delta_{q+1}^{1/2} \ell^\alpha \tag{4.5}$$

$$\|\bar{v}_q - v_\ell\|_{N+\alpha} \lesssim \tau_q \delta_{q+1} \ell^{-1-N+\alpha} \tag{4.6}$$

$$\|\bar{v}_q\|_{1+N} \lesssim \delta_q^{1/2} \lambda_q \ell^{-N} \tag{4.7}$$

for all $N \geq 0$.

In particular, this lemma shows that the claimed estimates (2.29)–(2.30) indeed hold.

Proof of Proposition 4.3. By definition

$$\bar{v}_q - v_\ell = \sum_i \chi_i (v_i - v_\ell).$$

Therefore Proposition 3.3 implies

$$\|\bar{v}_q - v_\ell\|_{N+\alpha} \lesssim \tau_q \delta_{q+1} \ell^{-1-N+\alpha}. \tag{4.8}$$

Note that using the definition of ℓ in (2.19) and τ_q in (2.26) and the comparison (2.21)

$$\delta_{q+1}^{1/2} \tau_q \ell^{-1} = \ell^{2\alpha} \lambda_q^{3\alpha/2} \leq \lambda_q^{-\alpha/2} \leq 1. \tag{4.9}$$

Therefore we obtain (4.5), and furthermore, for any $N \geq 0$

$$\|\bar{v}_q - v_\ell\|_{1+N+\alpha} \lesssim \delta_{q+1} \tau_q \ell^{-N-2+\alpha} = \delta_q^{1/2} \lambda_q (\ell \lambda_q)^{3\alpha} \ell^{-N} \leq \delta_q^{1/2} \lambda_q \ell^{-N}.$$

Then it also follows using (2.23) that

$$\|\bar{v}_q\|_{1+N} \lesssim \|v_\ell\|_{1+N} + \|v_\ell - \bar{v}_q\|_{1+N+\alpha} \lesssim \delta_q^{1/2} \lambda_q \ell^{-N}. \quad \square$$

4.4. **Estimates on the stress tensor.** We are now in a position to estimate the glued stress tensor $\overset{\circ}{\bar{R}}_q$:

Proposition 4.4. *The stress tensor $\overset{\circ}{\bar{R}}_q$ satisfies the following bounds for any $N \geq 0$:*

$$\left\| \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} \lesssim \delta_{q+1} \ell^{-N+\alpha} \quad (4.10)$$

$$\left\| (\partial_t + \bar{v}_q \cdot \nabla) \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} \lesssim \delta_{q+1} \delta_q^{1/2} \lambda_q \ell^{-N-\alpha}. \quad (4.11)$$

This shows that the claimed estimates (2.31)–(2.32) are indeed obeyed by $\overset{\circ}{\bar{R}}_q$.

Proof of Proposition 4.4. Recall that $v_i - \int_{\mathbb{T}^3} v_i = \text{curl } z_i$. Since $v_{i+1} - v_i$ has zero mean (cf. Remark 4.2, for $t \in I_i$ we have $\text{curl}(z_{i+1} z_i) = v_{i+1} - v_i$ and we may write:

$$\overset{\circ}{\bar{R}}_q = \partial_t \chi_i (\mathcal{R} \text{curl})(z_i - z_{i+1}) - \chi_i (1 - \chi_i) (v_i - v_{i+1}) \overset{\circ}{\otimes} (v_i - v_{i+1}).$$

Note that $\mathcal{R} \text{curl}$ is a zero-order operator. Therefore we obtain from Propositions 3.3 and 3.4 for any $N \geq 0$ with $t \in I_i$

$$\begin{aligned} \left\| \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} &\lesssim \tau_q^{-1} \|z_i - z_{i+1}\|_{N+\alpha} + \|v_i - v_{i+1}\|_{N+\alpha} \|v_i - v_{i+1}\|_\alpha \\ &\lesssim \delta_{q+1} \ell^{-N+\alpha} + \tau_q^2 \delta_{q+1}^2 \ell^{-2-N+2\alpha} \lesssim \delta_{q+1} \ell^{-N+\alpha}. \end{aligned}$$

Here we used again (4.9). Next, we calculate

$$\begin{aligned} D_{t,\ell} \overset{\circ}{\bar{R}}_q &= \partial_t^2 \chi_i (\mathcal{R} \text{curl})(z_i - z_{i+1}) \\ &\quad + \partial_t \chi_i (\mathcal{R} \text{curl}) D_{t,\ell} (z_i - z_{i+1}) + \partial_t \chi_i [v_\ell \cdot \nabla, \mathcal{R} \text{curl}](z_i - z_{i+1}) \\ &\quad - \partial_t (\chi_i (1 - \chi_i)) (v_i - v_{i+1}) \overset{\circ}{\otimes} (v_i - v_{i+1}) \\ &\quad - \chi_i (1 - \chi_i) \left((D_{t,\ell} (v_i - v_{i+1})) \overset{\circ}{\otimes} (v_i - v_{i+1}) - (v_i - v_{i+1}) \overset{\circ}{\otimes} (D_{t,\ell} (v_i - v_{i+1})) \right), \end{aligned}$$

where $[v_\ell \cdot \nabla, \mathcal{R} \text{curl}]$ denotes the commutator. Hence, using Proposition D.1 and Propositions 3.3 and 3.4 we deduce

$$\begin{aligned} \left\| D_{t,\ell} \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} &\lesssim \tau_q^{-2} \|z_i - z_{i+1}\|_{N+\alpha} + \tau_q^{-1} \|D_{t,\ell} (z_i - z_{i+1})\|_{N+\alpha} \\ &\quad + \tau_q^{-1} \|v_\ell\|_\alpha \|z_i - z_{i+1}\|_{N+\alpha} + \tau_q^{-1} \|v_\ell\|_{N+\alpha} \|z_i - z_{i+1}\|_\alpha \\ &\quad + \tau_q^{-1} \|v_i - v_{i+1}\|_{N+\alpha} \|v_i - v_{i+1}\|_\alpha \\ &\quad + \|D_{t,\ell} (v_i - v_{i+1})\|_{N+\alpha} \|v_i - v_{i+1}\|_\alpha + \|v_i - v_{i+1}\|_{N+\alpha} \|D_{t,\ell} (v_i - v_{i+1})\|_\alpha \\ &\lesssim \tau_q^{-1} \delta_{q+1} \ell^{-N+\alpha} + (\tau_q^2 \delta_{q+1} \ell^{-2}) \tau_q^{-1} \delta_{q+1} \ell^{-N+2\alpha} \\ &\lesssim \tau_q^{-1} \delta_{q+1} \ell^{-N+\alpha}. \end{aligned}$$

Finally, we deduce using (4.6):

$$\begin{aligned} \left\| (\partial_t + \bar{v}_q \cdot \nabla) \overset{\circ}{\bar{R}}_q \right\|_{N+\alpha} &\lesssim \|(v_\ell - \bar{v}_q) \cdot \nabla \overset{\circ}{\bar{R}}_q\|_{N+\alpha} + \|D_{t,\ell} \overset{\circ}{\bar{R}}_q\|_{N+\alpha} \\ &\lesssim \|v_\ell - \bar{v}_q\|_{N+\alpha} \|\overset{\circ}{\bar{R}}_q\|_{1+\alpha} + \|v_\ell - \bar{v}_q\|_\alpha \|\overset{\circ}{\bar{R}}_q\|_{N+1+\alpha} + \|D_{t,\ell} \overset{\circ}{\bar{R}}_q\|_{N+\alpha} \\ &\lesssim \tau_q \delta_{q+1}^2 \ell^{-N-2+2\alpha} + \tau_q^{-1} \delta_{q+1} \ell^{-N+\alpha} \\ &\lesssim \tau_q^{-1} \delta_{q+1} \ell^{-N+\alpha} = \delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q \ell^{-N-\alpha} \end{aligned}$$

again using (4.9). □

To finish this section we show that \bar{v}_q has approximately the same energy as v_ℓ :

Proposition 4.5. *The difference of the energies of \bar{v}_q and v_ℓ satisfies*

$$\left| \int_{\mathbb{T}^3} |\bar{v}_q|^2 - |v_\ell|^2 dx \right| \lesssim \delta_{q+1} \ell^\alpha \quad (4.12)$$

Proof of Proposition 4.5. Observe that for $t \in I_i$

$$\begin{aligned} \bar{v}_q \otimes \bar{v}_q &= (\chi_i v_i + (1 - \chi_i) v_{i+1}) \otimes (\chi_i v_i + (1 - \chi_i) v_{i+1}) \\ &= \chi_i v_i \otimes v_i + (1 - \chi_i) v_{i+1} \otimes v_{i+1} - \chi_i (1 - \chi_i) (v_i - v_{i+1}) \otimes (v_i - v_{i+1}), \end{aligned}$$

so that, taking the trace:

$$|\bar{v}_q|^2 - |v_\ell|^2 = \chi_i (|v_i|^2 - |v_\ell|^2) + (1 - \chi_i) (|v_{i+1}|^2 - |v_\ell|^2) - \chi_i (1 - \chi_i) |v_i - v_{i+1}|^2$$

Next, recall that v_i and v_ℓ are smooth solutions of (3.1) and (2.20) respectively, therefore

$$\begin{aligned} \left| \frac{d}{dt} \int_{\mathbb{T}^3} |v_i|^2 - |v_\ell|^2 dx \right| &= 2 \left| \int_{\mathbb{T}^3} \nabla v_\ell : \dot{R}_\ell dx \right| \lesssim \|\nabla v_\ell\|_0 \|\dot{R}_\ell\|_0 \\ &\lesssim \delta_q^{1/2} \lambda_q \delta_{q+1} \lesssim \tau_q^{-1} \delta_{q+1} \ell^\alpha, \end{aligned}$$

where we have used (2.24) and (3.3). Moreover, $v_i = v_\ell$ for $t = t_i$. Therefore, after integrating in time we deduce

$$\left| \int_{\mathbb{T}^3} |v_i|^2 - |v_\ell|^2 dx \right| \lesssim \delta_{q+1} \ell^\alpha.$$

Furthermore, using (3.4) and (4.9)

$$\int_{\mathbb{T}^3} |v_i - v_{i+1}|^2 dx \lesssim \|v_i - v_{i+1}\|_\alpha^2 \lesssim \tau_q^2 \delta_{q+1}^2 \ell^{-2+2\alpha} \stackrel{(4.9)}{\lesssim} \delta_{q+1} \ell^{2\alpha},$$

Therefore

$$\left| \int |\bar{v}_q|^2 - |v_\ell|^2 dx \right| \lesssim \delta_{q+1} \ell^\alpha,$$

which concludes the proof. \square

5. PERTURBATION STEP

In this section, we will outline the construction of the perturbation w_{q+1} , where

$$v_{q+1} := w_{q+1} + \bar{v}_q,$$

As already explained in the outline of the proof, the perturbation w_{q+1} is highly oscillatory and will be based on the Mikado flows introduced in [DS17], which are designed to cancel the low frequency error \bar{R}_q and are Lie-advected by the mean flow of \bar{v}_q .

5.1. Mikado flows. We begin by recalling the construction of Mikado flows given in [DS17].

Lemma 5.1. *For any compact subset $\mathcal{N} \subset\subset \mathcal{S}_+^{3 \times 3}$ there exists a smooth vector field*

$$W : \mathcal{N} \times \mathbb{T}^3 \rightarrow \mathbb{R}^3,$$

such that, for every $R \in \mathcal{N}$

$$\begin{cases} \operatorname{div}_\xi (W(R, \xi) \otimes W(R, \xi)) = 0 \\ \operatorname{div}_\xi W(R, \xi) = 0, \end{cases} \quad (5.1)$$

and

$$\oint_{\mathbb{T}^3} W(R, \xi) d\xi = 0, \quad (5.2)$$

$$\oint_{\mathbb{T}^3} W(R, \xi) \otimes W(R, \xi) d\xi = R. \quad (5.3)$$

Using the fact that $W(R, \xi)$ is \mathbb{T}^3 -periodic and has zero mean in ξ , we write

$$W(R, \xi) = \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} a_k(R) e^{ik \cdot \xi} \quad (5.4)$$

for some smooth functions $R \rightarrow a_k(R) \in \mathbb{C}^3$, satisfying $a_k(R) \cdot k = 0$. From the smoothness of W , we further infer

$$\sup_{R \in \mathcal{N}} |D_R^N a_k(R)| \leq \frac{C(\mathcal{N}, N, m)}{|k|^m} \quad (5.5)$$

for some constant C , which depends, as highlighted in the statement, on \mathcal{N} , N and m .

Remark 5.2. *Later in the proof the estimates (5.5) will be used with a specific choice of the compact set \mathcal{N} and of the integers N and m : this specific choice will then determine the universal constant M appearing in Proposition 2.1.*

Using the Fourier representation we see that from (5.3)

$$W(R, \xi) \otimes W(R, \xi) = R + \sum_{k \neq 0} C_k(R) e^{ik \cdot \xi} \quad (5.6)$$

where

$$C_k k = 0 \quad \text{and} \quad \sup_{R \in \mathcal{N}} |D_R^N C_k(R)| \leq \frac{C(\mathcal{N}, N, m)}{|k|^m} \quad (5.7)$$

for any $m, N \in \mathbb{N}$.

It will also be useful to write the Mikado flows in terms of a potential. We note

$$\operatorname{curl}_\xi \left(\left(\frac{ik \times a_k}{|k|^2} \right) e^{ik \cdot \xi} \right) = -i \left(\frac{ik \times a_k}{|k|^2} \right) \times k e^{ik \cdot \xi} = -\frac{k \times (k \times a_k)}{|k|^2} e^{ik \cdot \xi} = a_k e^{ik \cdot \xi} \quad (5.8)$$

5.2. Squiggling stripes and the stress tensor $\tilde{R}_{q,i}$. Recall that \tilde{R}_q is supported in the set $\mathbb{T}^3 \times \bigcup_i I_i$, whereas, from (4.2) it follows that $[0, T] \setminus \bigcup_i I_i = \bigcup_i J_i$, where the open intervals J_i have length $|J_i| = \frac{2}{3}\tau_q$ each, except for the first and last one, which might be shortened by the intersection with $[0, T]$, more precisely

$$J_i = (t_i - \frac{1}{3}\tau_q, t_i + \frac{1}{3}\tau_q) \cap [0, T].$$

We start by defining smooth non-negative cut-off functions $\eta_i = \eta_i(x, t)$ with the following properties

- (i) $\eta_i \in C^\infty(\mathbb{T}^3 \times [0, T])$ with $0 \leq \eta_i(x, t) \leq 1$ for all (x, t) ;
- (ii) $\operatorname{supp} \eta_i \cap \operatorname{supp} \eta_j = \emptyset$ for $i \neq j$;
- (iii) $\mathbb{T}^3 \times I_i \subset \{(x, t) : \eta_i(x, t) = 1\}$;
- (iv) $\operatorname{supp} \eta_i \subset \mathbb{T}^3 \times I_i \cup J_i \cup J_{i+1} = \mathbb{T}^3 \times (t_i - \frac{1}{3}\tau_q, t_{i+1} + \frac{1}{3}\tau_q) \cap [0, T]$;




FIGURE 1. The support of $\overset{\circ}{R}_q$ is given by the blue regions. The support of the cut-off functions η_i are encapsulated in the red region.

(v) There exists a positive geometric constant $c_0 > 0$ such that for any $t \in [0, T]$

$$\sum_i \int_{\mathbb{T}^3} \eta_i^2(x, t) dx \geq c_0.$$

In view of (iv) we set

$$\tilde{I}_i = (t_i - \frac{1}{3}\tau_q, t_{i+1} + \frac{1}{3}\tau_q) \cap [0, T].$$

Lemma 5.3. *There exists cut-off functions $\{\eta_i\}_i$ with the properties (i)-(v) above and such that for any i and $n, m \geq 0$*

$$\|\partial_t^n \eta_i\|_m \leq C(n, m) \tau_q^{-n}$$

where $C(n, m)$ are geometric constants depending only upon m and n .

Proof of Lemma 5.3. First of all we consider the sharp cutoffs $\tilde{\eta}_i$ defined by

$$\begin{aligned} \tilde{\eta}_i &= \mathbf{1}_{\tilde{\Omega}_i} \\ \tilde{\Omega}_i &= \left\{ (x, t) : t_i + \frac{\tau_q}{6} (\sin(2\pi x_1) + \frac{1}{2}) \leq t \leq t_{i+1} + \frac{\tau_q}{6} (\sin(2\pi x_1) - \frac{1}{2}) \right\} \end{aligned}$$

Next we fix a standard mollifier \varkappa in time and the standard mollifier ψ in space already used so far. Hence we define η_i by mollifying $\tilde{\eta}_i$ in space and time as follows:

$$\eta_i(x, t) = \int \tilde{\eta}_i(y, s) \psi\left(\frac{x-y}{c_1}\right) \varkappa\left(\frac{t-s}{c_2\tau_q}\right) dy ds,$$

where c_1 and c_2 are positive geometric constants. One may check that a suitable choice of c_1 and c_2 yields the desired conclusions (see Figure 1). \square

Define

$$\rho_q(t) := \frac{1}{3} \left(e(t) - \frac{\delta_{q+2}}{2} - \int_{\mathbb{T}^3} |\bar{v}_q|^2 dx \right)$$

and

$$\rho_{q,i}(x, t) := \frac{\eta_i^2(x, t)}{\sum_j \int_{\mathbb{T}^3} \eta_j^2(y, t) dy} \rho_q(t)$$

Define the backward flows Φ_i for the velocity field \bar{v}_q as the solution of the transport equation

$$\begin{cases} (\partial_t + \bar{v}_q \cdot \nabla)\Phi_i = 0 \\ \Phi_i(x, t_i) = x. \end{cases}$$

Define

$$R_{q,i} := \rho_{q,i} \text{Id} - \eta_i^2 \overset{\circ}{R}_q$$

and

$$\tilde{R}_{q,i} = \frac{\nabla \Phi_i R_{q,i} (\nabla \Phi_i)^T}{\rho_{q,i}}. \quad (5.9)$$

We note that, because of properties (ii)-(iv) of η_i ,

- $\text{supp } R_{q,i} \subset \text{supp } \eta_i$ and on $\text{supp } \eta_i$ we have $R_{q,i} = \rho_{q,i} \text{Id} - \overset{\circ}{R}_q$;
- $\text{supp } \tilde{R}_{q,i} \subset \mathbb{T}^3 \times (t_i - \frac{1}{3}\tau_q, t_{i+1} + \frac{1}{3}\tau_q)$;
- $\text{supp } \tilde{R}_{q,i} \cap \text{supp } \tilde{R}_{q,j} = \emptyset$ for all $i \neq j$.

Lemma 5.4. *For $a \gg 1$ sufficiently large we have*

$$\|\nabla \Phi_i - \text{Id}\|_0 \leq \frac{1}{2} \quad \text{for } t \in \text{supp}(\eta_i). \quad (5.10)$$

Furthermore, for any $N \geq 0$

$$\frac{\delta_{q+1}}{8\lambda_q^\alpha} \leq |\rho_q(t)| \leq \delta_{q+1} \quad \text{for all } t, \quad (5.11)$$

$$\|\rho_{q,i}\|_0 \leq \frac{\delta_{q+1}}{c_0}, \quad (5.12)$$

$$\|\rho_{q,i}\|_N \lesssim \delta_{q+1}, \quad (5.13)$$

$$\|\partial_t \rho_q\|_0 \lesssim \delta_{q+1} \delta_q^{1/2} \lambda_q, \quad (5.14)$$

$$\|\partial_t \rho_{q,i}\|_N \lesssim \delta_{q+1} \tau_q^{-1}. \quad (5.15)$$

Moreover, for all (x, t)

$$\tilde{R}_{q,i}(x, t) \in B_{1/2}(\text{Id}) \subset \mathcal{S}_+^{3 \times 3},$$

where $B_{1/2}(\text{Id})$ denotes the metric ball of radius 1/2 around the identity Id in the space $\mathcal{S}^{3 \times 3}$.

Proof of Lemma 5.4. Note that (5.11) is a trivial consequence of estimate (2.34) and the inequality $4\delta_{q+2} \leq \delta_{q+1}$. Note that by the definition of the cut-off functions η_i

$$c_0 \leq \sum_i \int_{\mathbb{T}^3} \eta_i^2(y, t) dy \leq 2 \quad (5.16)$$

and hence we obtain (5.12). Since $|\nabla^N \eta_j| \lesssim 1$, the bound (5.13) also follows.

Next, note that by applying (2.30) and (B.5) we obtain

$$\|\nabla \Phi_i - \text{Id}\|_0 \lesssim \tau_q \delta_q^{1/2} \lambda_q = \ell^{2\alpha}.$$

Furthermore, by definition we have

$$\begin{aligned} \tilde{R}_{q,i} - \text{Id} &= \nabla \Phi_i \left(\frac{R_{q,i}}{\rho_{q,i}} - \text{Id} \right) \nabla \Phi_i^T + \nabla \Phi_i \nabla \Phi_i^T - \text{Id} \\ &= -\nabla \Phi_i \frac{\eta_i^2 \overset{\circ}{R}_q}{\rho_{q,i}} \nabla \Phi_i^T + \nabla \Phi_i \nabla \Phi_i^T - \text{Id} \end{aligned}$$

Using (2.31) we see that

$$\left| \frac{\eta_i^2 \overset{\circ}{R}_q}{\rho_{q,i}} \right| \lesssim \frac{1}{\delta_{q+1}} \left| \overset{\circ}{R}_q \right| \lesssim \ell^\alpha.$$

Consequently we obtain

$$|\tilde{R}_{q,i} - \text{Id}| \lesssim \ell^\alpha$$

so that, choosing a sufficiently large, we ensure that $\tilde{R}_{q,i}(x, t)$ is contained in the ball of symmetric matrices $B_{1/2}(\text{Id})$.

Finally, to prove (5.15) we first note that

$$\left| \frac{d}{dt} \int |\bar{v}_q(x, t)|^2 dx \right| = \left| 2 \int \nabla \bar{v}_q \cdot \overset{\circ}{R}_q dx \right| \lesssim \delta_{q+1} \delta_q^{1/2} \lambda_q$$

Thus

$$\|\partial_t \rho_q\|_0 \lesssim \delta_{q+1} \delta_q^{1/2} \lambda^6$$

Then, since $\|\partial_t \eta_j\|_N \lesssim \tau_q^{-1}$ and $\delta_q^{1/2} \lambda_q \leq \tau_q^{-1}$, using (5.16), the estimate (5.15) follows. \square

5.3. The perturbation and the constant M . The principal term of the perturbation can be written as

$$w_o := \sum_i (\rho_{q,i}(x, t))^{1/2} (\nabla \Phi_i)^{-1} W(\tilde{R}_{q,i}, \lambda_{q+1} \Phi_i) = \sum_i w_{o,i}, \quad (5.17)$$

where Lemma 5.1 is applied with $\mathcal{N} = \bar{B}_{1/2}(\text{Id})$, namely the closed ball (in the space of symmetric 3×3 matrices) of radius $1/2$ centered at the identity matrix.

From Lemma 5.4 it follows that $W(\tilde{R}_{q,i}, \lambda_{q+1} \Phi_i)$ is well defined. Using the Fourier series representation of the Mikado flows (5.4) we obtain

$$w_{o,i} := \sum_{k \neq 0} (\rho_{q,i}(x, t))^{1/2} (\nabla \Phi_i)^{-1} a_k(\tilde{R}_{q,i}) e^{i\lambda_{q+1} k \cdot \Phi_i}.$$

The choice of w_o is motivated by the fact that the vector fields

$$U_{i,k} := (\nabla \Phi_i)^{-1} a_k(\tilde{R}_{q,i}) e^{i\lambda_{q+1} k \cdot \Phi_i}$$

are Lie-advected by the flow up to a ‘‘lower order term’’ in λ_{q+1} :

$$(\partial_t + \bar{v}_q \cdot \nabla) U_{i,k} = D\bar{v}_q U_{i,k} + (\nabla \Phi_i)^{-1} [(\partial_t + \bar{v}_q \cdot \nabla) a_k(\tilde{R}_{q,i})] e^{i\lambda_{q+1} k \cdot \Phi_i}. \quad (5.18)$$

Hence (again up to lower order term in λ_{q+1}), the vector fields $U_{i,k}$ remain divergence free. For notational convenience we set

$$b_{i,k}(x, t) := (\rho_{q,i}(x, t))^{1/2} a_k(\tilde{R}_{q,i}(x, t)) A_k$$

so that we may write

$$w_{o,i} := \sum_{k \neq 0} (\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i}.$$

The following is a crucial point of our construction, which ensures that the constant M of Proposition 2.1 is geometric and in particular independent of all the parameters of the construction.

Lemma 5.5. *There is a geometric constant \bar{M} such that*

$$\|b_{i,k}\|_0 \leq \frac{\bar{M}}{|k|^4} \delta_{q+1}^{1/2}. \quad (5.19)$$

⁶Note that $\|\partial_t e\|_0 \leq C \leq \delta_{q+1} \delta_q^{1/2} \lambda_q$ since $\delta_{q+1} \delta_q^{1/2} \lambda_q = \lambda_{q+1}^{-2\beta} \lambda_q^{1-\beta} \geq C a^{bq} (1 - \beta - 2\beta b) \geq C$ (recall that $b < \frac{1-\beta}{2\beta}$).

Proof of Lemma 5.5. First of all, applying (5.5) with $N = 0, m = 4$ and $\mathcal{N} = \bar{B}_{1/2}(\text{Id})$, we achieve

$$\|a_k(\tilde{R}_{q,i})\|_0 \leq \frac{\bar{C}}{|k|^4},$$

where \bar{C} is a geometric constant (cf. Remark 5.2). Hence, considering the bound (5.12), the constant \bar{M} is given by $\bar{C}c_0^{-1/2}$. \square

We are finally ready to define the constant M of Proposition 2.1: from Lemma 5.5 it follows trivially that the constant is indeed geometric and hence independent of all the parameters entering in the statement of Proposition 2.1.

Definition 5.6. *The constant M is defined as*

$$M = 64\bar{M} \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4},$$

where \bar{M} is the constant of Lemma 5.5.

In order to ensure w_{q+1} is divergence free, we correct our principal perturbation w_o by w_c , i.e. $w_{q+1} = w_o + w_c$ so that w_{q+1} is the curl of a vector field. In particular, in view of the identity (5.8) we define

$$w_c := \frac{-i}{\lambda_{q+1}} \sum_{i,k \neq 0} \left[\text{curl} \left((\rho_{q,i})^{1/2} \frac{\nabla \Phi_i^T (k \times a_k(\tilde{R}_{q,i}))}{|k|^2} \right) \right] e^{i\lambda_{q+1}k \cdot \Phi_i} =: \sum_{i,k \neq 0} c_{i,k} e^{i\lambda_{q+1}k \cdot \Phi_i}.$$

Then from (5.8) and the identity (see for instance [DS17])

$$\text{curl}(\nabla \Phi^T U(\Phi_i)) = \text{cof} \nabla \Phi_i^T (\text{curl} U)(\Phi_i) = \nabla \Phi^{-1}(\text{curl} U)(\Phi_i)$$

one can check that

$$w_{q+1} = w_o + w_c = \frac{-1}{\lambda_{q+1}} \text{curl} \left(\sum_{i,k \neq 0} (\nabla \Phi_i)^T \left(\frac{ik \times b_{k,i}}{|k|^2} \right) e^{i\lambda_{q+1}k \cdot \Phi_i} \right). \quad (5.20)$$

5.4. The final Reynolds stress. Upon letting

$$\bar{R}_q = \sum_i R_{q,i},$$

the new Reynolds stress is thus defined as

$$\mathring{R}_{q+1} = \underbrace{\mathcal{R}(w_{q+1} \cdot \nabla \bar{v}_q)}_{\text{Nash error}} + \underbrace{\mathcal{R}(\partial_t w_{q+1} + \bar{v}_q \cdot \nabla w_{q+1})}_{\text{Transport error}} + \underbrace{\mathcal{R} \text{div}(-\bar{R}_q + (w_{q+1} \otimes w_{q+1}))}_{\text{Oscillation error}}. \quad (5.21)$$

Notice that all three terms in (5.21) are of the form $\mathcal{R}f$, where f is either a divergence or a curl, and thus has zero mean. With this definition and Proposition 4.1, one may verify that

$$\begin{cases} \partial_t v_{q+1} + \text{div}(v_{q+1} \otimes v_{q+1}) + \nabla p_{q+1} = \text{div}(\mathring{R}_{q+1}), \\ \text{div} v_{q+1} = 0, \end{cases}$$

where the new pressure is defined by

$$p_{q+1} = \bar{p}_q + |w_{q+1}|^2 - \sum_i \rho_{q,i}. \quad (5.22)$$

5.5. Estimates on the perturbation.

Proposition 5.7. *For $t \in \tilde{I}_i$ and any $N \geq 0$*

$$\|(\nabla\Phi_i)^{-1}\|_N + \|\nabla\Phi_i\|_N \lesssim \ell^{-N}, \quad (5.23)$$

$$\|\tilde{R}_{q,i}\|_N \lesssim \ell^{-N}, \quad (5.24)$$

$$\|b_{i,k}\|_N \lesssim \delta_{q+1}^{1/2} |k|^{-6} \ell^{-N}, \quad (5.25)$$

$$\|c_{i,k}\|_N \lesssim \delta_{q+1}^{1/2} \lambda_{q+1}^{-1} |k|^{-6} \ell^{-N-1}. \quad (5.26)$$

It is important to notice that the symbol \lesssim denotes a dependence of the constants in the estimates from N , α , β and M , but not upon k or a .

Proof of Proposition 5.7. From (2.30), (B.5) and (B.6) we obtain

$$\|\nabla\Phi_i\|_N \lesssim 1 + \tau_q \|D\bar{v}_q\|_N \lesssim 1 + \tau_q \delta_q^{1/2} \lambda_q \ell^{-N}.$$

Using the fact that $\|\nabla\Phi_i - \text{Id}\|_0 \leq 1/2$ (see (5.10)), the estimate (5.23) follows (indeed it gives the slightly better estimate $\lesssim 1 + \ell^{-N+2\alpha}$, but the other is still enough for our purposes).

Recalling property (iv) of η_i we see that $\rho_{q,i}$ is a function of t only on $\text{supp } \overset{\circ}{\bar{R}}_q$, i.e.

$$\rho_{q,i}(x, t) = \frac{\rho_q(t)}{\sum_j \int_{\mathbb{T}^3} \eta_j^2(y, t) dy}.$$

Thus,

$$\frac{R_{q,i}}{\rho_{q,i}} = \text{Id} - \frac{\sum_j \int_{\mathbb{T}^3} \eta_j^2(y, t) dy}{\rho_q(t)} \overset{\circ}{\bar{R}}_q, \quad (5.27)$$

so that by (5.11) and (4.10) we obtain

$$\left\| \frac{R_{q,i}}{\rho_{q,i}} \right\|_N \lesssim 1 + \frac{\lambda_q^\alpha}{\delta_{q+1}} \left\| \overset{\circ}{\bar{R}}_q \right\|_N \ell^{-N} \lesssim \ell^{-N}, \quad (5.28)$$

where we have applied the crude estimate $\lesssim 1 + \|\overset{\circ}{\bar{R}}_q\|_{N+\alpha} \lambda_q^\alpha \delta_{q+1}^{-1} \lesssim 1 + \ell^{-N+\alpha} \lambda_q^\alpha \lesssim \ell^{-N}$.

Therefore, using Lemma 5.4 and property (v):

$$\left\| \tilde{R}_{q,i} \right\|_N \lesssim \|\nabla\Phi_i\|_N \|\nabla\Phi_i\|_0 + \left\| \frac{R_{q,i}}{\rho_{q,i}} \right\|_N \lesssim \|\nabla\Phi_i\|_N \|\nabla\Phi_i\|_0 + \ell^{-N}.$$

The estimate (5.24) then follows from (5.23).

The estimate (5.25) follows as a consequence of (5.5), (5.12) and (5.24). The estimate (5.26) follows as a consequence of (5.5), (5.12), (5.23) and (5.24). \square

Corollary 5.8. *Assuming a is sufficiently large, the perturbations w_o , w_c and w_q satisfy the following estimates*

$$\|w_o\|_0 + \frac{1}{\lambda_{q+1}} \|w_o\|_1 \leq \frac{M}{4} \delta_{q+1}^{1/2} \quad (5.29)$$

$$\|w_c\|_0 + \frac{1}{\lambda_{q+1}} \|w_c\|_1 \lesssim \delta_{q+1}^{1/2} \ell^{-1} \lambda_{q+1}^{-1} \quad (5.30)$$

$$\|w_{q+1}\|_0 + \frac{1}{\lambda_{q+1}} \|w_{q+1}\|_1 \leq \frac{M}{2} \delta_{q+1}^{1/2} \quad (5.31)$$

where the constant M depends solely on the constant c_0 in (5.16). In particular, we obtain (2.35).

Proof of Corollary 5.8. Taking into account (5.10), we conclude $\|(\nabla\Phi_i)^{-1}\|_0 \leq 2$ on $\text{supp}(\eta_i)$. Thus, taking into account that the $w_{o,i}$ have disjoint supports, from Lemma 5.5 we conclude

$$\|w_o\| \leq 2\delta_{q+1}^{1/2} \sum_{k \neq 0} \frac{\bar{M}}{|k|^4} \leq \frac{M}{32} \delta_{q+1}^{1/2}. \quad (5.32)$$

To estimate $\|w_o\|_1$ we observe first that

$$\left\| \nabla(e^{i\lambda_{q+1}k \cdot \Phi_i}) \right\|_0 \leq \lambda_{q+1} |k| \|\nabla\Phi_i\|_0 \leq 2\lambda_{q+1} |k|. \quad (5.33)$$

Compute now

$$\nabla w_{o,i} = \sum_k (\nabla\Phi_i)^{-1} b_{i,k} \nabla(e^{i\lambda_{q+1}k \cdot \Phi_i}) + \sum_k \nabla((\nabla\Phi_i)^{-1} b_{i,k}) e^{i\lambda_{q+1}k \cdot \Phi_i}.$$

In particular, from (5.33), Lemma 5.5 and Proposition 5.7 (taking into account that the supports of the $w_{o,i}$ are disjoint), we conclude

$$\|\nabla w_o\|_0 \leq 4\delta_{q+1}^{1/2} \lambda_{q+1} \sum_{k \neq 0} \frac{\bar{M}}{|k|^4} + C\delta_{q+1}^{1/2} \ell^{-1} \sum_{k \neq 0} \frac{1}{|k|^6} \leq \frac{M}{16} \delta_{q+1}^{1/2} \lambda_{q+1} + \bar{C} \delta_{q+1}^{1/2} \ell^{-1}, \quad (5.34)$$

where the constant \bar{C} depends upon β, α and M , but not upon a . In particular, summing (5.32) and (5.34) we achieve

$$\|w_o\|_0 + \lambda_{q+1}^{-1} \|w_o\|_1 \leq \frac{M}{8} \delta_{q+1}^{1/2} + \bar{C} (\lambda_{q+1} \ell)^{-1} \delta_{q+1}^{1/2}. \quad (5.35)$$

By our definition of the various parameters we get

$$(\ell \lambda_{q+1})^{-1} = \frac{\delta_q^{1/2} \lambda_q^{1+3\alpha/2}}{\delta_{q+1}^{1/2} \lambda_{q+1}} = \frac{\lambda_q^{1-\beta+3\alpha/2}}{\lambda_{q+1}^{1-\beta}} \leq \tilde{C} a^{b^q(1-\beta+3\alpha/2)-b^{q+1}(1-\beta)}$$

where the constant \tilde{C} depends on (2.18). Having chosen α small enough so that $b > 1 - \beta + 3\alpha/2 / 1 - \beta$, for a sufficiently large we achieve that the right hand side of (5.35) is smaller than $M/4\delta_{q+1}^{1/2}$.

The estimate (5.30) follows as a direct consequence of (5.26) and (5.33).

Combining (5.29) and (5.30) we achieve

$$\|w_{q+1}\|_0 + \frac{1}{\lambda_{q+1}} \|w_{q+1}\|_1 \leq \delta_{q+1}^{1/2} \left(\frac{M}{4} + C(\ell \lambda_{q+1})^{-1} \right), \quad (5.36)$$

where the constant C depends upon β, α and M , but not upon a . Hence, arguing as above, if $b > 1 - \beta + 3\alpha/2 / 1 - \beta$ then (5.31) holds for a sufficiently large (depending on β, α and M). \square

Let us define $D_{t,q} := \partial_t + \bar{v}_q \cdot \nabla$ to be the material derivative associated with \bar{v}_q . We then have

Proposition 5.9. *For $t \in \tilde{I}_i$ and $N \geq 0$ we have*

$$\|D_{t,q} \nabla \Phi_i\|_N \lesssim \delta_q^{1/2} \lambda_q \ell^{-N} \quad (5.37)$$

$$\|D_{t,q} \tilde{R}_{q,i}\|_N \lesssim \tau_q^{-1} \ell^{-N} \quad (5.38)$$

$$\|D_{t,q} c_{i,k}\|_N \lesssim \delta_{q+1}^{1/2} \tau_q^{-1} \ell^{-N-1} \lambda_{q+1}^{-1} |k|^{-6}. \quad (5.39)$$

Proof of Proposition 5.9. Observe that

$$D_{t,q}\nabla\Phi_i = -\nabla\Phi_i D\bar{v}_q$$

In particular,

$$\|D_{t,q}\nabla\Phi_i\|_N \lesssim \|\nabla\Phi_i\|_0 \|\bar{v}_q\|_{N+1} + \|\nabla\Phi_i\|_N \|\bar{v}_q\|_1.$$

Thus (5.37) follows from (4.7) and (5.23). Next, we observe that

$$D_{t,q}\rho_{q,i} = \partial_t\rho_{q,i} + \bar{v}_q \cdot \nabla\rho_{q,i}$$

and thus we can estimate

$$\|D_{t,q}\rho_{q,i}\|_N \lesssim \|\partial_t\rho_{q,i}\|_N + \|\rho_{q,i}\|_{N+1} \|\bar{v}_q\|_0 + \|\bar{v}_q\|_N \|\rho_{q,i}\|_1.$$

Recall that $\|\bar{v}_q\|_0 \leq \|v_\ell\|_0 + \|v_\ell - v_q\|_0 \lesssim 1 \lesssim \tau_q^{-1}$ and so from (4.7) we conclude $\|\bar{v}_q\|_N \leq \tau_q^{-1}\ell^{-N}$. Combining the latter estimate with (5.13) and (5.15) we achieve

$$\|D_{t,q}\rho_{q,i}\|_N \lesssim \delta_{q+1}\tau_q^{-1}\ell^{-N}. \quad (5.40)$$

Differentiating (5.27) we have

$$D_{t,q}(\rho_{q,i}^{-1}R_{q,i}) = -\left(\partial_t \frac{\sum_j \int_{\mathbb{T}^3} \eta_j^2(y,t) dy}{\rho_q(t)}\right) \bar{R}_q - \frac{\sum_j \int_{\mathbb{T}^3} \eta_j^2(y,t) dy}{\rho_q(t)} D_{t,q} \bar{R}_q. \quad (5.41)$$

Thus we can estimate, using (4.10) and (4.11):

$$\begin{aligned} \|D_{t,q}(\rho_{q,i}^{-1}R_{q,i})\|_N &\lesssim \delta_{q+1}^{-1} \delta_q^{1/2} \lambda_q^{1+2\alpha} \|\bar{R}_q\|_N + \tau_q^{-1} \delta_{q+1}^{-1} \lambda_q^\alpha \|\bar{R}_q\|_N + \delta_{q+1}^{-1} \lambda_q^\alpha \|D_{t,q} \bar{R}_q\|_N \\ &\lesssim \delta_q^{1/2} \lambda_q^{1+2\alpha} \ell^{-N+\alpha} + \tau_q^{-1} \lambda_q^\alpha \ell^{-N+\alpha} + \lambda_q^\alpha \delta_q^{1/2} \lambda_q \ell^{-N-\alpha} \lesssim \tau_q^{-1} \ell^{-N}. \end{aligned} \quad (5.42)$$

Differentiating (5.9) we achieve

$$D_{t,q} \tilde{R}_{q,i} = D_{t,q} \nabla\Phi_i(\rho_{q,i}^{-1}R_{q,i}) \nabla\Phi_i^T + \nabla\Phi_i D_{t,q}(\rho_{q,i}^{-1}R_{q,i}) \nabla\Phi_i^T + \nabla\Phi_i(\rho_{q,i}^{-1}R_{q,i})(D_{t,q} \nabla\Phi_i)^T.$$

Thus we can estimate

$$\begin{aligned} \|D_{t,q} \tilde{R}_{q,i}\|_N &\lesssim \|D_{t,q} \nabla\Phi_i\|_N \|(\rho_{q,i}^{-1}R_{q,i})\|_0 + \|D_{t,q} \nabla\Phi_i\|_0 \|(\rho_{q,i}^{-1}R_{q,i})\|_N \\ &\quad + \|D_{t,q} \nabla\Phi_i\|_0 \|(\rho_{q,i}^{-1}R_{q,i})\|_0 \|\nabla\Phi_i\|_N + \|D_{t,q}(\rho_{q,i}^{-1}R_{q,i})\|_N + \|D_{t,q}(\rho_{q,i}^{-1}R_{q,i})\|_0 \|\nabla\Phi_i\|_N. \end{aligned}$$

Using (5.37), (5.42), (5.28) and (5.23), we conclude (5.38).

Finally, the estimate (5.39) follows as a consequence of (5.5), Lemma 5.4, Proposition 5.7, (5.37), and (5.38). \square

6. PROOF OF PROPOSITION 2.1

In this section we complete the proof of Proposition 2.1 by proving the remaining estimates (2.36) and (2.37).

6.1. Estimates of the new Reynolds stress error. In the proposition below we prove the inductive estimates on \mathring{R}_{q+1} :

Proposition 6.1. *The Reynolds stress error \mathring{R}_{q+1} defined in (5.21) satisfies the estimate*

$$\left\| \mathring{R}_{q+1} \right\|_0 \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-4\alpha}}. \quad (6.1)$$

In particular, (2.36) holds.

6.1.1. *Nash error.* We just write this term as

$$\mathcal{R}(w_{q+1} \cdot \nabla \bar{v}_q) = \sum_i \sum_{k \neq 0} \mathcal{R} \left(\left((\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i} + c_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i} \right) \cdot \nabla \bar{v}_q \right).$$

Using Proposition C.2 we bound for $t \in \tilde{I}_i$

$$\begin{aligned} & \left\| \mathcal{R} \left((\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i} \cdot \nabla \bar{v}_q \right) \right\|_\alpha \\ & \lesssim \frac{\| \nabla \Phi_i^{-1} b_{i,k} \cdot \nabla \bar{v}_q \|_0}{\lambda_{q+1}^{1-\alpha}} + \frac{\| \nabla \Phi_i^{-1} b_{i,k} \cdot \nabla \bar{v}_q \|_{N+\alpha} + \| \nabla \Phi_i^{-1} b_{i,k} \cdot \nabla \bar{v}_q \|_0 \| \Phi_i \|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} \\ & \lesssim \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{1-\alpha} |k|^6} + \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{N-\alpha} \ell^{N+\alpha} |k|^6}. \end{aligned}$$

Now, provided α is sufficiently small we claim that we can first fix a suitable N and then choose a large enough, so that

$$\frac{1}{\lambda_{q+1}^{N-\alpha} \ell^{N+\alpha}} \leq \frac{1}{\lambda_{q+1}^{1-\alpha}}.$$

Such choice is equivalent to $\lambda_{q+1}^{(N-1)-(N-\alpha)\beta} \geq \lambda_q^{(1-\beta+3\alpha/2)(N+\alpha)}$. Taking the logarithms in base a , we need the condition

$$b^{q+1}((N-1) - (N-\alpha)\beta) > b^q(N+\alpha) \left(1 - \beta + \frac{3\alpha}{2} \right),$$

which would determine the needed N . In order to show that for α sufficiently small we can choose such an N , we just need to verify the existence of N such that $b((N-1) - N\beta) > N(1-\beta)$. The latter is equivalent to $(b-1)(N-1)(1-\beta) > (1-\beta) + b\beta$ which in turn, since $\beta < 1/3$ and $b > 1$, can certainly be satisfied for N large enough. Finally, having chosen first $\alpha > 0$ and then N according to the above requirement, we can then take $a \gg 1$ large enough to beat the eventual geometric constant due to (2.18). Hence we achieve

$$\left\| \mathcal{R} \left((\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i} \cdot \nabla \bar{v}_q \right) \right\|_\alpha \lesssim \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{1-\alpha} |k|^6}. \quad (6.2)$$

For the second term in the Nash error we again use Corollary C.2 to obtain

$$\begin{aligned} \left\| \mathcal{R} \left(c_{i,k} e^{i\lambda_{q+1} k \cdot \Phi_i} \cdot \nabla \bar{v}_q \right) \right\|_\alpha & \lesssim \frac{\| c_{i,k} \cdot \nabla \bar{v}_q \|_0}{\lambda_{q+1}^{1-\alpha}} + \frac{\| c_{i,k} \cdot \nabla \bar{v}_q \|_{N+\alpha} + \| c_{i,k} \cdot \nabla \bar{v}_q \|_0 \| \Phi_i \|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} \\ & \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\ell \lambda_{q+1}^{2-\alpha} |k|^6} + \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\ell^{N+1-\alpha} \lambda_{q+1}^{N+1-\alpha} |k|^6} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha} |k|^6} \end{aligned} \quad (6.3)$$

where again we assume to have fixed first N and then a large enough. We also implicitly used that

$$\ell \lambda_{q+1} \geq 1, \quad (6.4)$$

which is equivalent to $\lambda_{q+1}^{1-\beta} \geq \lambda_q^{1-\beta+3\alpha/2}$. The latter inequality follows from (2.11) and (2.18), upon taking logarithms in base a , choosing first α so that $(b-1)(1-\beta) \geq 3\alpha$, and then a sufficiently large so that $\frac{(b-1)}{10} \geq \log_a(4\pi)$.

Summing over the frequencies and using that $\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} |k|^{-6} < \infty$, we achieve

$$\mathcal{R}(w_{q+1} \cdot \nabla \bar{v}_q) \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha}}. \quad (6.5)$$

6.1.2. *Transport error.* We split the transport error into two parts

$$(\partial_t + \bar{v}_q \cdot \nabla)w_{q+1} = (\partial_t + \bar{v}_q \cdot \nabla)w_o + (\partial_t + \bar{v}_q \cdot \nabla)w_c.$$

Applying (5.18) yields

$$\begin{aligned} (\partial_t + \bar{v}_q \cdot \nabla)w_o &= \sum_{i,k} (\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1}k \cdot \Phi_i} \\ &\quad + \sum_{i,k} (\nabla \Phi_i)^{-1} (\partial_t + \bar{v}_q \cdot \nabla) \left(\rho_{q,i}^{1/2} a_k(\tilde{R}_i) \right) e^{i\lambda_{q+1}k \cdot \Phi_i}. \end{aligned} \quad (6.6)$$

We then apply Corollary C.2 to obtain for the first term in (6.6)

$$\begin{aligned} &\left\| \mathcal{R} \left((\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1}k \cdot \Phi_i} \right) \right\|_{\alpha} \\ &\lesssim \frac{\|(\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k}\|_0}{\lambda_{q+1}^{1-\alpha}} + \frac{\|(\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k}\|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} + \frac{\|(\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k}\|_0 \|\Phi_i\|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}}. \end{aligned} \quad (6.7)$$

We use Proposition 5.7 and Proposition 4.3 to estimate

$$\begin{aligned} \|(\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k}\|_{N+\alpha} &\lesssim \|\nabla \bar{v}_q\|_{N+\alpha} \|(\nabla \Phi_i)^{-1}\|_{\alpha} \|b_{i,k}\|_{\alpha} \\ &\quad + \|\nabla \bar{v}_q\|_{\alpha} \|(\nabla \Phi_i)^{-1}\|_{N+\alpha} \|b_{i,k}\|_{\alpha} + \|\nabla \bar{v}_q\|_{\alpha} \|(\nabla \Phi_i)^{-1}\|_{\alpha} \|b_{i,k}\|_{N+\alpha} \\ &\lesssim \delta_{q+1}^{1/2} \ell^{-N-3\alpha} \lesssim \delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q \ell^{-N-1-3\alpha}. \end{aligned}$$

Arguing in a similar fashion for the third summand in (6.7), we achieve

$$\left\| \mathcal{R} \left((\nabla \bar{v}_q)^T (\nabla \Phi_i)^{-1} b_{i,k} e^{i\lambda_{q+1}k \cdot \Phi_i} \right) \right\|_{\alpha} \lesssim \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{1-\alpha} |k|^6} + \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{N-\alpha} \ell^{N+1+3\alpha} |k|^6} \lesssim \frac{\lambda_q \delta_{q+1}^{1/2} \delta_q^{1/2}}{\lambda_{q+1}^{1-\alpha} |k|^6},$$

where in the last inequality, as in the previous section, we have assumed α sufficiently small and N appropriately chosen.

For the second term in (6.6), let us define

$$d_{i,k}(x, t) := (\nabla \Phi_i(x, t))^{-1} D_{t,q} \left((\rho_{q+1,i}(x, t))^{1/2} a_k(\tilde{R}_i(x, t)) \right).$$

Using (5.23), (5.13), (5.15) and (5.38) and again assuming N sufficiently large, arguing as above we conclude

$$\begin{aligned} \left\| \mathcal{R} \left(d_{i,k} e^{i\lambda_{q+1}k \cdot \Phi_i} \right) \right\|_{\alpha} &\lesssim \frac{\|d_{i,k}\|_0}{\lambda_{q+1}^{1-\alpha}} + \frac{\|d_{i,k}\|_{N+\alpha} + \|d_{i,k}\|_0 \|\Phi_i\|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} \\ &\lesssim \frac{\delta_{q+1}^{1/2}}{\tau_q \lambda_{q+1}^{1-\alpha} |k|^6} = \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha}} \ell^{-2\alpha} |k|^{-6} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-4\alpha}} |k|^{-6}, \end{aligned}$$

where we have used $\ell^{-2\alpha} \leq \lambda_q^{3\alpha} \leq \lambda_{q+1}^{3\alpha}$ (see (2.21)).

Now we consider the term involving the material derivative of the correction. Observe

$$(\partial_t + \bar{v}_q \cdot \nabla)w_c = \sum_{i,k} (D_{t,q} c_{i,k}) e^{i\lambda_{q+1}k \cdot \Phi_i}$$

Then applying Corollary C.2 and (5.39) yields

$$\begin{aligned} \left\| \mathcal{R} \left((D_{t,q} c_{i,k}) e^{i\lambda_{q+1} k \cdot \Phi_i} \right) \right\|_{\alpha} &\lesssim \frac{\|D_{t,q} c_{i,k}\|_0}{\lambda_{q+1}^{1-\alpha}} + \frac{\|D_{t,q} c_{i,k}\|_{N+\alpha} + \|D_{t,q} c_{i,k}\|_0 \|\Phi_i\|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} \\ &\lesssim \frac{\delta_{q+1}^{1/2}}{\tau_q \ell \lambda_{q+1}^{2-\alpha} |k|^6} \lesssim \frac{\delta_{q+1}^{1/2}}{\tau_q \lambda_{q+1}^{1-\alpha} |k|^6} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-3\alpha}} |k|^{-6} \end{aligned}$$

where we used (5.23), (5.39) and (6.4).

Again, summing over $k \neq 0$ we reach the inequality

$$\|\mathcal{R}(\partial_t w_{q+1} + \bar{v}_q \cdot \nabla w_{q+1})\|_{\alpha} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-3\alpha}}. \quad (6.8)$$

6.1.3. *Oscillation error.* Recall the oscillation error may be written as

$$\begin{aligned} &\mathcal{R} \operatorname{div} \left(-\bar{R}_q + w_{q+1} \otimes w_{q+1} \right) \\ &= \underbrace{\mathcal{R} \operatorname{div} \left(-\bar{R}_q + w_o \otimes w_o \right)}_{=: \mathcal{O}_1} + \underbrace{\mathcal{R} \operatorname{div} \left(w_o \otimes w_c + w_c \otimes w_o + w_c \otimes w_c \right)}_{=: \mathcal{O}_2}. \end{aligned}$$

For the second term we proceed as follows:

$$\begin{aligned} \|\mathcal{O}_2\|_{\alpha} &\lesssim \|w_o \otimes w_c + w_c \otimes w_o + w_c \otimes w_c\|_{\alpha} \\ &\lesssim \|w_o\|_0 \|w_c\|_{\alpha} + \|w_o\|_{\alpha} \|w_c\|_0 + \|w_c\|_{\alpha}^2 \lesssim \frac{\delta_{q+1}}{\ell \lambda_{q+1}^{1-\alpha}} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha}}. \end{aligned} \quad (6.9)$$

Now consider \mathcal{O}_1 . Due to the supports of the cutoffs η_j being mutually disjoint, we have

$$\mathcal{O}_1 = \mathcal{R} \operatorname{div} \left(-\bar{R}_q + \sum_i w_{o,i} \otimes w_{o,i} \right).$$

Using the definition of $w_{o,i}$ and (5.6), the tensor $w_{o,i} \otimes w_{o,i}$ may be written as

$$\begin{aligned} w_{o,i} \otimes w_{o,i} &= \rho_{q,i} \nabla \Phi_i^{-1} (W \otimes W) (\tilde{R}_{q,i}, \lambda_{q+1} \Phi_i) \nabla \Phi_i^{-T} \\ &= \rho_{q,i} \nabla \Phi_i^{-1} \tilde{R}_{q,i} \nabla \Phi_i^{-T} + \sum_{k \neq 0} \rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T} e^{i\lambda_{q+1} k \cdot \Phi_i} \\ &= R_{q,i} + \sum_{k \neq 0} \rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T} e^{i\lambda_{q+1} k \cdot \Phi_i}. \end{aligned} \quad (6.10)$$

On the other hand, recalling (5.7)

$$\nabla \Phi_i^{-1} C_k \nabla \Phi_i^{-T} \nabla \Phi_i^T k = 0,$$

consequently

$$\operatorname{div} \left(\sum_i w_{o,i} \otimes w_{o,i} - R_{q,i} \right) = \sum_{i, k \neq 0} \operatorname{div} (\rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T}) e^{i\lambda_{q+1} k \cdot \Phi_i}$$

Thus, by Proposition C.2

$$\begin{aligned}
\|\mathcal{O}_1\|_\alpha &\lesssim \sum_i \sum_{k \neq 0} \left\| \frac{\operatorname{div}(\rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T})}{\lambda_{q+1}^{1-\alpha}} \right\|_0 \\
&+ \sum_i \sum_{k \neq 0} \frac{\left\| \operatorname{div}(\rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T}) \right\|_{N+\alpha}}{\lambda_{q+1}^{N-\alpha}} + \left\| \operatorname{div}(\rho_{q,i} \nabla \Phi_i^{-1} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T}) \right\|_0 \|\Phi_i\|_{N+\alpha} \\
&\lesssim \sum_i \sum_{k \neq 0} \frac{\delta_{q+1}}{\ell \lambda_{q+1}^{1-\alpha} |k|^6} \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha}}, \tag{6.11}
\end{aligned}$$

where we have used (5.7) and, as in the previous sections, a large choice of N to absorb the estimates of the second line in that for the first line. Clearly, (6.9) and (6.11) give

$$\|\mathcal{R} \operatorname{div}(-\bar{R}_q + w_{q+1} \otimes w_{q+1})\|_\alpha \lesssim \frac{\delta_{q+1}^{1/2} \delta_q^{1/2} \lambda_q}{\lambda_{q+1}^{1-\alpha}}. \tag{6.12}$$

6.1.4. *Conclusion.* Clearly (6.1) follows from (6.5), (6.8), (6.12) and (5.21).

6.2. **Energy iterate.** (2.10) in the following proposition:

Proposition 6.2. *The energy of v_{q+1} satisfies the following estimate:*

$$\left| e(t) - \int_{\mathbb{T}^3} |v_{q+1}|^2 dx - \frac{\delta_{q+2}}{2} \right| \lesssim \frac{\delta_q^{1/2} \delta_{q+1}^{1/2} \lambda_q^{1+2\alpha}}{\lambda_{q+1}}.$$

In particular, the estimate (2.37) holds.

Proof of Proposition 6.2. By definition we have

$$\int_{\mathbb{T}^3} |v_{q+1}|^2 dx = \int_{\mathbb{T}^3} |\bar{v}_q|^2 dx + 2 \int_{\mathbb{T}^3} w_{q+1} \cdot \bar{v}_q dx + \int_{\mathbb{T}^3} |w_{q+1}(x, t)|^2 dx$$

We also recall that

$$\sum_i \int_{\mathbb{T}^3} \operatorname{tr} R_{q,i}(x, t) dx = 3 \sum_i \int_{\mathbb{T}^3} \rho_{q,i}(x, t) dx = 3\rho_q(t) = e(t) - \frac{\delta_{q+2}}{2} - \int_{\mathbb{T}^3} |\bar{v}_q|^2 dx.$$

By integrating by parts once and using the identity (5.20) and the estimates (5.23) and (5.25) we obtain

$$\left| \int_{\mathbb{T}^3} w_{q+1} \cdot \bar{v}_q dx \right| = \frac{1}{\lambda_{q+1}} \sum_i \sum_{k \neq 0} \left\| (\nabla \Phi_i)^T \left(\frac{ik \times b_k}{|k|^2} \right) \right\|_0 \|\bar{v}_q\|_1 \lesssim \frac{\delta_q^{1/2} \delta_{q+1}^{1/2} \lambda_q}{\lambda_{q+1}}.$$

Using (5.29) and (5.30) yields

$$\left| \int_{\mathbb{T}^3} 2w_o \cdot w_c + |w_c|^2 dx \right| \lesssim \frac{\delta_{q+1}}{\ell \lambda_{q+1}} \stackrel{(2.19)}{=} \frac{\delta_q^{1/2} \delta_{q+1}^{1/2} \lambda_q^{1+2\alpha}}{\lambda_{q+1}}.$$

Finally, recall from (6.10) that

$$\int_{\mathbb{T}^3} |w_o(x, t)|^2 dx = \sum_i \int_{\mathbb{T}^3} \operatorname{tr} R_{q,i}(x, t) dx + \int_{\mathbb{T}^3} \sum_{i, k \neq 0} \rho_{q,i} \nabla \Phi_i^{-1} \operatorname{tr} C_k(\tilde{R}_{q,i}) \nabla \Phi_i^{-T} e^{i\lambda_{q+1}k \cdot \Phi_i} dx$$

and thus it remains to bound the second term. Set $e_{q,i} := \rho_{q,i} \nabla \Phi_i^{-1} \operatorname{tr} C_k(\tilde{R}_i) \nabla \Phi_i^{-T}$ and use Proposition 5.7 and Lemma 5.4 to conclude

$$\|e_{q,i}\|_N \lesssim \delta_{q+1} \ell^{-N}.$$

Next observe that at any given time at most two $e_{q,i}$ are nonvanishing. Hence use (C.1) in Proposition C.2 to bound

$$\left| \int_{\mathbb{T}^3} \sum_i \sum_{k \neq 0} \rho_i \nabla \Phi_i^{-1} \operatorname{tr} C_k(\tilde{R}_i) \nabla \Phi_i^{-T} e^{i\lambda_{q+1}k \cdot \Phi_i} dx \right| \lesssim \sum_{k \neq 0} \frac{\delta_{q+1} \ell^{-N}}{\lambda_{q+1}^N |k|^N}.$$

As already argued several time, we can choose N such that $\delta_{q+1} \ell^{-N} / \lambda_{q+1}^N \leq \delta_{q+1} \delta_q^{1/2} \lambda_q / \lambda_{q+1}$. Assuming in addition that N is larger than 4 (so that the series is summable), we obtain the desired estimate. \square

7. AN h -PRINCIPLE

In order to prove Theorem 1.3, let us first state a variant of Proposition 3.1 from [DS17] that follows from the estimates in Section 5 used to prove the proposition in [DS17]:

Theorem 7.1. *Let $(\bar{v}, \bar{p}, \bar{R})$ be a smooth strict subsolution of the Euler equations on $\mathbb{T}^3 \times [0, T]$ and fix $0 < \gamma < 1$. Then there exists $\varepsilon_0 > 0$ such that for any $\varepsilon < \varepsilon_0$, and for any sufficiently large λ depending on ε_0 and $(\bar{v}, \bar{p}, \bar{R})$, we have the following: There exists a smooth solution (v, p, R) of (1.3) satisfying the estimates*

$$\begin{aligned} \|v - \bar{v}\|_{H^{-1}} &\leq C\lambda^{-1} \\ \|v\|_0 + \lambda^{-1} \|v\|_1 &\leq C \\ \|v \otimes v + R - \bar{v} \otimes \bar{v} - \bar{R}\|_{H^{-1}} &\leq C\lambda^{\gamma-1} \\ \|\mathring{R}\|_0 &\leq C\lambda^{\gamma-1} \\ \|\operatorname{tr} R\|_0 &\leq \varepsilon, \end{aligned}$$

where C depends solely on $(\bar{v}, \bar{p}, \bar{R})$, and \mathring{R} is the traceless part of R . Moreover setting

$$e(t) := \int_{\mathbb{T}^3} |\bar{v}|^2 + \operatorname{tr} \bar{R} dx \tag{7.1}$$

for any $t \in [0, T]$ we have

$$\frac{\varepsilon}{2} \leq e(t) - \int_{\mathbb{T}^3} |v|^2 dx \leq \varepsilon.$$

Proof of Theorem 1.3. Fix $k \geq 1$ and let $\varepsilon_k < \varepsilon_0$. We apply Theorem 7.1 with $\gamma = \alpha$ and $\lambda = \lambda_0$, where here (α, λ_0) are given in the statement of Proposition 2.1, and where we take a sufficiently

large such that λ_0 is sufficiently large (in terms of ε_k and $(\bar{v}, \bar{p}, \bar{R})$), so that the hypothesis of Theorem 7.1 is satisfied. We obtain (v, p, R) satisfying

$$\|v - \bar{v}\|_{H^{-1}} \leq C\lambda_0^{-1} \quad (7.2)$$

$$\|v\|_0 + \lambda_0^{-1}\|v\|_1 \leq C \quad (7.3)$$

$$\|v \otimes v + R - \bar{v} \otimes \bar{v} - \bar{R}\|_{H^{-1}} \leq C\lambda_0^{\alpha-1} \quad (7.4)$$

$$\|\mathring{R}\|_0 \leq C\lambda_0^{\alpha-1} \quad (7.5)$$

$$\|\text{tr } R\|_0 \leq \varepsilon_k, \quad (7.6)$$

and the function $e(t)$ as defined by (7.1) obeys

$$\frac{\varepsilon_k}{2} \leq e(t) - \int_{\mathbb{T}^3} |v|^2 dx \leq \varepsilon_k. \quad (7.7)$$

Analogous to the proof of Theorem 1.1, we set

$$\Gamma = \frac{\delta_1^{1/2}}{\varepsilon_k^{1/2}}$$

and rescale (v, p, R) to obtain

$$\tilde{v}_0(x, t) := \Gamma v(x, \Gamma t), \quad \tilde{p}_0(x, t) := \Gamma^2 p(x, \Gamma t) \quad \text{and} \quad \tilde{R}_0(x, t) := \Gamma^2 R(x, \Gamma t),$$

so that (v_0, p_0, R_0) also solves (1.3). Moreover, we have the estimates

$$\begin{aligned} \|\tilde{v}_0\|_0 + \lambda_0^{-1}\|\tilde{v}_0\|_1 &\leq \frac{C\delta_1^{1/2}}{\varepsilon_k^{1/2}} \\ \|\mathring{\tilde{R}}_0\|_0 &\leq \frac{C\delta_1}{\varepsilon_k\lambda_0^{1-\alpha}}. \end{aligned} \quad (7.8)$$

Choosing α sufficiently small and choosing a sufficiently large depending on ε_k , C , and M , we obtain

$$\frac{C\delta_1^{1/2}}{\varepsilon_k^{1/2}} \leq \min(M\delta_0^{1/2}, 1 - \delta_0) \quad \text{and} \quad \frac{C}{\varepsilon_k\lambda_0^{1-\alpha}} \leq \lambda_0^{-3\alpha}.$$

from which we obtain (2.7), (2.8), and (2.9).

If in addition we set

$$\tilde{e}(t) = \Gamma^2 e(\Gamma t)$$

then from (7.7) we obtain

$$\frac{\delta_1}{2} \leq \tilde{e}(t) - \int_{\mathbb{T}^3} |\tilde{v}_0|^2 dx \leq \delta_1,$$

and hence we obtain (2.10) for $q = 0$. Letting a be sufficiently large, we also obtain (2.1).

Applying Proposition 2.1 and arguing as was done in the proof of Theorem 1.1 we obtain a solution (\tilde{v}, \tilde{p}) to the Euler equations satisfying

$$\int_{\mathbb{T}^3} |\tilde{v}|^2 dx = \tilde{e}(t). \quad (7.9)$$

Moreover, by (2.12) we have the estimate

$$\|\tilde{v} - \tilde{v}_0\|_0 \lesssim \delta_1^{1/2}. \quad (7.10)$$

Lastly, we define (v_k, p_k) by the rescaling

$$v_k := \Gamma^{-1}\tilde{v}(x, \Gamma^{-1}t) \quad \text{and} \quad p_k := \Gamma^{-2}\tilde{p}(x, \Gamma^{-1}t).$$

Then (v_k, p_k) is a solution to the Euler equations, satisfying (1.4) as a consequence of rescaling (7.9). The sequence v_k is uniformly bounded in C^0 since

$$\|v_k\|_0 \leq \Gamma^{-1}(\|\tilde{v}\|_0 + \|\tilde{v} - \tilde{v}_0\|_0) \lesssim \varepsilon_k^{1/2} \delta_1^{-1/2} (\delta_1^{1/2} + C\delta_1^{1/2} \varepsilon_k^{-1/2}) \lesssim \varepsilon_0^{1/2} + C.$$

Thus $(v_k \otimes v_k)$ is also uniformly bounded in C^0 . By Banach-Alaoglu v_k and $v_k \otimes v_k$ have weak- $*$ convergent subsequences.

Moreover, by rescaling (7.10) and using (7.2) we have

$$\|v_k - \bar{v}\|_{H^{-1}} \lesssim \|v_k - v\|_0 + \|v - \bar{v}\|_{H^{-1}} \lesssim \Gamma^{-1} \delta_1^{1/2} + C\lambda_0^{-1} \lesssim \varepsilon_k^{1/2} + C\lambda_0^{-1} \lesssim \varepsilon_k^{1/2} \quad (7.11)$$

by choosing a (and thus λ_0) sufficiently large in terms of ε_k . Moreover, from (7.4)–(7.6), (7.8), and (7.10) we obtain

$$\begin{aligned} \|v_k \otimes v_k - v \otimes v - \bar{R}\|_{H^{-1}} &\lesssim \|v_k \otimes v_k - v \otimes v\|_0 + \|R\|_0 + \|v \otimes v + R - \bar{v} \otimes \bar{v} - \bar{R}\|_{H^{-1}} \\ &\lesssim \Gamma^{-2} \|\tilde{v} \otimes \tilde{v} - \tilde{v}_0 \otimes \tilde{v}_0\|_0 + \left\| \dot{R} \right\|_0 + \|\text{tr } R\|_0 + C\lambda_0^{\alpha-1} \\ &\lesssim \varepsilon_k \delta_1^{-1/2} (C\delta_1^{1/2} \varepsilon_k^{-1/2} + \delta_1^{1/2}) + \varepsilon_k + C\lambda_0^{\alpha-1} \lesssim C\varepsilon_k^{1/2}. \end{aligned} \quad (7.12)$$

Since the H^{-1} topology uniquely captures the weak- $*$ limit, the theorem is completed upon passing $\varepsilon_k \rightarrow 0$ in (7.11)–(7.12). \square

APPENDIX A. HÖLDER SPACES

In the following $m = 0, 1, 2, \dots$, $\alpha \in (0, 1)$, and θ is a multi-index. We introduce the usual (spatial) Hölder norms as follows. First of all, the supremum norm is denoted by $\|f\|_0 := \sup_{\mathbb{T}^3 \times [0, T]} |f|$. We define the Hölder seminorms as

$$\begin{aligned} [f]_m &= \max_{|\theta|=m} \|D^\theta f\|_0, \\ [f]_{m+\alpha} &= \max_{|\theta|=m} \sup_{x \neq y, t} \frac{|D^\theta f(x, t) - D^\theta f(y, t)|}{|x - y|^\alpha}, \end{aligned}$$

where D^θ are *space derivatives* only. The Hölder norms are then given by

$$\begin{aligned} \|f\|_m &= \sum_{j=0}^m [f]_j \\ \|f\|_{m+\alpha} &= \|f\|_m + [f]_{m+\alpha}. \end{aligned}$$

Moreover, we will write $[f(t)]_\alpha$ and $\|f(t)\|_\alpha$ when the time t is fixed and the norms are computed for the restriction of f to the t -time slice.

Recall the following elementary inequalities:

$$[f]_s \leq C(\varepsilon^{r-s} [f]_r + \varepsilon^{-s} \|f\|_0) \quad (\text{A.1})$$

for $r \geq s \geq 0$, $\varepsilon > 0$, and

$$[fg]_r \leq C([f]_r \|g\|_0 + \|f\|_0 [g]_r) \quad (\text{A.2})$$

for $r \geq 0$. From (A.1) with $\varepsilon = \|f\|_0^{1/r} [f]_r^{-1/r}$ we obtain the standard interpolation inequalities

$$[f]_s \leq C \|f\|_0^{1-s/r} [f]_r^{s/r}. \quad (\text{A.3})$$

Next we collect two classical estimates on the Hölder norms of compositions. These are also standard, for instance in applications of the Nash-Moser iteration technique (for a detailed proof the reader might consult [DLS14, Proposition 4.1]).

Proposition A.1. *Let $\Psi : \Omega \rightarrow \mathbb{R}$ and $u : \mathbb{R}^n \rightarrow \Omega$ be two smooth functions, with $\Omega \subset \mathbb{R}^N$. Then, for every $m \in \mathbb{N} \setminus \{0\}$ there is a constant C (depending only on m, N and n) such that*

$$[\Psi \circ u]_m \leq C([\Psi]_1 \|Du\|_{m-1} + \|D\Psi\|_{m-1} \|u\|_0^{m-1} \|u\|_m) \quad (\text{A.4})$$

$$[\Psi \circ u]_m \leq C([\Psi]_1 \|Du\|_{m-1} + \|D\Psi\|_{m-1} [u]_1^m). \quad (\text{A.5})$$

We also recall the quadratic commutator estimate of [CET94] (cf. also [CDLS12, Lemma 1]):

Proposition A.2. *Let $f, g \in C^\infty(\mathbb{T}^3 \times \mathbb{T})$ and ψ a standard radial smooth and compactly supported kernel. For any $r \geq 0$ we have the estimate*

$$\left\| (f * \psi_\ell)(g * \psi_\ell) - (fg) * \psi_\ell \right\|_r \leq C\ell^{2-r} \|f\|_1 \|g\|_1,$$

where the constant C depends only on r .

APPENDIX B. ESTIMATES FOR TRANSPORT EQUATIONS

In this section we recall some well known results regarding smooth solutions of the *transport equation*:

$$\begin{cases} \partial_t f + v \cdot \nabla f = g, \\ f(\cdot, 0) = f_0, \end{cases} \quad (\text{B.1})$$

where $v = v(t, x)$ is a given smooth vector field. We will consider solutions on the entire space \mathbb{R}^3 and treat solutions on the torus simply as periodic solution in \mathbb{R}^3 . The following proposition contains standard estimates for such solutions (for a detailed proof, the reader might consult [BDLISJ15, Appendix D]).

Proposition B.1. *Assume $|t| \|v\|_1 \leq 1$. Then, any solution f of (B.1) satisfies*

$$\|f(t)\|_0 \leq \|f_0\|_0 + \int_0^t \|g(\cdot, \tau)\|_0 d\tau, \quad (\text{B.2})$$

$$\|f(t)\|_\alpha \leq e^\alpha \left(\|f_0\|_\alpha + \int_0^t \|g(\cdot, \tau)\|_\alpha d\tau \right), \quad (\text{B.3})$$

for all $0 \leq \alpha \leq 1$, and, more generally, for any $N \geq 1$ and $0 \leq \alpha < 1$

$$[f(t)]_{N+\alpha} \lesssim [f_0]_{N+\alpha} + |t| [v]_{N+\alpha} [f_0]_1 + \int_0^t \left([g(\tau)]_{N+\alpha} + (t-\tau) [v]_{N+\alpha} [g(\tau)]_1 \right) d\tau. \quad (\text{B.4})$$

Define $\Phi(\cdot, t)$ to be the inverse of the flux X of v starting at time 0 as the identity (i.e. $d/dt X = v(X, t)$ and $X(x, 0) = x$). Under the same assumptions as above we have:

$$\|\nabla \Phi(t) - \text{Id}\|_0 \lesssim |t| [v]_1, \quad (\text{B.5})$$

$$[\Phi(t)]_N \lesssim |t| [v]_N \quad \forall N \geq 2. \quad (\text{B.6})$$

APPENDIX C. POTENTIAL THEORY ESTIMATES

We recall the definition of the standard class of periodic Calderón-Zygmund operators. Let K be an \mathbb{R}^3 kernel which obeys the properties

- $K(z) = \Omega\left(\frac{z}{|z|}\right) |z|^{-3}$, for all $z \in \mathbb{R}^3 \setminus \{0\}$
- $\Omega \in C^\infty(\mathbb{S}^2)$
- $\int_{|\hat{z}|=1} \Omega(\hat{z}) d\hat{z} = 0$.

From the \mathbb{R}^3 kernel K , use Poisson summation to define the periodic kernel

$$K_{\mathbb{T}^3}(z) = K(z) + \sum_{\ell \in \mathbb{Z}^3 \setminus \{0\}} (K(z + \ell) - K(\ell)).$$

Then the operator

$$T_K f(x) = p.v. \int_{\mathbb{T}^3} K_{\mathbb{T}^3}(x - y) f(y) dy$$

is a \mathbb{T}^3 -periodic Calderón-Zygmund operator, acting on \mathbb{T}^3 -periodic functions f with zero mean on \mathbb{T}^3 . The following proposition, proving the boundedness of periodic Calderón-Zygmund operators on periodic Hölder spaces is classical (see e.g. [CZ54]):

Proposition C.1. *Fix $\alpha \in (0, 1)$. Periodic Calderón-Zygmund operators are bounded on the space of zero mean \mathbb{T}^3 -periodic C^α functions.*

The following is a simple consequence of classical stationary phase techniques. For a detailed proof the reader might consult [DS17, Lemma 2.2].

Proposition C.2. *Let $\alpha \in (0, 1)$ and $N \geq 1$. Let $a \in C^\infty(\mathbb{T}^3)$, $\Phi \in C^\infty(\mathbb{T}^3; \mathbb{R}^3)$ be smooth functions and assume that*

$$\hat{C}^{-1} \leq |\nabla \Phi| \leq \hat{C}$$

holds on \mathbb{T}^3 . Then

$$\left| \int_{\mathbb{T}^3} a(x) e^{ik \cdot \Phi} dx \right| \lesssim \frac{\|a\|_N + \|a\|_0 \|\Phi\|_N}{|k|^N}, \quad (\text{C.1})$$

and for the operator \mathcal{R} defined in (4.4), we have

$$\left\| \mathcal{R} \left(a(x) e^{ik \cdot \Phi} \right) \right\|_\alpha \lesssim \frac{\|a\|_0}{|k|^{1-\alpha}} + \frac{\|a\|_{N+\alpha} + \|a\|_0 \|\Phi\|_{N+\alpha}}{|k|^{N-\alpha}},$$

where the implicit constant depends on \hat{C} , α and N , but not on k .

APPENDIX D. COMMUTATORS INVOLVING SINGULAR INTEGRALS

The following lemma is a variant of Lemma 1 from [Con15]:

Proposition D.1. *Let $\alpha \in (0, 1)$ and $N \geq 0$. Let T_K be a Calderón-Zygmund operator with kernel K . Let $b \in C^{N+1, \alpha}(\mathbb{T}^3)$ a vectorfield. Then we have*

$$\| [T_K, b \cdot \nabla] f \|_{N+\alpha} \lesssim \|b\|_{1+\alpha} \|f\|_{N+\alpha} + \|b\|_{N+1+\alpha} \|f\|_\alpha$$

for any $f \in C^{N+\alpha}(\mathbb{T}^3)$, where the implicit constant depends on α, N and K .

Proof of Proposition D.1. The case $N = 0$ is precisely Lemma 1 in [Con15], except that in the former paper, the proof is given for Calderón-Zygmund operators defined on \mathbb{R}^3 , and for functions in $C^\alpha(\mathbb{R}^3) \cap L^p(\mathbb{R}^3)$. However, note that if f is the 1-periodic extension to all of \mathbb{R}^3 of the function f on \mathbb{T}^3 , and if $\chi(y)$ is a smooth cutoff function, which is identically 1 on $[-1 - 1/20, 1 + 1/20]^3$, and vanishes on the complement of $[-1 - 1/10, 1 + 1/10]^3$, we then have that

$$T_K f(x) = p.v. \int_{\mathbb{R}^3} K(x - y) \chi(y) f(y) dy + T_{\text{smooth}} f(x)$$

where

$$T_{\text{smooth}}: C^0(\mathbb{T}^3) \rightarrow C^N(\mathbb{T}^3)$$

is a bounded operator, for any $N \geq 0$. Thus, modulo using the smoothing property of T_{smooth} , we may apply directly the proof in [Con15] to the periodic case of this paper.

Let us now consider the case $N \geq 1$, and to this end let θ be a multi-index with $|\theta| = N$. Then, by the Leibniz rule

$$\begin{aligned} \partial^\theta [T_K, b \cdot \nabla] f &= T_K(\partial^\theta (b \cdot \nabla f)) - \partial^\theta (b \cdot \nabla T_K f) \\ &= \sum_{\theta'} \binom{\theta}{\theta'} \left\{ T_K(\partial^{\theta'} b \cdot \nabla \partial^{\theta - \theta'} f) - \partial^{\theta'} b \cdot \nabla \partial^{\theta - \theta'} T_K f \right\} \\ &= \sum_{\theta'} \binom{\theta}{\theta'} \left\{ [T_K, \partial^{\theta'} b \cdot \nabla] \partial^{\theta - \theta'} f \right\}. \end{aligned}$$

Therefore we obtain from the case $N = 0$:

$$\left\| \partial^\theta [T_K, b \cdot \nabla] f \right\|_\alpha \lesssim \sum_{j=0}^N \|b\|_{j+1+\alpha} \|f\|_{N-j+\alpha}.$$

Furthermore, by interpolation

$$\|b\|_{j+1+\alpha} \lesssim \|b\|_{1+\alpha}^{1-j/N} \|b\|_{N+1+\alpha}^{j/N}, \quad \text{and} \quad \|f\|_{N-j+\alpha} \lesssim \|f\|_{N+\alpha}^{1-j/N} \|f\|_\alpha^{j/N},$$

so that, for any $j = 0, \dots, N$

$$\|b\|_{j+1+\alpha} \|f\|_{N-j+\alpha} \lesssim \|b\|_{1+\alpha} \|f\|_{N+\alpha} + \|b\|_{N+1+\alpha} \|f\|_\alpha.$$

This concludes the proof. \square

REFERENCES

- [BDLS11] Y. Brenier, C. De Lellis, and L. Székelyhidi, Jr. Weak-strong uniqueness for measure-valued solutions. *Comm. Math. Phys.*, 305(2):351–361, 2011.
- [Buc14] T. Buckmaster. *Onsager’s Conjecture*. PhD thesis, Universität Leipzig, 2014.
- [Buc15] T. Buckmaster. Onsager’s conjecture almost everywhere in time. *Comm. Math. Phys.*, 333(3):1175–1198, 2015.
- [BDLISJ15] T. Buckmaster, C. De Lellis, P. Isett, and L. Székelyhidi Jr. Anomalous dissipation for 1/5-holder Euler flows. *Annals of Mathematics*, 182(1):127–172, 2015.
- [BDLS13] T. Buckmaster, C. De Lellis, and L. Székelyhidi, Jr. Transporting microstructure and dissipative Euler flows. *arXiv preprint arXiv:1302.2815*, 2013.
- [BDLS16] T. Buckmaster, C. De Lellis, and L. Székelyhidi, Jr. Dissipative Euler flows with Onsager-critical spatial regularity. *Comm. Pure Appl. Math.*, 69(9):1613–1670, 2016.
- [BSV16] T. Buckmaster, S. Shkoller, and V. Vicol. Nonuniqueness of weak solutions to the SQG equation. *arXiv preprint arXiv:1610.00676*, 2016.
- [CZ54] A.P. Calderón and A. Zygmund. Singular integrals and periodic functions. *Studia Math.*, 14:249–271, 1954.
- [CCF16] A. Castro, D. Córdoba, and D. Faraco. Mixing solutions for the Muskat problem. *arXiv preprint arXiv:1605.04822*, 2016.
- [CCFS08] A. Cheskidov, P. Constantin, S. Friedlander, and R. Shvydkoy. Energy conservation and Onsager’s conjecture for the Euler equations. *Nonlinearity*, 21(6):1233–1252, 2008.
- [Chi14] E. Chiodaroli. A counterexample to well-posedness of entropy solutions to the compressible Euler system. *J. Hyper. Differential Equations* 11(3):493–519, 2014.
- [CS14] A. Choffrut and L. Székelyhidi Jr. Weak solutions to the stationary incompressible Euler equations. *SIAM J. Math. Anal.*, 46(6):4060–4074, 2014.
- [Con15] P. Constantin. Lagrangian–Eulerian methods for uniqueness in hydrodynamic systems. *Adv. Math.*, 278:67–102, 2015.
- [CET94] P. Constantin, W. E, and E.S. Titi. Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Comm. Math. Phys.*, 165(1):207–209, 1994.
- [CDLS12] S. Conti, C. De Lellis, and L. Székelyhidi Jr. h -principle and rigidity for $C^{1,\alpha}$ isometric embeddings. In *Nonlinear partial differential equations*, pages 83–116. Springer, 2012.
- [CFG11] D. Córdoba, D. Faraco, and F. Gancedo. Lack of uniqueness for weak solutions of the incompressible porous media equation. *Arch. Ration. Mech. Anal.*, 200(3):725–746, 2011.

- [DS17] S. Daneri and L. Székelyhidi. Non-uniqueness and h-principle for hölder-continuous weak solutions of the euler equations. *Archive for Rational Mechanics and Analysis*, 224(2):471–514, 2017.
- [DLS09] C. De Lellis and L. Székelyhidi, Jr. The Euler equations as a differential inclusion. *Ann. of Math. (2)*, 170(3):1417–1436, 2009.
- [DL10] C. De Lellis and L. Székelyhidi Jr. On admissibility criteria for weak solutions of the Euler equations. *Arch. Ration. Mech. Anal.*, 195(1):225–260, 2010.
- [DLS12] C. De Lellis and L. Székelyhidi, Jr. The h -principle and the equations of fluid dynamics. *Bull. Amer. Math. Soc. (N.S.)*, 49(3):347–375, 2012.
- [DLS13] C. De Lellis and L. Székelyhidi, Jr. Dissipative continuous Euler flows. *Invent. Math.*, 193(2):377–407, 2013.
- [DLS14] C. De Lellis and L. Székelyhidi, Jr. Dissipative Euler flows and Onsager’s conjecture. *J. Eur. Math. Soc. (JEMS)*, 16(7):1467–1505, 2014.
- [Eyi94] G. L. Eyink. Energy dissipation without viscosity in ideal hydrodynamics. I. Fourier analysis and local energy transfer. *Phys. D*, 78(3-4):222–240, 1994.
- [Ise13a] P. Isett. *Holder continuous Euler flows with compact support in time*. ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)–Princeton University.
- [Ise13b] P. Isett. Regularity in time along the coarse scale flow for the incompressible Euler equations. *arXiv preprint*, 1307.0565, 2013.
- [Ise16] P. Isett. A proof of Onsager’s conjecture. *arXiv preprint* arXiv:1608.08301, 2016.
- [IO16] P. Isett and S.-J. Oh. On nonperiodic Euler flows with Hölder regularity. *Arch. Ration. Mech. Anal.*, 221(2):725–804, 2016.
- [IV15] P. Isett and V. Vicol. Hölder continuous solutions of active scalar equations. *Annals of PDE*, 1(1):1–77, 2015.
- [Kui55] N. H. Kuiper. On C^1 -isometric imbeddings. I, II. *Nederl. Akad. Wetensch. Proc. Ser. A*. **58** = *Indag. Math.*, 17:545–556, 683–689, 1955.
- [Lio96] P.-L. Lions. *Mathematical Topics in Fluid Mechanics: Volume 1: Incompressible Models*. Oxford University Press, June 1996.
- [LXX16] T. Luo, C. Xie, Z. Xin. Non-uniqueness of admissible weak solutions to compressible Euler systems with source terms. *Advances in Math.*, 291:542–583, 2016.
- [MB02] A.J. Majda and A.L. Bertozzi. *Vorticity and incompressible flow*, volume 27 of *Cambridge Texts in Applied Mathematics*. Cambridge University Press, Cambridge, 2002.
- [Nas54] J. Nash. C^1 isometric imbeddings. *Ann. of Math. (2)*, 60:383–396, 1954.
- [Ons49] L. Onsager. Statistical hydrodynamics. *Il Nuovo Cimento (1943-1954)*, 6:279–287, 1949.
- [Sch93] V. Scheffer. An inviscid flow with compact support in space-time. *J. Geom. Anal.*, 3(4):343–401, 1993.
- [Shn97] A. Shnirelman. On the nonuniqueness of weak solution of the Euler equation. *Comm. Pure Appl. Math.*, 50(12):1261–1286, 1997.
- [Shv11] R. Shvydkoy. Convex integration for a class of active scalar equations. *J. Amer. Math. Soc.*, 24(4):1159–1174, 2011.
- [Szé12] L. Székelyhidi Jr. Relaxation of the incompressible porous media equation. *Ann. Sci. Éc. Norm. Supér. (4)*, 45(3):491–509, 2012.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NY 10012, USA

E-mail address: buckmaster@cims.nyu.edu

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, CH-8057 ZÜRICH, SWITZERLAND

E-mail address: camillo.delellis@math.unizh.ch

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT LEIPZIG, D-04103 LEIPZIG, GERMANY

E-mail address: laszlo.szekelyhidi@math.uni-leipzig.de

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544, USA

E-mail address: vvicol@math.princeton.edu