

An exact sequence for $K_*^M/2$ with applications to quadratic forms

By D. ORLOV,^{*} A. VISHIK,^{**} and V. VOEVODSKY^{***}

Contents

1. Introduction
2. An exact sequence for $K_*^M/2$
3. Reduction to points of degree 2
4. Some applications
 - 4.1. Milnor's Conjecture on quadratic forms
 - 4.2. The Kahn-Rost-Sujatha Conjecture
 - 4.3. The J -filtration conjecture

1. Introduction

Let k be a field of characteristics zero. For a sequence $\underline{a} = (a_1, \dots, a_n)$ of invertible elements of k consider the homomorphism

$$K_*^M(k)/2 \rightarrow K_{*+n}^M(k)/2$$

in Milnor's K -theory modulo elements divisible by 2 defined by multiplication with the symbol corresponding to \underline{a} . The goal of this paper is to construct a four-term exact sequence (18) which provides information about the kernel and cokernel of this homomorphism.

The proof of our main theorem (Theorem 3.2) consists of two independent parts. Let $Q_{\underline{a}}$ be the norm quadric defined by the sequence \underline{a} (see below). First, we use the techniques of [13] to establish a four term exact sequence (1) relating the kernel and cokernel of multiplication by \underline{a} with Milnor's K -theory of the closed and the generic points of $Q_{\underline{a}}$ respectively. This is done in the first section. Then, using elementary geometric arguments, we show that the sequence can be rewritten in its final form (18) which involves only the generic point and the closed points with residue fields of degree 2.

^{*}Supported by NSF grant DMS-97-29992.

^{**}Supported by NSF grant DMS-97-29992 and RFFI-99-01-01144.

^{***}Supported by NSF grants DMS-97-29992 and DMS-9901219 and the Ambrose Monell Foundation.

As an application we establish, for fields of characteristics zero, the validity of three conjectures in the theory of quadratic forms - the Milnor conjecture on the structure of the Witt ring, the Khan-Rost-Sujatha conjecture and the J -filtration conjecture. All these results require only the first form of our exact sequence. Using the final form of the sequence we also show that the kernel of multiplication by \underline{a} is generated, as a $K_*^M(k)$ -module, by its components of degree ≤ 1 .

This paper is a natural extension of [13] and we feel free to refer to the results of [13] without reproducing them here. Most of the mathematics used in this paper was developed in the spring of 1995 when all three authors were at Harvard. In its present form the paper was written while the authors were members of the Institute for Advanced Study in Princeton. We would like to thank both institutions for their support.

2. An exact sequence for $K_*^M/2$

Let $\underline{a} = (a_1, \dots, a_n)$ be a sequence of elements of k^* . Recall that the n -fold Pfister form $\langle\langle a_1, \dots, a_n \rangle\rangle$ is defined as the tensor product

$$\langle 1, -a_1 \rangle \otimes \cdots \otimes \langle 1, -a_n \rangle$$

where $\langle 1, -a_i \rangle$ is the norm form in the quadratic extension $k(\sqrt{a_i})$. Denote by $Q_{\underline{a}}$ the projective quadric of dimension $2^{n-1} - 1$ defined by the form $q_{\underline{a}} = \langle\langle a_1, \dots, a_{n-1} \rangle\rangle - \langle a_n \rangle$. This quadric is called the small Pfister quadric or the norm quadric associated with the symbol \underline{a} . Denote by $k(Q_{\underline{a}})$ the function field of $Q_{\underline{a}}$ and by $(Q_{\underline{a}})_0$ the set of closed points of $Q_{\underline{a}}$. The following result is the main theorem of the paper.

THEOREM 2.1. *Let k be a field of characteristic zero. Then for any sequence of invertible elements (a_1, \dots, a_n) the following sequence of abelian groups is exact*

$$(1) \quad \coprod_{x \in (Q_{\underline{a}})_0} K_*^M(k(x))/2 \xrightarrow{\text{Tr}_{k(x)/k}} K_*^M(k)/2 \xrightarrow{\cdot \underline{a}} K_{*+n}^M(k)/2 \rightarrow K_{*+n}^M(k(Q_{\underline{a}}))/2.$$

The proof goes as follows. We first construct two exact sequences of the form

$$(2) \quad 0 \rightarrow K \rightarrow K_{*+n}^M(k)/2 \rightarrow K_{*+n}^M(k(Q_{\underline{a}}))/2$$

and

$$(3) \quad \coprod_{x \in (Q_{\underline{a}})_0} K_*^M(k(x))/2 \xrightarrow{\text{Tr}_{k(x)/k}} K_*^M(k)/2 \rightarrow I \rightarrow 0$$

and then construct an isomorphism $I \rightarrow K$ such that the composition

$$K_*^M(k)/2 \rightarrow I \rightarrow K \rightarrow K_{*+n}^M(k)/2$$

is multiplication by \underline{a} .

Our construction of the sequence (2) makes sense for any smooth scheme X and we shall do it in this generality. Recall that we denote by $\check{C}(X)$ the simplicial scheme such that $\check{C}(X)_n = X^{n+1}$ and that faces and degeneracy morphisms are given by partial projections and diagonal embeddings respectively. We will use repeatedly the following lemma which is an immediate corollary of [13, Lemma 7.2] and [13, Cor. 6.7].

LEMMA 2.2. *For any smooth scheme X over k and any $p \leq q$ the homomorphism*

$$H^{p,q}(\mathrm{Spec}(k), \mathbf{Z}/2) \rightarrow H^{p,q}(\check{C}(X), \mathbf{Z}/2)$$

defined by the canonical morphism $\check{C}(X) \rightarrow \mathrm{Spec}(k)$, is an isomorphism.

PROPOSITION 2.3. *For any $n \geq 0$ there is an exact sequence of the form*

$$(4) \quad 0 \rightarrow H^{n,n-1}(\check{C}(X), \mathbf{Z}/2) \rightarrow K_n^M(k)/2 \rightarrow K_n^M(k(X))/2.$$

Proof. The computation of motivic cohomology of weight 1 shows that

$$\mathrm{Hom}(\mathbf{Z}/2, \mathbf{Z}/2(1)) \cong H^{0,1}(\mathrm{Spec}(k), \mathbf{Z}/2) \cong \mathbf{Z}/2.$$

The nontrivial element $\tau : \mathbf{Z}/2 \rightarrow \mathbf{Z}/2(1)$ together with multiplication morphism $\mathbf{Z}(n-1) \otimes \mathbf{Z}/2(1) \xrightarrow{\sim} \mathbf{Z}/2(n)$ defines a morphism

$$\tau : \mathbf{Z}/2(n-1) \rightarrow \mathbf{Z}/2(n).$$

The Beilinson-Lichtenbaum conjecture implies immediately the following result.

LEMMA 2.4. *The morphism τ extends to a distinguished triangle in DM_-^{eff} of the form*

$$(5) \quad \mathbf{Z}/2(n-1) \xrightarrow{\tau} \mathbf{Z}/2(n) \rightarrow \underline{H}^{n,n}(\mathbf{Z}/2)[-n],$$

where $\underline{H}^n(\mathbf{Z}/2(n))$ is the n^{th} cohomology sheaf of the complex $\mathbf{Z}/2(n)$.

Consider the long sequence of morphisms in the triangulated category of motives from the motive of $\check{C}(X)$ to the distinguished triangle (5). It starts as

$$0 \rightarrow H^{n,n-1}(\check{C}(X), \mathbf{Z}/2) \rightarrow H^{n,n}(\check{C}(X), \mathbf{Z}/2) \rightarrow H^0(\check{C}(X), \underline{H}^{n,n}(\mathbf{Z}/2)).$$

By Lemma 2.2 there are isomorphisms

$$H^n(\check{C}(X), \mathbf{Z}/2(n)) = H^{n,n}(\mathrm{Spec}(k), \mathbf{Z}/2) = K_n^M(k)/2.$$

On the other hand, since $\underline{H}^{n,n}(\mathbf{Z}/2)$ is a homotopy invariant sheaf with transfers, we have an embedding

$$H^0(\check{C}(X), \underline{H}^{n,n}(\mathbf{Z}/2)) \hookrightarrow \underline{H}^{n,n}(\mathbf{Z}/2)(\mathrm{Spec}(k(X))).$$

The right-hand side is isomorphic to $H^{n,n}(\mathrm{Spec}(k(X)), \mathbf{Z}/2) = K_n^M(k(X))/2$. This completes the proof of the proposition. \square

Let us now construct the exact sequence (3). Denote the standard simplicial scheme $\check{C}(Q_{\underline{a}})$ by $\mathcal{X}_{\underline{a}}$. Recall that we have a distinguished triangle of the form

$$(6) \quad M(\mathcal{X}_{\underline{a}})(2^{n-1} - 1)[2^n - 2] \xrightarrow{\varphi} M_{\underline{a}} \xrightarrow{\psi} M(\mathcal{X}_{\underline{a}}) \xrightarrow{\mu'} M(\mathcal{X}_{\underline{a}})(2^{n-1} - 1)[2^n - 1]$$

where $M_{\underline{a}}$ is a direct summand of the motive of the quadric $Q_{\underline{a}}$. Denote the composition

$$(7) \quad M(\mathcal{X}_{\underline{a}}) \xrightarrow{\mu'} M(\mathcal{X}_{\underline{a}})(2^{n-1} - 1)[2^n - 1] \xrightarrow{pr} \mathbf{Z}/2(2^{n-1} - 1)[2^n - 1]$$

by $\mu \in H^{2^n-1, 2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$. By Lemma 2.2,

$$H^{i,i}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) = H^{i,i}(\mathrm{Spec}(k), \mathbf{Z}/2) = K_i^M(k)/2.$$

Therefore, multiplication with μ defines a homomorphism

$$K_i^M(k)/2 \xrightarrow{\cdot\mu} H^{i+2^n-1, i+2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2).$$

PROPOSITION 2.5. *The sequence*

$$(8) \quad \coprod_{x \in (Q_{\underline{a}})_{(0)}} K_i^M(k(x))/2 \xrightarrow{\mathrm{Tr}_{k(x)/k}} K_i^M(k)/2 \xrightarrow{\cdot\mu} H^{i+2^n-1, i+2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \rightarrow 0$$

is exact.

Proof. Let us consider morphisms in the triangulated category of motives from the distinguished triangle (6) to the object $\mathbf{Z}/2(i+2^{n-1}-1)[i+2^n-1]$. By definition, $M_{\underline{a}}$ is a direct summand of the motive of the smooth projective variety $Q_{\underline{a}}$ of dimension $2^{n-1}-1$. Therefore, the group $H^{i+2^n-1, i+2^{n-1}-1}(M_{\underline{a}}, \mathbf{Z}/2)$ is trivial by [13, Lemma 4.11] and [9]. Using this fact, we obtain the following exact sequence:

$$(9) \quad H^{i+2^n-2, i+2^{n-1}-1}(M_{\underline{a}}, \mathbf{Z}/2) \xrightarrow{\varphi^*} H^{i,i}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \xrightarrow{\mu'^*} \\ \rightarrow H^{i+2^n-1, i+2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \rightarrow 0.$$

By definition (see [13, p. 22]) the morphism φ is given by the composition

$$(10) \quad M(\mathcal{X}_{\underline{a}})(2^{n-1} - 1)[2^n - 2] \xrightarrow{pr} \mathbf{Z}(2^{n-1} - 1)[2^n - 2] \rightarrow M_{\underline{a}}$$

and the composition of the second arrow with the canonical embedding $M_{\underline{a}} \rightarrow M(Q_{\underline{a}})$ is the fundamental cycle map

$$\mathbf{Z}(2^{n-1} - 1)[2^n - 2] \rightarrow M(Q_{\underline{a}})$$

which corresponds to the fundamental cycle on $Q_{\underline{a}}$ under the isomorphism

$$\mathrm{Hom}(\mathbf{Z}(2^{n-1} - 1)[2^n - 2], M(Q_{\underline{a}})) = \mathrm{CH}_{2^{n-1}-1}(Q_{\underline{a}}) \cong \mathbf{Z}$$

(see [13, Th. 4.4]). On the other hand by Lemma 2.2 the homomorphism

$$H^{i,i}(\mathrm{Spec}(k), \mathbf{Z}/2) \rightarrow H^{i,i}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$$

defined by the first arrow in (10) is an isomorphism. This implies immediately that the exact sequence (9) defines an exact sequence of the form

$$(11) \quad H^{i+2^n-2, i+2^{n-1}-1}(Q_{\underline{a}}, \mathbf{Z}/2) \xrightarrow{\varphi^*} H^{i,i}(\mathrm{Spec}(k), \mathbf{Z}/2) \xrightarrow{\mu'^*} \\ \rightarrow H^{i+2^n-1, i+2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \rightarrow 0.$$

By [13, Lemma 4.11] there is an isomorphism

$$H^{i+2^n-2, i+2^{n-1}-1}(Q_{\underline{a}}, \mathbf{Z}/2) \cong H^{2^{n-1}-1}(Q_{\underline{a}}, \underline{K}_{i+2^{n-1}-1}^M/2).$$

The Gersten resolution for the sheaf $\underline{K}_m^M/2$ (see, for example, [9]) shows that the group $H^{2^{n-1}-1}(Q_{\underline{a}}, \underline{K}_{i+2^{n-1}-1}^M/2)$ can be identified with the cokernel of the map:

$$\coprod_{y \in (Q_{\underline{a}})_{(1)}} K_{i+1}^M(k(y))/2 \xrightarrow{\partial} \coprod_{x \in (Q_{\underline{a}})_{(0)}} K_i^M(k(x))/2,$$

and the map $H^{i+2^n-2, i+2^{n-1}-1}(Q_{\underline{a}}, \mathbf{Z}/2) \rightarrow H^{i,i}(\mathrm{Spec}(k), \mathbf{Z}/2)$ defined by the fundamental cycle corresponds in this description to the map

$$\coprod_{x \in (Q_{\underline{a}})_{(0)}} K_i^M(k(x))/2 \xrightarrow{\mathrm{Tr}_{k(x)/k}} K_i^M(k)/2 = H^{i,i}(\mathrm{Spec}(k), \mathbf{Z}/2).$$

This finishes the proof of Proposition 2.5. \square

We are going to show now that the map $K_*^M(k)/2 \xrightarrow{\alpha} K_{*+n}^M(k)/2$ glues the exact sequences (4) and (8) in one. Denote by $\mathbb{H}^i(\mathcal{X}_{\underline{a}})$ the direct sum $\oplus_m H^{m+i, m}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$. It has a natural structure of a graded module over the ring $K_*^M(k)/2$ and one can easily see that the sequences (4) and (8) define sequences of $K_*^M(k)/2$ -modules of the form

$$(12) \quad 0 \rightarrow \mathbb{H}^1(\mathcal{X}_{\underline{a}}) \rightarrow K_*^M(k)/2 \rightarrow K_*^M(k(Q_{\underline{a}}))/2,$$

$$(13) \quad \coprod_{x \in (Q_{\underline{a}})_{(0)}} K_*^M(k(x))/2 \xrightarrow{\mathrm{Tr}_{k(x)/k}} K_*^M(k)/2 \xrightarrow{\mu} \mathbb{H}^{2^{n-1}}(\mathcal{X}_{\underline{a}}) \rightarrow 0.$$

Consider cohomological operations

$$Q_i : H^{\bullet, *}(-, \mathbf{Z}/2) \rightarrow H^{\bullet+2^{i+1}-1, *+2^i-1}(-, \mathbf{Z}/2)$$

introduced in [12]. The composition $Q_{n-2} \cdots Q_0$ defines a homomorphism of graded abelian groups $d : \mathbb{H}^1(\mathcal{X}_{\underline{a}}) \rightarrow \mathbb{H}^{2^{n-1}}(\mathcal{X}_{\underline{a}})$. Now, [12, Prop. 13.4] together with the fact that $H^{p,q}(\mathrm{Spec}(k), \mathbf{Z}/2) = 0$ for $p > q$ implies that d is a homomorphism of $K_*^M(k)/2$ -modules. We are going to show that d is an isomorphism and that the composition

$$(14) \quad K_*^M(k)/2 \xrightarrow{\mu} \mathbb{H}^{2^{n-1}}(\mathcal{X}_{\underline{a}}) \xrightarrow{d^{-1}} \mathbb{H}^1(\mathcal{X}_{\underline{a}}) \rightarrow K_*^M(k)/2$$

is multiplication with \underline{a} .

LEMMA 2.6. *The homomorphism d is injective.*

Proof. We have to show that the composition of operations

$$Q_{n-2} \dots Q_0 : H^{*+n, *+n-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \rightarrow H^{*+2^{n-1}, *+2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$$

is injective. Let $\tilde{\mathcal{X}}_{\underline{a}}$ be the simplicial cone of the morphism $\mathcal{X}_{\underline{a}} \rightarrow \mathrm{Spec}(k)$ which we consider as a pointed simplicial scheme. The long exact sequence of cohomology defined by the cofibration sequence

$$(15) \quad (\mathcal{X}_{\underline{a}})_+ \rightarrow \mathrm{Spec}(k)_+ \rightarrow \tilde{\mathcal{X}}_{\underline{a}} \rightarrow \Sigma_s^1((\mathcal{X}_{\underline{a}})_+)$$

together with the fact that $H^{p,q}(\mathrm{Spec}(k), \mathbf{Z}/2) = 0$ for $p > q$ shows that for $p > q + 1$ we have a natural isomorphism $H^{p,q}(\tilde{\mathcal{X}}_{\underline{a}}, \mathbf{Z}/2) = H^{p-1,q}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$ compatible with the actions of cohomological operations. Therefore, it is sufficient to prove injectivity of the composition $Q_{n-2} \dots Q_0$ on motivic cohomology groups of the form $H^{*+n+1, *+n-1}(\tilde{\mathcal{X}}_{\underline{a}}, \mathbf{Z}/2)$. To show that $Q_{n-2} \dots Q_0$ is a monomorphism it is sufficient to check that the operation Q_i acts monomorphically on the group

$$H^{*+n-i+2^{i+1}-1, *+n-i+2^i-2}(\tilde{\mathcal{X}}_{\underline{a}}, \mathbf{Z}/2)$$

for all $i = 0, \dots, n-2$. For any $i \leq n-1$ we have $\ker(Q_i) = \mathrm{Im}(Q_i)$ by [13, Cor. 3.5]. Therefore, the kernel of Q_i on our group is the image of $H^{*+n-i, *+n-i-1}(\tilde{\mathcal{X}}_{\underline{a}}, \mathbf{Z}/2)$. On the other hand, the cofibration sequence (15) together with Lemma 2.2 implies that for $p \leq q+1$ we have $H^{p,q}(\tilde{\mathcal{X}}_{\underline{a}}, \mathbf{Z}/2) = 0$ which proves the lemma. \square

Denote by γ the element of $H^{n, n-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$ which corresponds to the symbol \underline{a} under the embedding into $K_n^M(k)/2$ (sequence (4)). To prove that d is surjective and that the composition (14) is multiplication with \underline{a} we use the following lemma.

LEMMA 2.7. *The composition $K_*^M(k)/2 \xrightarrow{\gamma} \mathbb{H}^1(\mathcal{X}_{\underline{a}}) \xrightarrow{d} \mathbb{H}^{2^{n-1}}(\mathcal{X}_{\underline{a}})$ coincides with multiplication by μ .*

Proof. Since our maps are homomorphisms of $K_*^M(k)$ -modules it is sufficient to verify that the cohomological operation d sends $\gamma \in H^{n, n-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$ to $\mu \in H^{2^{n-1}, 2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2)$. By Lemma 2.6, d is injective. Therefore, the element $d(\gamma)$ is nonzero. On the other hand, sequence (8) shows that

$$H^{2^{n-1}, 2^{n-1}-1}(\mathcal{X}_{\underline{a}}, \mathbf{Z}/2) \cong K_0^M(k)/2 \cong \mathbf{Z}/2$$

and μ is a generator of this group. Therefore, $d(\gamma) = \mu$. \square

LEMMA 2.8. *The homomorphism d is surjective.*

Proof. This follows immediately from Lemma 2.7 and surjectivity of multiplication by μ (Proposition 2.5). \square

LEMMA 2.9. *The composition (14) is multiplication with \underline{a} .*

Proof. Since all the maps in (14) are morphisms of $K_*^M(k)$ -modules, it is sufficient to check the condition for the generator $1 \in K_0^M(k)/2$. And the later follows from Lemma 2.7 and the definition of γ . \square

This finishes the proof of Theorem 2.1.

The following statement, which is easily deduced from the exact sequence (1), is the key to many applications.

Let E/k be a field. For any element $h \in K_n^M(k)$ denote by $h|_E$, as usual, the restriction of h on E , i.e., the image of h under the natural morphism $K_n^M(k) \rightarrow K_n^M(E)$.

THEOREM 2.10. *For any field k and any nonzero $h \in K_n^M(k)/2$ there exist a field E/k and a pure symbol $\alpha = \{a_1, \dots, a_n\} \in K_n^M(k)/2$ such that $h|_E = \alpha|_E$ is a nonzero pure symbol of $K_n^M(E)/2$.*

Proof. Let $h = \alpha_1 + \dots + \alpha_l$, where α_i are pure symbols corresponding to sequences $\underline{a}_i = (a_{1i}, \dots, a_{ni})$. Let $Q_{\underline{a}_i}$ be the norm quadric corresponding to the symbol α_i . For any $0 < i \leq l$ denote by E_i the field $k(Q_{\underline{a}_1} \times \dots \times Q_{\underline{a}_i})$. It is clear that $h|_{E_i} = 0$. Let us fix i such that $h|_{E_{i+1}} = 0$ and $h|_{E_i}$ is a nonzero element. Then $h|_{E_i}$ belongs to

$$\ker(K_n^M(E_i)/2 \rightarrow K_n^M(E_{i+1})/2).$$

By Theorem 2.1, the kernel is covered by $K_0^M(E_i) \cong \mathbf{Z}/2$ and is generated by $\alpha_{i+1}|_{E_i}$. Thus, we have $\alpha_{i+1}|_{E_i} = h|_{E_i} \neq 0$. \square

3. Reduction to points of degree 2

In this section we prove the following result.

THEOREM 3.1. *Let k be a field such that $\text{char}(k) \neq 2$ and Q be a smooth quadric over k . Let $Q_{(0)}$ be the set of closed points of Q and $Q_{(0, \leq 2)}$ the subset in $Q_{(0)}$ of points x such that $[k_x : k] \leq 2$. Then, for any $n \geq 0$, the image of the map*

$$(16) \quad \oplus \text{tr}_{k_x/k} : \oplus_{x \in Q_{(0)}} K_n^M(k_x) \rightarrow K_n^M(k)$$

coincides with the image of the map

$$(17) \quad \oplus \text{tr}_{k_x/k} : \oplus_{x \in Q_{(0, \leq 2)}} K_n^M(k_x) \rightarrow K_n^M(k).$$

Combining Theorem 2.1 with Theorem 3.1 we get the following result.

THEOREM 3.2. *Let k be a field of characteristic zero and $\underline{a} = (a_1, \dots, a_n)$ a sequence of invertible elements of k . Then the sequence*

$$(18) \quad \bigoplus_{x \in (Q_{\underline{a}})_{(0, \leq 2)}} K_i^M(k_x)/2 \rightarrow K_i^M(k)/2 \xrightarrow{\alpha} K_{i+n}^M(k)/2 \rightarrow K_{i+n}^M(k(Q_{\underline{a}}))/2$$

is exact.

Theorem 3.2 together with the well known result of Bass and Tate (see [1, Cor. 5.3]) implies the following.

THEOREM 3.3. *Let k be a field of characteristic zero and $\underline{a} = (a_1, \dots, a_n)$ a sequence of invertible elements of k such that the corresponding elements of $K_n^M(k)/2$ are not zero. Then the kernel of the homomorphism $K_n^M(k)/2 \xrightarrow{\alpha} K_{*+n}^M(k)/2$ is generated, as a module over $K_*^M(k)$, by the kernel of the homomorphism $K_1^M(k)/2 \rightarrow K_{1+n}^M(k)/2$.*

Let us start the proof of Theorem 3.1 with the following two lemmas.

LEMMA 3.4. *Let E be an extension of k of degree n and V a k -linear subspace in E such that $2\dim(V) > n$. Then, for any $n > 0$, $K_n^M(E)$ is generated, as an abelian group, by elements of the form (x_1, \dots, x_n) where all x_i 's are in V .*

Proof. It is sufficient to prove the statement for $n = 1$. Let x be an invertible element of E . Since $2\dim(V) > \dim_k E$ we have $V \cap xV \neq 0$. Therefore x is a quotient of two elements of $V \cap E^*$. \square

LEMMA 3.5. *Let k be an infinite field and p a closed, separable point in \mathbf{P}_k^n , $n \geq 2$ of degree m . Then there exists a rational curve C of degree $m - 1$ such that $p \in C$ and C is either nonsingular, or has one rational singular point.*

Proof. We may assume that p lies in $\mathbf{A}^n \subset \mathbf{P}^n$. Then there exists a linear function x_1 on \mathbf{A}^n such that the map of the residue fields $k_{x_1(p)} \rightarrow k_p$ is an isomorphism. Let (x_1, \dots, x_n) be a coordinate system starting with x_1 . Since the restriction of x_1 to p is an isomorphism the inverse gives a collection of regular functions $\bar{x}_2, \dots, \bar{x}_n$ on $x_1(p) \subset \mathbf{A}^1$. Each of these functions has a representative f_i in $k[x_1]$ of degree at most $m - 1$. The projective closure of the affine curve given by the equations $x_i = f_i(x_1)$, $i = 2, \dots, n$, satisfies the conditions of the lemma. \square

Let Q be any quadric over k . If Q has a rational point (or even a point of odd degree, which is the same by Springer's theorem, [3, VII, Th. 2.3]), then Theorem 3.1 for Q holds for obvious reasons. Therefore we may assume that Q has no points of odd degree. It is well known (see e.g. [11, Th. 2.3.8, p. 39]) that any smooth quadric of dimension > 0 over a finite field of odd characteristic has a rational point. Since the statement of the theorem is

obvious for $\dim(Q) = 0$ we may assume that k is infinite. By the theorem of Springer, for finite extension of odd degree E/F , the quadric Q_F is isotropic if and only if Q_E is. Hence, we can assume that E/k is separable.

Let e be a point on Q with the residue field E . We have to show that the image of the transfer map $K_n^M(E) \rightarrow K_n^M(k)$ lies in the image of the map (17). We proceed by induction on d where $2d = [E : k]$. If $d = 1$ there is nothing to prove. Assume by induction that for any closed point f of Q such that $[k_f : k] < 2d$ the image of the transfer map $K_n^M(k_f) \rightarrow K_n^M(k)$ lies in the image of (17).

If $\dim(Q) = 0$ our statement is obvious. Consider the case of a conic $\dim(Q) = 1$. Let D be any effective divisor on Q of degree $2d - 2$. Denote by $h^0(D)$ the linear space $H^0(Q, \mathcal{O}(D))$ which can be identified with the space of rational functions f such that $D + (f)$ is effective. Evaluating elements of $h^0(D)$ on e we get a homomorphism $h^0(D) \rightarrow E$ which is injective since $\deg(D) < 2d$. By the Riemann-Roch theorem, $\dim(h^0(D)) = 2d - 1$ and therefore, by Lemma 3.4, $K_n^M(E)$ is generated by elements of the form $\{f_1(e), \dots, f_n(e)\}$ where $f_i \in h^0(D)$. Let now D' be an effective divisor on Q of degree 2 (it exists since Q is a conic). Using again the Riemann-Roch theorem we see that $\dim(|e - D'|) > 0$, i.e. that there exists a rational function f with a simple pole in e and a zero in D' . In particular, the degrees of all the points where f has singularities other than e is strictly less than $2d$. Consider the symbol $\{f_1, \dots, f_n, f\} \in K_{n+1}^M(k(Q))$. Let

$$\partial : K_{n+1}^M(k(Q)) \rightarrow \bigoplus_{x \in Q_{(0)}} K_n^M(k_x)$$

be the residue homomorphism. By [9] its composition with (16) is zero. On the other hand we have

$$\partial(\{f_1, \dots, f_n, f\}) = \{f_1(e), \dots, f_n(e)\} + u$$

where u is a sum of symbols concentrated in the singular points of f_1, \dots, f_n and singular points of f other than e . Therefore, by our construction u belongs to $\bigoplus_{x \in Q_{(0)}, < 2d} K_n^M(k_x)$ and we conclude that $\text{tr}_{E/k}\{f_1(e), \dots, f_n(e)\}$ lies in the image of (17) by induction.

Let now Q be a quadric in \mathbf{P}^n where $n \geq 3$. Let c be a rational point of \mathbf{P}^n outside Q and $\pi : Q \rightarrow \mathbf{P}^{n-1}$ be the projection with the center in c . The ramification locus of π is a quadric on \mathbf{P}^{n-1} which has no rational points.

Assume first that there exists e such that the degree of $\pi(e)$ is d . Then, by Lemma 3.5, we can find a (singular) rational curve C' in \mathbf{P}^{n-1} of degree $d - 1$ which contains $\pi(e)$. Consider the curve $C = \pi^{-1}(C') \subset Q$. Let \tilde{C}, \tilde{C}' be the normalizations of C and C' and $\tilde{\pi} : \tilde{C} \rightarrow \tilde{C}'$ the morphism corresponding to π . Since $\deg(e) = 2d$ and $\deg(\pi(e)) = d$ the point e does not belong to the ramification locus of $\pi : Q \rightarrow \mathbf{P}^{n-1}$. This implies that e lifts to a point \tilde{e} of \tilde{C}

of degree $2d$ and that $\tilde{e} = \tilde{\pi}^{-1}(\tilde{\pi}(\tilde{e}))$. Since the ramification locus of π has no rational points the singular point of C' is unramified. This implies that $\tilde{\pi}$ is ramified in $\leq 2(d-1)$ points and, therefore, \tilde{C} is a hyperelliptic curve of genus less than or equal to $d-2$.

Let D be an effective divisor on \tilde{C} of degree $2d-2$. By the Riemann-Roch theorem we have $\dim(h^0(D)) \geq d+1$. On the other hand, since $\deg(D) < 2d$, the homomorphism $h^0(D) \rightarrow E$ defined by evaluation at \tilde{e} is injective. Therefore, by Lemma 3.4, $K_n^M(E)$ is additively generated by the elements of the form $\{f_1(\tilde{e}), \dots, f_n(\tilde{e})\}$ for $f_i \in h^0(D)$.

Let D' be an effective divisor on \tilde{C} of degree 2. By the Riemann-Roch theorem we have $\dim(h^0(\tilde{e}-D')) \geq d+1 > 0$. Therefore, there exists a rational function f , with simple pole in \tilde{e} , such that all its other singularities are located in points of degree $< 2d$. We can conclude now that $\text{tr}_{E/k}\{f_1(\tilde{e}), \dots, f_n(\tilde{e})\}$ belongs to the image of (17) in the same way as in the case of $\dim(Q) = 1$.

Consider now the general case - we still assume that $n \geq 3$ but not that we can find a center of projection c such that $\deg(\pi(e)) = d$. Taking a general c we may assume that $\deg(\pi(e)) = 2d$ and that e does not belong to the ramification locus of π . By Lemma 3.5 we can find a rational curve C' in \mathbf{P}^{n-1} of degree $2d-1$ which contains $\pi(e)$. Consider the curve $C = \pi^{-1}(C') \subset Q$. Let \tilde{C}, \tilde{C}' be the normalizations of C and C' , and $\tilde{\pi} : \tilde{C} \rightarrow \tilde{C}'$ the morphism corresponding to π . Since the point e does not belong to the ramification locus of π it lifts to a point \tilde{e} of \tilde{C} of degree $2d$. Since the ramification locus of π does not have rational points and the only singular point of C' is rational, $\tilde{\pi}$ is ramified in no more than $2(2d-1)$ points; therefore, \tilde{C} is a hyperelliptic curve of genus $\leq 2d-2$.

Let D be an effective divisor on $\tilde{C}' = \mathbf{P}^1$ of degree d . We have $\dim(h^0(D)) = d+1$ and since $\tilde{\pi}(\tilde{e})$ has degree $2d$ the evaluation at $\tilde{\pi}(\tilde{e})$ gives an injective homomorphism $h^0(D) \rightarrow E$. By Lemma 3.4, we conclude that any element of $K_n^M(E)$ is a linear combination of elements of the form $\{f_1(\tilde{\pi}(\tilde{e})), \dots, f_n(\tilde{\pi}(\tilde{e}))\}$ where f_i are in $h^0(D)$.

Let D' be an effective divisor of degree 2 on \tilde{C} . By the Riemann-Roch theorem we have $\dim(h^0(\tilde{e}-D')) \geq 1$; i.e., there exists a rational function f with a simple pole in \tilde{e} and a zero in D' . If $d > 1$ then all the singular points of f , except \tilde{e} , are of degree $< 2d$ and by the same reasoning as in the previous two cases we conclude that $\text{tr}_{E/k}(\{f_1(\tilde{\pi}(\tilde{e})), \dots, f_n(\tilde{\pi}(\tilde{e}))\})$ is a linear combination of the form

$$\sum_{x \in \tilde{C}_{(0), < 2d}} \text{tr}_{k_x/k}(u_x) + \sum_{i,y \in (f_i \circ \tilde{\pi})} \text{tr}_{k_y/k}(v_{i,y}).$$

Summands of the first type are in the image of (17) by the inductive assumption. The fact that summands of the second type are in the image of (17) follows from the case $\deg(\pi(e)) = d$ considered above.

4. Some applications

4.1. *Milnor's Conjecture on quadratic forms.* As the first corollary of Theorem 2.10 we get *Milnor's Conjecture on quadratic forms*.

As usual, we denote by $W(k)$ the *Witt ring* of quadratic forms over k , and by $I \subset W(k)$ the ideal of even-dimensional forms. The filtration $W(k) \supset I \supset I^2 \supset \dots \supset I^n \supset \dots$ by the powers of I is called the I -filtration on W . We denote the associated graded ring by $\text{Gr}_I^*(W(k))$. Consider the map

$$K_1^M(k)/2 = k^*/(k^*)^2 \xrightarrow{\varphi_1} \text{Gr}_I^1(W(k))$$

which sends $\{a\}$ to $\langle 1, -a \rangle$. Since $(\langle 1, -a \rangle + \langle 1, -b \rangle - \langle 1, -ab \rangle) \in I^2$ it is a group-homomorphism and one can easily see that it is an isomorphism. For any $a \in k^* \setminus 1$, the form $\langle\langle a, 1 - a \rangle\rangle$ is hyperbolic and, therefore, the isomorphism φ_1 can be extended to a ring homomorphism $\varphi : K_*^M(k)/2 \rightarrow \text{Gr}_I^*(W(k))$. Since $\text{Gr}_I^*(W(k))$ is generated by the first-degree component φ is surjective. The *Milnor Conjecture on quadratic forms* states that φ is an isomorphism i.e. that it is injective. It was proven in degree 2 by J. Milnor [6], in degree 3 by M. Rost [8] and A. Merkurjev-A. Suslin [7], and in degree 4 by M. Rost. Moreover, R. Elman and T. Y. Lam [2] proved that the map φ is injective on *pure symbols*.

THEOREM 4.1. *Let k be a field of characteristic zero. Then, the natural map $\varphi : K_*^M(k)/2 \rightarrow \text{Gr}_I^*(W(k))$ is an isomorphism.*

Proof. We already know that φ is surjective. Let $h \neq 0$ be an element of $K_n^M(k)/2$. By Theorem 2.10 there exists a field extension E/k such that $h|_E$ is a nonzero pure symbol. By a result of R. Elman and T. Y. Lam ([2]) the map φ is injective on *pure symbols*. Hence $\varphi(h|_E)$ is a nonzero element of $\text{Gr}_I^n(W(E))$. Since the morphism φ is compatible with field extensions, the element $\varphi(h) \in \text{Gr}_I^n(W(k))$ is also nonzero. Therefore, φ is injective. \square

4.2. *The Kahn-Rost-Sujatha Conjecture.* In [5] B. Kahn, M. Rost and R. Sujatha proved that for any quadric Q of dimension m the $\ker(K_i^M(k)/2 \rightarrow K_i^M(k(Q))/2)$ is trivial for any $i < \log_2(m+2)$, if $i \leq 4$ (actually, in [5] the authors worked with $H_{et}^i(k, \mathbf{Z}/2)$ instead of $K_i^M(k)/2$, but because of [13] we can use $K_i^M(k)/2$ here). The authors also conjectured¹ (among other things) that the same is true without the restriction $i \leq 4$. The following result proves this conjecture.

THEOREM 4.2. *Let Q be an m -dimensional quadric over a field k of characteristic zero. Then $\ker(K_i^M(k)/2 \rightarrow K_i^M(k(Q))/2)$ is trivial for any $i < \log_2(m+2)$.*

¹only in the original version of the paper

Proof. Denote by q a quadratic form which defines the quadric Q . Assume that h is a nonzero element of $\ker(\mathbb{K}_i^M(k)/2 \rightarrow \mathbb{K}_i^M(k(Q))/2)$. Using Theorem 2.10 we can find an extension E/k such that $h|_E$ is a nonzero pure symbol of the form $\underline{a} = \{a_1, \dots, a_n\}$. Then, since $h|_{E(Q)} = 0$, the corresponding Pfister quadric $Q_{\underline{a}}/E$ becomes hyperbolic over $E(Q)$. Since $Q_{\underline{a}}|_{E(Q)}$ is hyperbolic the form $t \cdot q|_E$ is isomorphic to a subform of the Pfister form $\langle\langle a_1, \dots, a_n \rangle\rangle$ for some coefficient $t \in E^*$ by [11, Ch. 4, Th. 5.4]. In particular, $m + 2 = \dim(Q) + 2 = \dim(q) \leq 2^i$. Therefore, $i \geq \log_2(m + 2)$. \square

4.3. *The J -filtration conjecture.* Together with the I -filtration on $W(k)$ we can consider the following so-called J -filtration. Let $x \in W(k)$ be an element, q an anisotropic quadratic form which represents x and Q the corresponding projective quadric. Since Q has a point over the field $k(Q)$, we have a decomposition of the form

$$q|_{k(Q)} = q_1 \perp \underbrace{\mathbf{H} \perp \dots \perp \mathbf{H}}_{i_1(q)},$$

where q_1 is an anisotropic form over $k(Q)$, and \mathbf{H} is the elementary hyperbolic form. The number $i_1(q)$ is called the first *higher Witt index* of q . In the same way we can decompose $q_1|_{k(Q)(Q_1)}$ etc., obtaining a sequence of quadratic forms q, q_1, \dots, q_{s-1} , where each q_i is an anisotropic form defined over $k(Q) \dots (Q_{i-1})$, and

$$q_{s-1}|_{k(Q)\dots(Q_{s-1})} = \underbrace{\mathbf{H} \perp \dots \perp \mathbf{H}}_{i_s(q)}$$

is a hyperbolic form. By [4, Th. 5.8] (see also [11, Ch. 4, Th. 5.4]), any quadratic form q' over a field E , such that $q'|_{E(Q')}$ is hyperbolic, is proportional to some Pfister form. This implies that the form q_{s-1} is proportional to an n -fold Pfister form $\langle\langle a_1, \dots, a_n \rangle\rangle$, where $\{a_1, \dots, a_n\} \in \mathbb{K}_n^M(k(Q) \dots (Q_{s-2}))/2$. This procedure defines, for any element $x \in W(k)$, a natural number n which we will call the degree of x .

Let us define $J_n(W(k))$ as the subset of $W(k)$ consisting of all elements of degree $\geq n$. It can be easily checked that $I^n \subseteq J_n$. It was conjectured in [4, Question 6.7] and in [11] that the J coincides with the I . The following theorem proves this conjecture.

THEOREM 4.3. $J_n = I^n$.

Proof. Let x be an element of $J_n(W(k))$ which is represented by a quadratic form q . As above we have a sequence of quadrics Q, Q_1, \dots, Q_{s-1} such that $q|_{k(Q)(Q_1)\dots(Q_{s-1})}$ is hyperbolic. This means that x goes to 0 under the natural map from $W(k)$ to $W(k(Q)(Q_1) \dots (Q_{s-1}))$.

All quadrics Q, Q_1, \dots, Q_{s-1} have dimensions $\geq 2^n - 2 > 2^{n-1} - 2$. By Theorem 4.2, for any $0 \leq i \leq n-1$, the kernel $\ker(\mathbb{K}_i^M(k)/2 \rightarrow \mathbb{K}_i^M(k(Q) \dots (Q_{s-1})))$

is trivial. Therefore, applying the Milnor conjecture (Theorem 4.1), we conclude that the map

$$\mathrm{Gr}_J^i(W(k)) \rightarrow \mathrm{Gr}_J^i(W(k(Q) \dots (Q_{s-1})))$$

is a monomorphism for all $0 \leq i \leq n - 1$. Therefore the map

$$W(k)/I^n(W(k)) \rightarrow W(k(Q) \dots (Q_{s-1}))/I^n(W(k(Q) \dots (Q_{s-1})))$$

is a monomorphism as well. Therefore, x belongs to $I^n(W(k))$. \square

STEKLOV MATHEMATICAL INSTITUTE, 8 GUBKINA ST., MOSCOW, RUSSIA
E-mail address: orlov@mi.ras.ru

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS OF THE
 RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
E-mail address: vishik@mccme.ru

INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ
E-mail address: vladimir@math.ias.edu

REFERENCES

- [1] H. BASS and J. TATE, The Milnor ring of a global field, *Lecture Notes in Math.* **342** (1973), 340–446.
- [2] R. ELMAN and T. Y. LAM, Pfister forms and K -theory of fields, *J. Algebra* **23** (1972), 181–213.
- [3] T. Y. LAM, *Algebraic Theory of Quadratic Forms*, Benjamin/Cummings Publ. Co., Inc., Reading, Mass., 1973.
- [4] M. KNEBUSCH, Generic splitting of quadratic forms, *Proc. London Math. Soc.* **33** (1976), 67–93.
- [5] B. KAHN, M. ROST, and R. J. SUJATHA, Unramified cohomology of quadrics. I, *Amer. J. Math.* **12** (1998), 841–891.
- [6] J. MILNOR, Algebraic K -theory and quadratic forms, *Invent. Math.* **9** (1969/1970), 318–344.
- [7] A. S. MERKURJEV and A. A. SUSLIN, The norm-residue homomorphism of degree 3 (in Russian), *Izv. Akad. Nauk SSSR* **54** (1990), 339–356; English translation: *Math. USSR Izv.* **36** (1991), 349–368.
- [8] M. ROST, Hilbert theorem 90 for K_3^M for degree-two extensions, preprint, Regensburg, 1986; <http://www.mathematik.uni-bielefeld.de/~rost/K3-86.html>.
- [9] ———, Some new results on the Chow-groups of quadrics, preprint, Regensburg, 1990; <http://www.mathematik.uni-bielefeld.de/~rost/chowquadr.html>.
- [10] ———, Chow groups with coefficients, *Doc. Math.* **1** (1996), 319–393.
- [11] W. SCHARLAU, *Quadratic and Hermitian Forms*, Springer-Verlag, New York, 1985.
- [12] V. VOEVODSKY, Reduced power operations in motivic cohomology, *Publ. Math. Inst. Hautes Études Sci.* **98** (2003), 1–57.
- [13] ———, Motivic cohomology with $\mathbf{Z}/2$ -coefficients, *Publ. Math. IHES* **98** (2003), 59–104.

(Received October 23, 2001)