

Lecture 3: Untilting

October 5, 2018

In this lecture, we let C^b denote a perfectoid field of characteristic p .

Warning 1. We will often use the superscript b to signal that an object under consideration “lives in characteristic p ”. In particular, declaring that C^b is a perfectoid field of characteristic p is not meant to signal that C^b is given as the tilt of a perfectoid field C . In fact, our emphasis is the opposite: we take the field C^b as given, and would like to understand all possible *untilts* of C^b . Recall that an untilt of C^b is defined to be a pair (K, ι) , where K is a perfectoid field and $\iota : C \simeq K^b$ is an (continuous) isomorphism.

Question 2. Let C^b be a perfectoid field of characteristic p . How can one classify the untilts of C^b ?

Remark 3. Let K be a perfectoid field of characteristic zero. Note that giving a continuous isomorphism $\iota : C^b \simeq K^b$ is equivalent to giving an isomorphism of valuation rings

$$\mathcal{O}_C^b \rightarrow \mathcal{O}_K^b = \varprojlim(\cdots \rightarrow \mathcal{O}_K/p\mathcal{O}_K \xrightarrow{\varphi} \mathcal{O}_K/p\mathcal{O}_K \xrightarrow{\varphi} \mathcal{O}_K/p\mathcal{O}_K).$$

We saw in the previous lecture that this induces an isomorphism of quotient rings $\iota_0 : \mathcal{O}_C^b/(\pi) \simeq \mathcal{O}_K/(\pi)$ for some element $\pi \in C^b$ satisfying $0 < |\pi|_{C^b} < 1$. Conversely, any such isomorphism ι_0 can be lifted to an isomorphism of valuation rings $\mathcal{O}_C^b \simeq \mathcal{O}_K^b$, since \mathcal{O}_C^b is isomorphic to the inverse limit

$$\cdots \rightarrow \mathcal{O}_C^b/(\pi) \xrightarrow{\varphi} \mathcal{O}_C^b/(\pi) \xrightarrow{\varphi} \mathcal{O}_C^b/(\pi).$$

We may therefore rephrase Question 2 as follows: how can we classify perfectoid fields K of characteristic zero equipped with an isomorphism $\mathcal{O}_K/(\pi) \simeq \mathcal{O}_C^b/(\pi)$?

For an untilt (K, ι) of C , let us abuse notation by writing $\sharp : C^b \rightarrow K$ for the composite map $C^b \xrightarrow{\iota} K^b \xrightarrow{\sharp} K$. This map does not need to be surjective. However, it is not too far off. We saw in the previous lecture that every element $x \in \mathcal{O}_K$ is congruent modulo p to an element in the image of the map \sharp : that is, we can find an element $c_0 \in \mathcal{O}_C^b$ satisfying $x = c_0^\sharp + x'p$, for some $x' \in \mathcal{O}_K$. Applying the same argument to x' , we obtain $x = c_0^\sharp + c_1^\sharp p + x''p^2$, for some $x'' \in \mathcal{O}_K$. Iterating this argument, we obtain a description of x as an infinite sum

$$x = c_0^\sharp + c_1^\sharp p + c_2^\sharp p^2 + c_3^\sharp p^3 + \cdots,$$

for some sequence of elements $c_0, c_1, c_2, \dots \in \mathcal{O}_C^b$; note that this infinite sum makes sense because the ring \mathcal{O}_K is p -adically complete. The decomposition above is not at all unique: generally an element $x \in \mathcal{O}_K$ can be decomposed as a sum $\sum_{n \geq 0} c_n^\sharp p^n$ in many different ways. For example, if K is algebraically closed, then any element $x \in \mathcal{O}_K$ can be written in the form c_0^\sharp by choosing a compatible sequence of p^n th roots of x ; in characteristic zero, these p^n th roots are not unique.

One virtue of working with expressions like $\sum_{n \geq 0} c_n^\sharp p^n$ is that they make sense simultaneously in *every* untilt K of C^b . Moreover, it is possible to give work out formulas for adding and multiplying these expressions which are independent of the choice of K . To make this idea precise, it will be convenient to review the theory of Witt vectors.

Notation 4. Let R be a perfect ring of characteristic p : that is, a commutative ring such that $p = 0$ in R and every element $x \in R$ has a unique p th root. We let $W(R)$ denote the ring of Witt vectors of R . Then $W(R)$ is characterized up to (unique) isomorphism by the following properties:

- (1) There is an isomorphism $W(R)/pW(R) \simeq R$.
- (2) The element p is not a zero-divisor in $W(R)$.
- (3) The ring $W(R)$ is p -adically complete.

Example 5. Let $R = \mathbf{F}_p$ be the finite field with p -elements. Then $W(R) \simeq \mathbf{Z}_p$ can be identified with the ring of p -adic integers.

Notation 6. For every element $x \in R$, we let $[x]$ denote its *Teichmüller representative* in $W(R)$. Then $[x]$ is uniquely determined by the following properties:

- The quotient map $W(R) \rightarrow W(R)/pW(R) \simeq R$ carries $[x]$ to x .
- The element $[x] \in W(R)$ admits a p^n th root, for every $n \geq 0$.

Concretely, one can construct the Teichmüller representative $[x]$ as the limit $\lim_{n \rightarrow \infty} \overline{(x^{1/p^n})}^{p^n}$, where $\overline{x^{1/p^n}}$ is any element of $W(R)$ representing the p^n th root $x^{1/p^n} \in R$. The construction of Teichmüller representatives determines a map

$$[\bullet] : R \rightarrow W(R)$$

which is multiplicative (that is, we have $[xy] = [x][y]$) but not additive.

Remark 7. Let R be a perfect ring of characteristic p and let x be an element of $W(R)$. Then x has some image $c_0 \in R$ under the quotient map $W(R) \rightarrow W(R)/pW(R) \simeq R$. The Teichmüller lift $[c_0]$ is then congruent to x modulo p , so we can write $x = [c_0] + x'p$ for some $x' \in W(R)$. Iterating this observation, we obtain an identity

$$x = [c_0] + [c_1]p + [c_2]p^2 + [c_3]p^3 + \cdots,$$

called the *Teichmüller expansion* of x . Note that, in contrast to the situation before, this expansion is unique: if

$$\sum [c_n]p^n = \sum [c'_n]p^n,$$

then an easy induction shows that $c_n = c'_n$ for each n .

Remark 8. Let R be a perfect ring of characteristic p . Then the ring of Witt vectors $W(R)$ can be characterized by a universal property:

- (*) For any p -adically complete ring A , reduction modulo p induces a bijection

$$\mathrm{Hom}(W(R), A) \rightarrow \mathrm{Hom}(R, A/pA).$$

In other words, every ring homomorphism $R \rightarrow A/pA$ can be lifted uniquely to a ring homomorphism $W(R) \rightarrow A$.

Let us now specialize to the situation of interest to us.

Construction 9. Let C^b be a perfectoid field of characteristic p and let \mathcal{O}_C^b be the valuation ring of C^b . Then \mathcal{O}_C^b is a perfect ring of characteristic p . We let $\mathbf{A}_{\mathrm{inf}}$ denote the ring of Witt vectors $W(\mathcal{O}_C^b)$.

Remark 10. The ring $\mathbf{A}_{\mathrm{inf}}$ is one of Fontaine's *period rings*; it will play an essential role in this course.

Remark 11. Let C^b be a perfectoid field of characteristic p and let (K, ι) be an untilt of C^b . The map $\sharp : \mathcal{O}_C^b \rightarrow \mathcal{O}_K$ is not a ring homomorphism (unless K has characteristic p). However, it induces a ring homomorphism $\mathcal{O}_C^b \rightarrow \mathcal{O}_K/p\mathcal{O}_K$. Since \mathcal{O}_K is p -adically complete, the universal property of Remark 8 to lifts this to a ring homomorphism

$$\theta : \mathbf{A}_{\text{inf}} = W(\mathcal{O}_C^b) \rightarrow \mathcal{O}_K.$$

Concretely, this map is given by the formula

$$\theta([c_0] + [c_1]p + [c_2]p^2 + \dots) = c_0^\sharp + c_1^\sharp p + c_2^\sharp p^2 + \dots.$$

From the discussion at the beginning of the lecture, we deduce that θ is surjective.

Remark 12. In the situation of Remark 11, the map θ is *local*: that is, an element $\sum [c_n]p^n$ of \mathbf{A}_{inf} is invertible if and only if its image $\sum c_n^\sharp p^n \in \mathcal{O}_K$ is invertible (in both cases, invertibility is equivalent to the requirement that $|c_0|_{C^b} = 1$).

Remark 13. Let C^b be a perfectoid field of characteristic p and let K be a characteristic zero untilt of C^b . We saw in the previous lecture that it is possible to choose an element $\pi \in \mathcal{O}_C^b$ with $|\pi|_{C^b} = |p|_K$, and that the map $\sharp : C^b \rightarrow K$ induces an isomorphism of commutative rings $\mathcal{O}_C^b/\pi\mathcal{O}_C^b \simeq \mathcal{O}_K/p\mathcal{O}_K$. In other words, \mathcal{O}_C^b and \mathcal{O}_K have a common quotient ring. Remark 11 shows that both \mathcal{O}_C^b and \mathcal{O}_K can be realized as a quotient of the same ring $\mathbf{A}_{\text{inf}} = W(\mathcal{O}_C^b)$: one gets \mathcal{O}_C^b from \mathbf{A}_{inf} by reducing modulo p , and \mathcal{O}_K by reducing modulo the kernel $\ker(\theta)$. These quotient maps fit into a commutative diagram

$$\begin{array}{ccc} \mathbf{A}_{\text{inf}} & \xrightarrow{\theta} & \mathcal{O}_K \\ \downarrow & & \downarrow \\ \mathcal{O}_C^b & \xrightarrow{\sharp} & \mathcal{O}_K/p\mathcal{O}_K. \end{array}$$

In the situation of Remark 11, the map θ is never injective: in other words, it is always possible to write an element $x \in \mathcal{O}_K$ as a sum $\sum_{n \geq 0} c_n^\sharp p^n$ in multiple ways.

Example 14. Let C^b be a perfectoid field of characteristic p and let K be an untilt of C^b . We saw in the previous lecture that there exists an element $\pi \in \mathcal{O}_C^b$ satisfying $|\pi|_{C^b} = |p|_K$ (if K has characteristic p , we just take $\pi = 0$). It follows that the elements p and π^\sharp have the same absolute value in K , and therefore differ by multiplication by an invertible element $\bar{u} \in \mathcal{O}_K$. Write $\bar{u} = \theta(u)$ for $u \in \mathbf{A}_{\text{inf}}$ (so that u is also invertible, by Remark 12). The identity $\pi^\sharp = \bar{u}p$ then implies that $[\pi] - up \in \mathbf{A}_{\text{inf}}$ belongs to the kernel of θ .

Definition 15. Let C^b be a perfectoid field of characteristic p . We say that an element $\xi \in \mathbf{A}_{\text{inf}}$ is *distinguished* if it has the form $[\pi] - up$, where $|\pi|_{C^b} < 1$ and u is an invertible element of \mathbf{A}_{inf} . In other words, ξ is distinguished if its Teichmüller expansion

$$\xi = [c_0] + [c_1]p + [c_2]p^2 + \dots$$

has the property that $|c_0|_{C^b} < 1$ and $|c_1|_{C^b} = 1$.

Example 14 shows that, for every untilt K of C^b , the kernel of the induced map $\theta : \mathbf{A}_{\text{inf}} = W(\mathcal{O}_C^b) \rightarrow \mathcal{O}_K$ always contains a distinguished element ξ . We now establish a converse:

Proposition 16. *Let C^b be a perfectoid field of characteristic p and let ξ be a distinguished element of $\mathbf{A}_{\text{inf}} = W(\mathcal{O}_C^b)$. Then the quotient $\mathbf{A}_{\text{inf}}/(\xi)$ can be identified with the valuation ring \mathcal{O}_K in a perfectoid field K . Moreover, the canonical map*

$$\mathcal{O}_C^b = \mathbf{A}_{\text{inf}}/(p) \rightarrow \mathbf{A}_{\text{inf}}/(\xi, p) \simeq \mathcal{O}_K/(p)$$

exhibits K as an untilt of C^b (see Remark 3).

Corollary 17. *Let C^b be a perfectoid field of characteristic p , let K be an untilt of C^b , and let $\theta : \mathbf{A}_{\text{inf}} \rightarrow \mathcal{O}_K$ be as above. Then $\ker(\theta)$ is a principal ideal, generated by any choice of distinguished element $\xi \in \ker(\theta)$.*

Proof. It follows from Example 14 that $\ker(\theta)$ contains a distinguished element ξ , so that θ induces a surjection $\bar{\theta} : \mathbf{A}_{\text{inf}}/(\xi) \twoheadrightarrow \mathcal{O}_K$. Proposition 16 shows that we can identify $\mathbf{A}_{\text{inf}}/(\xi)$ with $\mathcal{O}_{K'}$ for some untilt K' of C^b . Since $\mathcal{O}_{K'}$ is an integral domain, the kernel of θ is a prime ideal of $\mathcal{O}_{K'}$, and is therefore either (0) (in which case $\bar{\theta}$ is an isomorphism) or the maximal ideal $\mathfrak{m}_{K'}$ (which is impossible, since \mathcal{O}_K is not a field). \square

Corollary 18. *Let C^b be a perfectoid field of characteristic p . Then the construction*

$$\xi \mapsto \text{Fraction field of } \mathbf{A}_{\text{inf}}/(\xi)$$

induces a bijection

$$\{\text{Distinguished elements of } \mathbf{A}_{\text{inf}}\}/\text{multiplication by units} \simeq \{\text{Untilts of } C^b\}/\text{isomorphism}.$$

To prove Proposition 16, we will need the following purely algebraic fact, which we leave to the reader.

Exercise 19. Let R be a commutative ring (not necessarily Noetherian!) containing a pair of elements x and y . Suppose that:

- The element x is not a zero-divisor in R , and R is x -adically complete.
- The image of y is not a zero-divisor in R/xR , and R/xR is y -adically complete.

Show that:

- The element y is not a zero-divisor in R , and R is y -adically complete.
- The image of x is not a zero-divisor in R/yR , and R/yR is x -adically complete.

Proof of Proposition 16. Let ξ be a distinguished element of \mathbf{A}_{inf} , so that we can write $\xi = [\pi] - up$ for some $\pi \in \mathfrak{m}_C^b$ and some invertible element u in \mathbf{A}_{inf} . If $\pi = 0$, then $\mathbf{A}_{\text{inf}}/(\xi) \simeq \mathbf{A}_{\text{inf}}/(p) \simeq \mathcal{O}_C^b$ and we have nothing to prove. Let us therefore assume that π is not zero. Let \mathcal{O}_K denote the quotient ring $\mathbf{A}_{\text{inf}}/\xi\mathbf{A}_{\text{inf}}$. (Beware that this notation is misleading, since we do not yet know that there is a valued field K having \mathcal{O}_K as its valuation ring.) We then have a canonical map $\theta : \mathbf{A}_{\text{inf}} \twoheadrightarrow \mathcal{O}_K$ (with kernel generated by ξ); for each element $x \in \mathcal{O}_C^b$, we will denote $\theta([x])$ by $x^\sharp \in \mathcal{O}_K$.

We now apply Exercise 19 to the elements $x = p$ and $y = \xi$ of the ring \mathbf{A}_{inf} . The construction of \mathbf{A}_{inf} as the ring of Witt vectors $W(\mathcal{O}_C^b)$ shows that \mathbf{A}_{inf} is p -adically complete and p -torsion free. Moreover, the image of ξ in the quotient $\mathbf{A}_{\text{inf}}/p\mathbf{A}_{\text{inf}} \simeq \mathcal{O}_C^b$ is π , satisfying $0 < |\pi|_{C^b} < 1$. It follows that \mathcal{O}_C^b is ξ -adically complete and ξ -torsion free. Applying Exercise 19, we deduce the following:

- The ring \mathbf{A}_{inf} is ξ -adically complete and ξ -torsion free.
- The quotient ring $\mathcal{O}_K = \mathbf{A}_{\text{inf}}/(\xi)$ is p -adically complete and p -torsion free.

We next prove the following:

(a) For any element $y \in \mathcal{O}_K$, there exists an element $x \in \mathcal{O}_C^b$ such that y is a unit multiple of x^\sharp .

To prove (a), we may assume without loss of generality that $y \neq 0$. Since \mathcal{O}_K is p -adically complete, we can write $y = p^n y'$ for some $y' \in \mathcal{O}_K$ which is not divisible by p . Replacing y by y' , we may assume that y is not divisible by p . Since θ is surjective, we can choose $x \in \mathcal{O}_C^b$ such that $y \equiv x^\sharp \pmod{p}$. Then x^\sharp is not divisible by p , so x is not divisible by π . We can therefore write $\pi = xx'$ for some $x' \in \mathfrak{m}_C^b$. We have $y = x^\sharp + \pi^\sharp w = x^\sharp(1 + x'^\sharp w)$ for some $w \in \mathcal{O}_K$. Since some power of x' is divisible by π in the ring \mathcal{O}_C^b , some power of x'^\sharp is divisible by p in the ring \mathcal{O}_K . It follows that $1 + x'^\sharp w$ is an invertible element of \mathcal{O}_K (with

inverse given by the p -adically convergent sum $1 - x'^{\sharp}w + (x'^{\sharp}w)^2 - (x'^{\sharp}w)^3 + \dots$. This proves that y is a unit multiple of x^{\sharp} , as desired.

Note that the element x appearing in (a) is not uniquely determined: we are free to multiply it by any unit in \mathcal{O}_C^b . However, this is our only freedom:

- (b) Let $x, x' \in \mathcal{O}_C^b$ be elements such that x^{\sharp} is divisible by x'^{\sharp} in \mathcal{O}_K . Then x is divisible by x' in \mathcal{O}_C^b : that is, we have $|x|_{C^b} \leq |x'|_{C^b}$.

Suppose otherwise. We then have $|x|_{C^b} > |x'|_{C^b}$, so we can write $x' = tx$ for some $t \in \mathfrak{m}_{C^b}$. Since x is not zero, it divides π^n for $n \gg 0$. Consequently, our assumption that x^{\sharp} is a multiple of x'^{\sharp} guarantees that $(\pi^n)^{\sharp}$ is a unit multiple of $(\pi^n t)^{\sharp}$ in \mathcal{O}_K . Since π^{\sharp} is a unit multiple of p and \mathcal{O}_K is p -torsion-free, it follows that t^{\sharp} is a unit in \mathcal{O}_K . This is impossible, since the image of t^{\sharp} is nilpotent in the ring $\mathcal{O}_K/p\mathcal{O}_K \simeq \mathcal{O}_C^b/\pi\mathcal{O}_C^b$.

We next claim:

- (c) The ring \mathcal{O}_K is an integral domain.

To prove (c), let y be any nonzero element of \mathcal{O}_K ; we wish to show that y is not a zero divisor. By virtue of (a), we may assume that $y = x^{\sharp}$ for some nonzero element $x \in \mathcal{O}_C^b$. Then x divides π^n for some large n ; we may therefore replace x by π^n . In this case, y is a unit multiple of p^n , which we have already seen is not a zero divisor in \mathcal{O}_K .

For each element $y \in \mathcal{O}_K$, let us define $|y|_K = |x|_{C^b}$, where x is any element of \mathcal{O}_C^b satisfying $y = x^{\sharp} \cdot \text{unit}$. It follows from (a) and (b) that $|y|_K$ is well-defined. Moreover, we have obvious identities

$$|0|_K = 0 \quad |1|_K = 1 \quad |y \cdot z|_K = |y|_K \cdot |z|_K.$$

Moreover, it follows from (b) that for each $y, z \in \mathcal{O}_K$, we have $|y|_K \leq |z|_K$ if and only if y is divisible by z . This immediately implies that $|y + z|_K \leq \max(|y|_K, |z|_K)$. We can therefore extend $|\bullet|_K$ uniquely to a non-archimedean absolute value on the fraction field K of \mathcal{O}_K . Moreover, an element $\frac{y}{z}$ of K satisfies $|\frac{y}{z}|_K = \frac{|y|_K}{|z|_K} \leq 1$ if and only if $|y|_K \leq |z|_K$: that is, if and only if y is divisible by $z \in \mathcal{O}_K$. It follows that \mathcal{O}_K is the valuation ring of K (with respect to the absolute value $|\bullet|_K$).

Note that $|p|_K = |\pi|_{C^b} < 1$, so that K has residue characteristic p . Moreover, since p is not a zero-divisor in \mathcal{O}_K , the field K has characteristic zero: that is, p is a pseudo-uniformizer of \mathcal{O}_K . Consequently, the assertion that \mathcal{O}_K is p -adically complete guarantees that it is complete with respect to its absolute value. The maximal ideal $\mathfrak{m} \subseteq \mathcal{O}_K$ is not generated by p : for example, it contains the element $(\pi^{1/p})^{\sharp}$, which is not divisible by p . Finally, we note that the isomorphisms

$$\mathcal{O}_K/p\mathcal{O}_K \simeq \mathbf{A}_{\text{inf}}/(\xi, p) \simeq (\mathbf{A}_{\text{inf}}/p\mathbf{A}_{\text{inf}})/(\xi) = \mathcal{O}_C^b/\pi\mathcal{O}_C^b$$

guarantee that the Frobenius map is surjective on $\mathcal{O}_K/p\mathcal{O}_K$. Consequently, K is a perfectoid field of characteristic zero which is an untilt of C (Remark 3). \square